

THE FINITE FRIEDMAN-STANLEY JUMPS: GENERIC DICHOTOMIES FOR BOREL HOMOMORPHISMS

ASSAF SHANI

ABSTRACT. Fix $n = 1, 2, 3, \dots$ or $n = \omega$. We prove a dichotomy for Borel homomorphisms from the n -th Friedman-Stanley jump $=^{+n}$ to an equivalence relation E which is classifiable by countable structures: if there is no reduction from $=^{+n}$ to E , then in fact all Borel homomorphisms are very far from a reduction. For this we use a different presentation of $=^{+n}$, equivalent up to Borel bi-reducibility, which is susceptible to Baire-category techniques.

This dichotomy is seen as a method for proving positive Borel reducibility results from $=^{+n}$. As corollaries we prove: (1) for $n \leq \omega$, $=^{+n}$ is in the spectrum of the meager ideal. This extends a result of Kanovei, Sabok, and Zapletal for $n = 1$; (2) $=^{+\omega}$ is a regular equivalence relation. This answers positively a question of Clemens; (3) for $n < \omega$, the equivalence relations, classifiable by countable structures, which do not Borel reduce $=^{+n}$ are closed under countable products. This extends a result of Kanovei, Sabok, and Zapletal for $n = 1$.

1. INTRODUCTION

This paper is a contribution to the study of equivalence relations on Polish spaces up to Borel reducibility. Given equivalence relations E and F on Polish spaces X and Y respectively, a map $f: X \rightarrow Y$ is said to be a **reduction** of E to F if for any $x_1, x_2 \in X$,

$$x_1 E x_2 \iff f(x_1) F f(x_2).$$

We say that E is **Borel reducible** to F , denoted $E \leq_B F$ if there is a Borel measurable function which is a reduction of E to F . In this case, we think of E as no more complicated than F . Borel reducibility is the most central concept in the study of equivalence relations on Polish spaces. Say that E and F are **Borel bireducible**, denoted $E \sim_B F$, if $E \leq_B F$ and $F \leq_B E$. An equivalence relation E on a Polish space X is **Borel** if E is a Borel subset of $X \times X$, with the product topology. More generally, E is **analytic** if E is an analytic subset of $X \times X$, that is, E is the projection of a Borel subset of $X \times X \times Y$ for some Polish space Y .

Date: November 22, 2024.

2010 Mathematics Subject Classification. Primary: 03E15, 03E75, 54H05, 03E25, 03E47.

Key words and phrases. Borel reducibility, Analytic equivalence relations, Borel homomorphisms, Friedman-Stanley jump, Hereditarily countable sets.

Research partially supported by NSF grant DMS-2246746 and NSERC grant RGPIN-2024-05827.

A central motivation for the field is to study the complexity of various classification problems in mathematics. Generally speaking, separable mathematical objects can be coded as members of some Polish space. Natural notions of equivalence, such as isomorphism between countable graphs, isometry between separable metric spaces, or homeomorphism between compact metric spaces, can then be seen as equivalence relations on Polish spaces. These are generally analytic, and sometimes Borel. Another point of view is the study of (Borel) definable cardinality between quotients of Polish spaces. A reduction of E to F corresponds to an injective map between quotient spaces X/E to Y/F . So we study injective maps between such quotient spaces, but only consider “sufficiently nice” maps, those which lift to a Borel map between Polish spaces.

Given an equivalence relation E on X , the **Friedman-Stanley jump** of E is the equivalence relation E^+ on $X^{\mathbb{N}}$ defined by

$$x E^+ y \iff \forall n \exists m (x(n) E y(n)) \text{ and } \forall n \exists m (y(n) E x(m)),$$

equivalently, if $\{[x(n)]_E : n \in \mathbb{N}\} = \{[y(n)]_E : n \in \mathbb{N}\}$. The quotient $E^+/X^{\mathbb{N}}$ may be identified with $\mathcal{P}_{\aleph_0}(E/X)$, the countable powerset of E/X .

The iterated Friedman-Stanley jumps, $=^{+\alpha}$, are defined recursively along the countable ordinals as follows (see [Gao09, 12.2.6]).

- $=^{+0}$ is the equality relation on \mathbb{R} , $=_{\mathbb{R}}$,
- $=^{+(\alpha+1)}$ is defined as $(=^{+\alpha})^+$,
- $=^{+\lambda}$ is defined as $\prod_{\alpha < \lambda} =^{+\alpha}$, for a limit ordinal λ .

The equivalence relation $=^{+1}$ is often denoted as $=^+$. The Friedman-Stanley jumps play a central role in the theory of equivalence relations. A classification problem is considered “classifiable using countable sets of reals as complete invariants” if it is Borel reducible to $=^+$; “classifiable using countable sets of countable sets of reals as complete invariants” if it is Borel reducible to $=^{+2}$; and so on.

An equivalence relation is **classifiable by countable structures** if it is Borel reducible to an isomorphism relation on a space of all countable \mathcal{L} -structures, for some countable language \mathcal{L} . (See [Kan08, 12.3], [Gao09, 3.6], [Hjo00]). A Borel equivalence relation which is classifiable by countable structures is Borel reducible to $=^{+\alpha}$ for some countable ordinal α (see [Fri00, Theorem 1.5]).

When studying some equivalence relation E , we would like to compare it, in terms of Borel reducibility, to a given Friedman-Stanley jump $=^{+\alpha}$. The results in [HKL98] provide a powerful tool for proving that E is Borel reducible to $=^{+\alpha}$. There are flexible tools to prove *irreducibility* results between some E and $=^{+\alpha}$, such as the study of pinned cardinals [LZ20, URL17] and the use of symmetric models in [Sha21].

Problem 1.1. Fix a countable ordinal α . Develop tools to construct a Borel reduction from $=^{+\alpha}$ to some other equivalence relation.

In this paper we provide such tools for $\alpha \leq \omega$. First, we note that Problem 1.1 is well understood for $=^+$, that is, $\alpha = 1$. There are many results reducing $=^+$

to other equivalence relations¹, for example, [For00, Theorem 65 part 2], [Kay17, Theorem 1.1], and [CMRS23, Proposition 3.5]. There are also three general results for constructing such a reduction:

- Marker [Mar07, Theorem 1.2] provides a model theoretic criterion for a first order isomorphism relation \cong_T to reduce $=^+$: if the type space $S(T)$ is uncountable.
- Larson and Zapletal [LZ20, Theorem 2.8.11] provide a set theoretic criterion for an analytic equivalence relation E to reduce $=^+$: if E is unpinned in the Solovay extension.
- Kanovei, Sabok, and Zapletal provided the following Baire-category tool.

Theorem 1.2 (Kanovei-Sabok-Zapletal [KSZ13, Theorem 6.24]). Let E be an analytic equivalence relation. Then either

- $=^+$ is Borel reducible to E , or
- any Borel homomorphism from $=^+$ to E maps a comeager set into a single E -class.

Given equivalence relations E and F on Polish spaces X and Y , a map $f: X \rightarrow Y$ is a **Borel homomorphism** from E to F , denoted $f: E \rightarrow_B F$, if for any $x_1, x_2 \in X$,

$$x_1 E x_2 \implies f(x_1) F f(x_2).$$

Theorem 1.2 says that if there is no Borel reduction of $=^+$ to E , then in fact all Borel homomorphisms from $=^+$ to E are trivial, on a comeager set.

We mention two immediate difficulties in generalizing this to the higher jumps.

Remark 1.3. For $n \geq 2$, $=^{+n}$ does not behave well, in terms of Baire-category, with the product topology given by the Friedman-Stanley jump. See Claim 1.22 below.

Remark 1.4. For $1 \leq k < n$, there is a natural Borel homomorphism from $=^{+n}$ to $=^{+k}$, which is not “completely trivial”. For example, the homomorphism $u: (\mathbb{R}^{\mathbb{N}})^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ from $=^{+2}$ to $=^+$, defined by $u(x)((n, m)) = x(n)(m)$, where $(,) : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ is a bijection. This is the “union homomorphism”, taking a set of sets of reals $\{\{x(n)(m) : m \in \mathbb{N}\} : n \in \mathbb{N}\}$ to an enumeration of their union $\{x(n)(m) : n, m \in \mathbb{N}\}$.

With these two modifications in mind, we provide a complete Baire-category analysis of all Borel homomorphisms from $=^{+n}$, $n \leq \omega$, to equivalence relations which are classifiable by countable structures. This is the main result of the paper.

Theorem 1.5. There are Borel equivalence relations F_n for $n \leq \omega$, and Borel homomorphisms $u_k^n : F_n \rightarrow_B F_k$ for $1 \leq k \leq n \leq \omega$, so that for each $n \leq \omega$, F_n and $=^{+n}$ are Borel bireducible, and the following dichotomy for Borel homomorphisms holds. For any equivalence relation E which is classifiable by countable structures, either

¹In the literature, $=^+$ takes many names, including Eq^+ , E_{ctbl} , F_2 , and \cong_2 .

- F_n is Borel reducible to E , or
- for any Borel homomorphism $f: F_n \rightarrow_B E$ there is some $k < n$ so that the homomorphism f factors through u_k^n on a comeager set, that is, there is a Borel homomorphism $h: F_k \rightarrow_B E$, defined on a comeager set, so that $h \circ u_k^n(x) \in E$ for a comeager set of x in the domain of F_n . (See Figure 1.)




FIGURE 1. $(\forall f: F_n \rightarrow_B E)(\exists k < n \exists h: F_k \rightarrow_B E)$

Remark 1.6. As remarked above the equivalence relations F_n , $n > 1$, are necessarily different than $=^{+n}$. For $n = 1$, the equivalence relation F_1 is simply $=^+$. Let F_0 be the trivial equivalence relation on a space $\{*\}$ with a single element, and consider the trivial homomorphism $u_0^1: F_1 \rightarrow_B F_0$. Then the second bullet of Theorem 1.2 for $=^+$ has the same form as Figure 1 with $k = 0$ and $n = 1$.

Remark 1.7. We see Theorem 1.5 as a tool to prove that $=^{+n}$ is Borel reducible to some equivalence relation E . In order to prove that such reduction exists, it suffices to find a Borel homomorphism which is “sufficiently different” from the homomorphisms u_k^n , for $k < n$.

The definition of the equivalence relations F_n , appearing in Theorem 1.5, is given in Section 1.1. A group action inducing F_n is presented in Section 1.2. We then prove the following corollaries of Theorem 1.5. The definitions and background are presented in each subsection.

- (1) (Section 1.3.) For $n \leq \omega$, $=^{+n}$ is in the spectrum of the meager ideal. This was proved for $n = 1$ in [KSZ13].
- (2) (Section 1.4) $=^{+\omega}$ is regular. This answers positively a question of Clemens [Cle22].
- (3) (Section 1.5) Fix $n < \omega$. Suppose G_k , $k \in \mathbb{N}$, are classifiable by countable structures and $=^{+n} \not\leq_B G_k$. Then $=^{+n} \not\leq_B \prod_{k \in \mathbb{N}} G_k$. This was proved for $n = 1$ in [KSZ13].

Several open questions related to these results are posed in the relevant subsections. In Section 1.6 we note that the Borel complexity of F_n is $\mathbf{\Pi}_{2+n}^0$, which is the optimal potential complexity of $=^{+n}$ by [HKL98].

We first focus on proving a corollary of Theorem 1.5, that F_n preserves its complexity on comeager sets (see Section 1.3), which is proved in Section 5 (Theorem 5.1). In Section 2 we sketch some ideas from [KSZ13], for proving that $=^+$ retains its complexity on comeager sets, and explain the main difficulties towards $n \geq 2$. The main construction, which will eventually lead to the necessary reductions of F_n , is

presented in Section 4. In Section 3 we present some technical results regarding Vaught transforms for the actions presented in Section 1.2.

In Section 6 we present some ideas behind the proof of Theorem 1.2 from [KSZ13], and explain the remaining difficulties towards extending these to the $n \geq 2$ case. In particular, we will use the following lemma.

Lemma 1.8. Let E be an equivalence relation which is classifiable by countable structures and let $f: F_n \rightarrow_B E$ be a Borel homomorphism which does not factor through u_k^n , $k < n$, on a comeager set. Then there are equivalence relations E_k , for $k < n$, Borel homomorphisms $\pi_k^n: E \rightarrow_B E_k$, $\pi_k^{k+1}: E_{k+1} \rightarrow_B E_k$, and $f_k: F_k \rightarrow_B E_k$ so that the following diagram commutes on comeager sets, and so that f_k does not factor through u_l^k for $l < k$.

$$\begin{array}{ccccccc}
 F_1 & \xleftarrow{u_1^2} & F_2 & \xleftarrow{u_2^3} & F_3 & \xleftarrow{u_3^4} & \cdots & \xleftarrow{u_k^n} & F_n \\
 \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & & & \downarrow f \\
 E_1 & \xleftarrow{\pi_1^2} & E_2 & \xleftarrow{\pi_2^3} & E_3 & \xleftarrow{\pi_3^4} & \cdots & \xleftarrow{\pi_k^n} & E
 \end{array}$$

The proof of Theorem 1.5 is then completed in Section 7.

Remark 1.9. The proof of Lemma 1.8 is the only place in which we use that E is classifiable by countable structures. Extending the lemma for a wider class of equivalence relations will similarly extend Theorem 1.5, as well as Theorem 1.29 and Proposition 1.32 below.

Question 1.10. Is Lemma 1.8 true for all analytic equivalence relations?

Problem 1.11. Find a model theoretic condition for an isomorphism relation \cong_T to reduce $=^{+n}$, extending the result [Mar07, Theorem 1.2] for $n = 1$.

Problem 1.12. Find a set theoretic condition for an equivalence relation E to reduce $=^{+n}$, extending the result [LZ20, Theorem 2.8.11] for $n = 1$.

1.1. The definition of F_n and u_k^n from Theorem 1.5. Consider the Polish space $((2^{\mathbb{N}})^{\mathbb{N}})^{\omega}$, with the natural product topology. We use the following standard notation: the space 2 is identified with the discrete space with two elements $\{0, 1\}$. The ordinal ω is identified with the set of natural numbers $\mathbb{N} = \{0, 1, 2, \dots\}$. The space $2^{\mathbb{N}}$ is identified with the space $\mathcal{P}(\mathbb{N})$ of all subsets of \mathbb{N} .

Given $x \in ((2^{\mathbb{N}})^{\mathbb{N}})^{\omega}$, we define a sequence A_n^x , $n = 1, 2, \dots$, as follows.

- $A_1^x = \{x(0)(k) : k \in \mathbb{N}\}$.
- For $l \in \mathbb{N}$, define $a_1^{x,l} = \{x(0)(k) : x(1)(l)(k) = 1\}$, a subset of A_1^x .

Given A_n^x and $a_n^{x,l}$ for $l \in \mathbb{N}$, define

- $A_{n+1}^x = \{a_n^{x,k} : k \in \mathbb{N}\}$, and
- $a_{n+1}^{x,l} = \{a_n^{x,k} : x(n+1)(l)(k) = 1\}$, a subset of A_{n+1}^x .

For $m < \omega$, and $x \in ((2^{\mathbb{N}})^{\mathbb{N}})^m$ we define similarly A_n^x for $n = 1, 2, \dots, m$.

Definition 1.13. For $2 \leq n \leq \omega$, define $X_n \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^n$ as the set of all x such that:

- (1) $(\forall 1 \leq i < n)(\forall m)(\exists k)x(i)(k)(m) = 1$;
- (2) $(\forall 1 \leq i < n)(\forall k)(\exists m)x(i)(k)(m) = 1$;
- (3) $(\forall 1 \leq i < n)(\forall k, l_1, l_2)(x(i-1)(l_1) = x(i-1)(l_2) \rightarrow x(i)(k)(l_1) = x(i)(k)(l_2))$.

Observation 1.14. From condition (1) it follows that for any $n \leq \omega$,

- If $x \in X_n$ then $A_k^x = \bigcup A_{k+1}^x$ for any $k < n$.
- For $m < \omega$, $m \leq n$, $x, y \in X_n$, if $A_m^x = A_m^y$ then $A_k^x = A_k^y$ for all $k < m$.

Observation 1.15. In the construction above, we used the binary sequence $x(i+1)(t) \in 2^{\mathbb{N}}$ to code a subset of A_{i+1}^x , via its enumeration $(a_i^{x,k} : k \in \mathbb{N})$. Condition (3) says that for $x \in X_n$, if l_1 and l_2 are identified in this enumeration, $a_i^{x,l_1} = a_i^{x,l_2}$, then they are also identified by $x(i+1)(t)$. Condition (2) says that for $x \in X_n$, for each $i < n$, $a_{i+1}^{x,l}$ is a non-empty subset of A_{i+1}^x .

Remark 1.16. Note that X_n is a dense G_δ subset of $((2^{\mathbb{N}})^{\mathbb{N}})^n$. For condition (3) this is true since for a dense G_δ set of $x \in ((2^{\mathbb{N}})^{\mathbb{N}})^n$, $x(i-1)(l_1) \neq x(i-1)(l_2)$, for $l_1 \neq l_2$.

Definition 1.17 (Main definition: F_n and u_k^n).

(1) For $n < \omega$, the equivalence relation F_n is defined on X_n by

$$x F_n y \iff A_n^x = A_n^y.$$

The equivalence relation F_ω is defined on X_ω by

$$x F_\omega y \iff \forall n < \omega (A_n^x = A_n^y).$$

(2) Given $m \leq n$, define $u_m^n : X_n \rightarrow X_m$ as the natural projection map to the first m copies of $(2^{\mathbb{N}})^{\mathbb{N}}$. Then $u_m^n : F_n \rightarrow_B F_m$ is a Borel homomorphism. For $n < \omega$, the homomorphism u_{n-1}^n can be seen as the union map, sending the set A_n^x to its union $A_{n-1}^x = A_{n-1}^{u_{n-1}^n(x)}$.

The map $x \mapsto A_n^x$ can be seen as a reduction of F_n to $=^{+n}$. Specifically, there is a Borel map from X_n to $(2^{\mathbb{N}})^{\omega^n}$ (the domain of $=^{+n}$), sending each $x \in X_n$ to some $z \in (2^{\mathbb{N}})^{\omega^n}$ “enumerating” the set A_n^x .

1.2. Group action. In this section we present the equivalence relations F_n as orbit equivalence relation, when restricted to a (large) subdomain.²

Consider the Polish group S_∞ of all permutations of \mathbb{N} , with its natural action $a : S_\infty \curvearrowright (2^{\mathbb{N}})^{\mathbb{N}}$, permuting the sequence of reals. The induced orbit equivalence relation on $(2^{\mathbb{N}})^{\mathbb{N}}$ is not $=^+$, but is Borel bireducible with $=^+$ (see [Gao09, Exercise

²In this section, and throughout the paper, we use colors for emphasis and clarification. The reader is advised to view these pages in color.

8.3.4]). The two equivalence relations in fact agree on the comeager set of all injective sequences of reals.

Consider also the natural action $b_0: S_\infty \curvearrowright 2^\mathbb{N}$, permuting binary sequence. Let $b: S_\infty \curvearrowright (2^\mathbb{N})^\mathbb{N}$ be the diagonal action, $g \cdot_b (x_n)_n = (g \cdot_{b_0} x_n)_n$. Recall the definition of F_2 on $(2^\mathbb{N})^\mathbb{N} \times (2^\mathbb{N})^\mathbb{N}$. When permuting a sequence of reals (the first coordinate) using a , the action of b on the second coordinate updates the binary sequences, so that they still carve out the same subset of reals as before. To recover all the symmetries of F_2 , we also want to allow S_∞ to act via a on the second coordinate.

Note that the actions a and b on $(2^\mathbb{N})^\mathbb{N}$ commute, and so give rise to the product action $c = (b, a)$ of the product group $S_\infty \times S_\infty$

$$c: S_\infty \times S_\infty \curvearrowright (2^\mathbb{N})^\mathbb{N}$$

Define an action

$$a_2: S_\infty \times S_\infty \curvearrowright (2^\mathbb{N})^\mathbb{N} \times (2^\mathbb{N})^\mathbb{N}$$

by

$$(g, h) \cdot_{a_2} x = (g \cdot_a x(0), (g, h) \cdot_c x(1)).$$

The corresponding orbit equivalence relation agrees with F_2 on the large, comeager set, of all $x \in (2^\mathbb{N})^\mathbb{N} \times (2^\mathbb{N})^\mathbb{N}$ for which $x(0)$ is an injective enumeration of A_1^x and $(a_1^{x,l} : l \in \mathbb{N})$ is an injective enumeration of A_2^x (recall the definitions from Section 1). More generally:

Definition 1.18. For $n \leq \omega$ define

$$a_n: (S_\infty)^n \curvearrowright ((2^\mathbb{N})^\mathbb{N})^n$$

so that for $g \in (S_\infty)^n$ and $x \in ((2^\mathbb{N})^\mathbb{N})^n$

$$(g \cdot_{a_n} x)(k+1) = (g(k), g(k+1)) \cdot_c x(k+1)$$

and $(g \cdot_{a_n} x)(0) = g(0) \cdot_a x(0)$.

Let X_n^{inj} be the set of all $x \in X_n$ so that $x(0)$ is an injective enumeration of A_x^1 and for each $k < n$, $(a_k^{x,l} : l \in \mathbb{N})$ is an injective enumeration of A_{k+1}^x .

Claim 1.19. For each $n \leq \omega$.

- (1) X_n^{inj} is a comeager subset of X_n , and is a_n -invariant.
- (2) On X_n^{inj} the orbit equivalence relation induced by a_n is F_n .

Remark 1.20. The group action provides another point of view that the presentation F_n is better behaved than $=^{+n}$. For example, $=^{+2}$, defined on $((2^\mathbb{N})^\mathbb{N})^\mathbb{N}$ as the Friedman-Stanley jump of $=^+$, is naturally induced (on a subdomain) by an action of the infinite support wreath product group $S_\infty \wr S_\infty$. This group is defined as the semi-direct product $S_\infty \rtimes (S_\infty)^\mathbb{N}$ with the natural permutation action of $S_\infty \curvearrowright (S_\infty)^\mathbb{N}$. Similarly, the higher jumps $=^{+n}$ can be presented (on a subdomain) as an orbit equivalence relation induced by a natural action of an iterated wreath product of S_∞ . (See [CC22, Proposition 2.3] for example, where variations of the Friedman-Stanley jump are considered.)

1.3. The spectrum of the meager ideal.

Definition 1.21 (Kanovei-Sabok-Zapletal [KSZ13, Definition 1.16]³). An analytic equivalence relation E is **in the spectrum of the meager ideal** if there is an equivalence relation F on a Polish space Y so that

- E and F are Borel bireducible;
- For any non-meager set $C \subseteq Y$, $F \upharpoonright C$ is Borel bireducible with F .

For F as in the second bullet, we say that F **retains its complexity on non-meager sets**.

Kanovei, Sabok, and Zapletal [KSZ13] concluded from Theorem 1.2 that $=^+$, on $\mathbb{R}^{\mathbb{N}}$, retains its complexity on non-meager sets, and is therefore in the spectrum of the meager ideal. The higher jumps *do not* retain their complexity, with the topology coming from the jump operation.

Claim 1.22. There is a comeager set C so that $(=^{+2} \upharpoonright C) \leq_B =^+$.

Proof. Recall that $=^{+2}$ is defined on the space $(\mathbb{R}^{\mathbb{N}})^{\mathbb{N}}$. Let $C \subseteq (\mathbb{R}^{\mathbb{N}})^{\mathbb{N}}$ be the set of all $x \in (\mathbb{R}^{\mathbb{N}})^{\mathbb{N}}$ so that for any $n, m, l, k \in \mathbb{N}$, $(n, m) \neq (l, k) \implies x(n)(m) \neq x(l)(k)$. C is a comeager subset of $(\mathbb{R}^{\mathbb{N}})^{\mathbb{N}}$.

Define $g: C \rightarrow (\mathbb{R} \times \mathbb{R})^{\mathbb{N}^3}$ by $g(x) = ((x(n)(m), x(n)(k)) : n, m, k \in \mathbb{N})$. Fix a bijection $e: \mathbb{N} \rightarrow \mathbb{N}^3$, which extends naturally to a homeomorphism $\hat{e}: (\mathbb{R} \times \mathbb{R})^{\mathbb{N}^3} \rightarrow (\mathbb{R} \times \mathbb{R})^{\mathbb{N}}$. Define $f: C \rightarrow (\mathbb{R} \times \mathbb{R})^{\mathbb{N}}$ by $f = \hat{e} \circ g$.

For $x \in D$, the sets $\{x(n)(m) : m \in \mathbb{N}\}$ are disjoint for different values of n . $f(x)$ is an enumeration of the equivalence relation partitioning the set of reals $\{x(n)(m) : n, m \in \mathbb{N}\}$ into the sets $\{x(n)(m) : m \in \mathbb{N}\} : n \in \mathbb{N}$. It follows that f is a reduction of $(=^{+2} \upharpoonright C)$ to $(=_{\mathbb{R} \times \mathbb{R}})^+$. Since $=_{\mathbb{R}} \sim_B =_{\mathbb{R} \times \mathbb{R}}$, we conclude that $(=^{+2} \upharpoonright C) \leq_B =^+$, as required. \square

Theorem 1.23. For each $1 \leq n \leq \omega$, $=^{+n}$ is in the spectrum of the meager ideal.

Since $F_n \sim_B =^{+n}$, it suffices to prove the following.

Proposition 1.24. F_n on X_n retains its complexity on non-meager sets.

Proof. First we make the following two observations.

- (1) For each $n \leq \omega$, each F_n class is meager in X_n .
- (2) For any $k < n \leq \omega$, u_k^n is not a reduction on any non-meager set.

Fix a non-meager set $Z \subseteq X_n$. We prove that $F_n \leq_B F_n \upharpoonright Z$. First, we claim that there is a Borel homomorphism $f: F_n \rightarrow_B F_n \upharpoonright Z$ which is a reduction on a non-meager set. For such f , the second bullet of Theorem 1.5 fails: if f factors through u_k^n on a comeager set, for $k < n$, then it would follow that u_k^n is a reduction on a non-meager set, contradicting (2) above. We then conclude, by Theorem 1.5, that F_n is Borel reducible to $F_n \upharpoonright Z$.

³Kanovei, Sabok, and Zapletal studied the behavior of equivalence relations on I -positive sets for various ideals I . Here we only mention the case where I is the meager ideal.

Note that if Z is F_n -invariant, it is easy to find a homomorphism f as claimed: simply let f be the identity on Z , and a constant function outside of Z . In general, we can find such a homomorphism using large section uniformization, as follows. We will use below category quantifiers and the Vaught transform. See [Kec95, 8.J] and [Gao09, 3.2].

Recall that, once restricted to a comeager invariant set X_n^{inj} , F_n can be presented as an orbit equivalence relation induced by a continuous action of the Polish group $G = (S_\infty)^n$ (see Section 1.2). We may assume that $Z \subseteq X_n^{\text{inj}}$. Fix a countable dense set $G_0 \subseteq G$. Since almost every orbit is dense, the set $Z' = G_0 \cdot Z$ is comeager. For any $g \in G$, $g^{-1}Z'$ is comeager, that is, $\forall^* x \in X (g \cdot x \in Z')$. We conclude that $\forall^* x \in X \forall^* g \in G (g \cdot x \in Z')$. That is, the invariant set $B = \{x : \forall^* g \in G (g \cdot x \in Z')\}$ is comeager. By [Kec95, Theorem 18.6] there is a Borel map $h: B \rightarrow G$ so that $h(x) \cdot x \in Z'$ for all $x \in B$.

Finally, define $f: X \rightarrow Z$ as follows. Fix $z_0 \in Z$ and an enumeration $(\gamma_n)_{n \in \mathbb{N}}$ of G_0 . If $x \in X \setminus B$, $f(x) = z_0$. If $x \in B$, define $f(x) = \gamma_n \cdot h(x) \cdot x$ for the minimal n so that $\gamma_n \cdot h(x) \cdot x \in Z$. Then f is a Borel homomorphism as claimed. \square

Conjecture 1.25. For each countable ordinal α , $=^{+\alpha}$ is in the spectrum of the meager ideal.

Question 1.26. Is the spectrum of the meager ideal closed under

- (1) the Friedman-Stanley jump operation;
- (2) countable products.

1.4. A question of Clemens. In the context of definable cardinality of quotients of Polish spaces, a Borel homomorphism corresponds to a definable map between two such quotients, and a Borel reduction corresponds to an injective definable map.

Definition 1.27 (Clemens [Cle22]). Let E and F be Borel equivalence relations on Polish spaces X and Y respectively. Say that E is **prime to** F if for any Borel homomorphism $f: E \rightarrow_B F$, E retains its complexity on a fiber, that is, there is $y \in Y$ so that E is Borel reducible to $E \upharpoonright \{x \in X : f(x) F y\}$.

Primeness is a strong form of Borel-irreducibility, which holds between many pairs of benchmark equivalence relations (see [Cle22, Theorem 1]).

In the classical context of cardinality, primeness corresponds to a pigeonhole principle: any function $f: A \rightarrow B$ has a fiber of cardinality $|A|$. This is true if and only if the cardinality $|B|$ is strictly smaller than the cofinality of $|A|$. Recall that the cardinality $|A|$ is regular if it is equal to its cofinality, that is, if for any $|B| < |A|$, any function from A to B has a fiber of size $|A|$.

Following this analogy Clemens defined **regular** equivalence relation as follows.

Definition 1.28 (Clemens [Cle22]). A Borel equivalence relation E is **regular** if for any Borel equivalence relation F , if $F <_B E$ then E is prime to F .

Clemens [Cle22, Question 7.3] asked if $=^{+\omega}$ is regular. We confirm this. (Notational warning: what we call here $=^{+\omega}$ is denoted by \mathbb{F}_ω in [Cle22].)

Theorem 1.29. For any equivalence relation E which is classifiable by countable structures, either $=^{+\omega} \leq_B E$ or $=^{+\omega}$ is prime to E . In particular, $=^{+\omega}$ is regular.

Proof. Note that all the properties above respect Borel bireducibility. In particular, if $E \sim_B E'$ then E is prime to F if and only if E' is prime to F . Therefore, it suffices to prove the theorem with F_ω instead of $=^{+\omega}$.

Fix E as in the theorem and assume that F_ω is not Borel reducible to E . Let $f: F_\omega \rightarrow_B E$ be a Borel homomorphism. By Theorem 1.5 there is some $k < \omega$, a Borel homomorphism $g: F_k \rightarrow_B E$, defined on a comeager set, and a comeager $C \subseteq X_\omega$, so that for any $x \in C$,

$$g(u_k^\omega(x)) E f(x).$$

View $((2^{\mathbb{N}})^{\mathbb{N}})^\omega$ as $((2^{\mathbb{N}})^{\mathbb{N}})^k \times ((2^{\mathbb{N}})^{\mathbb{N}})^{\omega \setminus k}$. Recall that u_k^ω is the projection from $((2^{\mathbb{N}})^{\mathbb{N}})^k \times ((2^{\mathbb{N}})^{\mathbb{N}})^{\omega \setminus k}$ to $((2^{\mathbb{N}})^{\mathbb{N}})^k$. By the Kuratowski-Ulam theorem (see [Kec95, Theorem 8.41 (iii)]) there is $y \in ((2^{\mathbb{N}})^{\mathbb{N}})^k$ so that $C_y = \{z \in ((2^{\mathbb{N}})^{\mathbb{N}})^{\omega \setminus k} : (y, z) \in C\}$ is comeager in $((2^{\mathbb{N}})^{\mathbb{N}})^{\omega \setminus k}$. Note that $\{y\} \times C_y$ is contained in the fiber $\{x \in X_\omega : f(x) E g(y)\}$. We will finish the proof by showing that F_ω is Borel reducible to $F_\omega \upharpoonright \{y\} \times C_y$.

Consider the homeomorphism $\phi: ((2^{\mathbb{N}})^{\mathbb{N}})^{\omega \setminus k} \rightarrow ((2^{\mathbb{N}})^{\mathbb{N}})^\omega$, defined by $\phi(z)(l) = z(l+k)$. Then for $z_1, z_2 \in ((2^{\mathbb{N}})^{\mathbb{N}})^{\omega \setminus k}$,

$$(y, z_1) F_\omega (y, z_2) \iff \phi(z_1) F_\omega \phi(z_2).$$

The set $\phi(C_y)$ is comeager in $((2^{\mathbb{N}})^{\mathbb{N}})^\omega$, as ϕ is a homeomorphism. Since F_ω retains its complexity of comeager sets, there is a Borel reduction $h: F_\omega \rightarrow F_\omega \upharpoonright \phi(C_y)$. Finally, the map

$$x \mapsto (y, \phi^{-1}(h(x)))$$

is a Borel reduction of F_ω to $F_\omega \upharpoonright \{y\} \times C_y$, as required. \square

Clemens [Cle22, Lemma 7.6] showed that if $\alpha \geq 2$ is not of the form ω^β , for some countable ordinal β , then $=^{+\alpha}$ is not regular.

Question 1.30 (See [Cle22, Question 7.3]). For a countable ordinal β , is $=^{+\omega^\beta}$ regular?

As in Definition 1.28, Clemens defined an equivalence relation E as **prime** if for any Borel equivalence relation F , either $E \leq_B F$ or E is prime F . In the context of definable cardinality, when not every two sizes are comparable, this is a strengthening of being regular. A positive answer to Question 1.10 will imply that $=^{+\omega}$ is prime.

Question 1.31. Is $=^{+\omega}$ prime?

1.5. Non-reduction to products. Given equivalence relations E_k on X_k , the **product equivalence relation** $\prod_k E_k$ is defined on the space $\prod_k X_k$ by

$$x \prod_k E_k y \iff x(k) E_k y(k) \text{ for all } k.$$

We write $E^{\mathbb{N}}$ for the product $\prod_k E_k$ where $E_k = E$ for all k . The product operation plays an important role in the study of Borel equivalence relations. For example, it follows from the dichotomy theorem proved by Hjorth and Kechris [HK01] that the equivalence relation $E_0^{\mathbb{N}}$, also known as E_3 , is an immediate successor of E_0 with respect to \leq_B . When studying jump operations on Borel equivalence relations, a product is often used to define the limit stages of iterated jumps. The definition of $=^{+\omega}$ as $\prod_n =^{+n}$ is one such example. The following result shows that, for $n < \omega$, $=^{+n}$ cannot be presented as a product of strictly simpler equivalence relations.

Proposition 1.32. Fix $n < \omega$. For $k < \omega$, let E_k be an equivalence relation, classifiable by countable structures, so that $=^{+n} \not\leq_B E_k$. Then $=^{+n} \not\leq_B \prod_k E_k$.

Proof. We may replace $=^{+n}$ by F_n . Note that a Borel homomorphism $f: F_n \rightarrow_B \prod_k E_k$ can be identified with a sequence of Borel homomorphisms $f_k: F_n \rightarrow_B E_k$. By Theorem 1.5, each f_k factors, on a comeager set, through u_{n-1}^n . It follows that f factors through u_{n-1}^n on a comeager set. In particular, a Borel homomorphism $f: F_n \rightarrow \prod_k E_k$ cannot be a reduction. \square

This was proved (for all analytic equivalence relations) for $n = 1$ by Kanovei, Sabok, and Zapletal [KSZ13, Corollary 6.30]. The result is phrased there in terms of intersections of equivalence relations. Given equivalence relations E_k on a common space X , let their intersection $\bigcap_k E_k$ be the equivalence relation on X defined by $x \bigcap_k E_k y \iff x E_k y$ for every k .

There is a close relationship between products and intersections. Note that $\prod_k E_k$ can be written as an intersection of equivalence relations E'_k on $\prod_k X_k$ so that $E'_k \sim_B E_k$ for each k . Furthermore, the intersection $\bigcap_k E_k$ is Borel reducible to $\prod_k E_k$, witnessed by the diagonal map $X \rightarrow X^{\mathbb{N}}$, $x \mapsto (x, x, \dots)$. Therefore Proposition 1.32 is equivalent to a similar result for intersections:

Corollary 1.33. Fix $n < \omega$ and a Polish space X . For $k < \omega$, let E_k be an equivalence relation on X , classifiable by countable structures, so that $=^{+n} \not\leq_B E_k$. Then $=^{+n} \not\leq_B \bigcap_k E_k$.

1.6. Borel complexity. The equivalence relations $=^{+n}$, $n < \omega$, are naturally written as $\mathbf{\Pi}_{2n+1}^0$ relations on their domains, where $=^+$ is $\mathbf{\Pi}_3^0$, and each application of the Friedman-Stanley jump operator adds an alternating $\forall \exists$ quantification. The equivalence relations $=^{+n}$ are in fact simpler, in terms of **potential complexity** [HKL98]. We refer the reader to [Lou94] or [HKL98] for the definition. An equivalent definition is: E is **potentially Γ** , for a point-class Γ , if E is Borel reducible to some equivalence relation F , where F is in Γ .

Hjorth, Kechris, and Louveau [HKL98] proved that the optimal potential complexity of $=^{+n}$ is precisely $\mathbf{\Pi}_{2+n}^0$. In fact, they proved that among S_∞ -actions $=^{+n}$ is a maximal equivalence relation with this potential complexity. Moreover, they extended these results for the transfinite jumps and completely classified the possible potential complexities of Borel equivalence relations induced by an S_∞ action.

Here we simply note that the equivalence relations F_n , defined to optimize Baire-category considerations, naturally have the optimal potential complexity.

Proposition 1.34. The relation F_n is $\mathbf{\Pi}_{2+n}^0$ as a subset of $X_n \times X_n$.

Proof. For F_1 , which is $=^+$, a direct computation shows that it is $\mathbf{\Pi}_3^0$.

We define relations Q_n on $X_n \times 2^{\mathbb{N}}$, for $0 \leq n < \omega$, so that

- Q_n is $\mathbf{\Pi}_{n+1}$, and
- for $1 \leq n$, for $x, y \in X_{n+1}$, $x F_{n+1} y$ if and only if $u_n^{n+1}(x) F_n u_n^{n+1}(y)$ and

$$\forall n_1 \exists n_2 \forall l_1, l_2 [(u_{n-1}^{n+1}(x), x(n-1)(l_1)) Q_{n-1} (u_{n-1}^{n+1}(y), y(n-1)(l_2)) \rightarrow x(n)(n_1)(l_1) = y(n)(n_2)(l_2)].$$

Assuming this, for $1 \leq n$, as Q_{n-1} is $\mathbf{\Pi}_n$, the expression in the square brackets is Σ_n , which shows that F_{n+1} is $\mathbf{\Pi}_{n+3}^0$, as required.

For $n = 0$, define Q_0 as equality on $2^{\mathbb{N}}$. Note that we identify X_0 as a space with 1 member, and so we identify $X_0 \times 2^{\mathbb{N}}$ with $2^{\mathbb{N}}$. For $n \geq 1$, given $(x, v) \in X_n \times 2^{\mathbb{N}}$, recall the definition of $A_n^x = \{a_{n-1}^{x,l} : l \in \mathbb{N}\}$. Define $a_n^{x,v} = \{a_{n-1}^{x,t} : v(t) = 1\} \subseteq A_n^x$. Given $(x, v), (y, w) \in X_n \times 2^{\mathbb{N}}$, define

$$(x, v) Q_n (y, w) \iff A_n^x = A_n^y \wedge a_n^{x,v} = a_n^{y,w}.$$

The relation $(x, v) Q_{n+1} (y, w)$ is true if and only if

$$(x F_n y) \wedge \forall l_1, l_2 [(u_{n-1}^n(x), x(n-1)(l_1)) Q_n (u_{n-1}^n(y), y(n-1)(l_2)) \rightarrow v(l_1) = w(l_2)].$$

Inductively, F_n is $\mathbf{\Pi}_{n+2}^0$ and Q_n is $\mathbf{\Pi}_{n+1}^0$. We conclude that Q_{n+1} is $\mathbf{\Pi}_{n+2}^0$. \square

2. COMPLEXITY ON COMEAGER SETS: SOME IDEAS AND SOME OBSTACLES

One obstacle towards the $n > 1$ case was already encountered. The natural topology coming from the Friedman-Stanley jump operation does not work (see 1.22), and we therefore had to find the “correct” presentation of these equivalence relations, as in Section 1.1.

Let us focus on a corollary of the main theorem, that the equivalence relations F_n retain their complexity on comeager sets (see Section 1.3). In this section we sketch some ideas behind the proof for F_1 , and explain why a different type of construction is necessary to deal with F_n for $n > 1$.

2.1. The case $n = 1$. The fact that F_1 (which is $=^+$) retains its complexity on comeager sets was proven in [KSZ13]. Given a comeager set $C \subseteq (2^{\mathbb{N}})^{\mathbb{N}}$, let C^* be its Vaught transform (see [Gao09, 3.2.2]), $C^* = \{a \in (2^{\mathbb{N}})^{\mathbb{N}} : (\forall^* g \in S_\infty) g \cdot a \in C\}$.

Fix a map $g: (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}}$. Define $f_0: (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N} \times \mathbb{N}}$ by

$$f_0(x)(k, l) = g(x(k))(l).$$

Fix a bijection $e: \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$. This extends naturally to a homeomorphism $\hat{e}: (2^{\mathbb{N}})^{\mathbb{N} \times \mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}}$. Let $C: (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}}$ be a Borel map so that for $x \in (2^{\mathbb{N}})^{\mathbb{N}}$, if

the set enumerated by x is finite, then $C(x) = x$, and if the set enumerated by x is infinite, then $C(x)$ is an injective enumeration of the same set. Define

$$f = C \circ \hat{e} \circ f_0.$$

For a “sufficiently generic” choice of map g , it can be verified that $f(x) \in C^*$ for all $x \in (2^{\mathbb{N}})^{\mathbb{N}}$. It follows that there is a Borel map $\rho: (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow S_{\infty}$ so that $\rho(x) \cdot f(x) \in C$. Finally, the map $x \mapsto \rho(x) \cdot f(x)$ is a reduction of F_1 to $F_1 \upharpoonright C$.

2.2. The case $n > 1$. Below we explain why a direct generalization of the construction in Section 2.1, to construct a map reducing F_2 to some comeager subset of F_2 , does not work. Fix $(x, y) \in (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$ in the domain of F_2 . We would want to define $f(x, y)$ to be of the form (u, v) so that (u, v) is “sufficiently generic”, in the sense that it lands in the Vaught transform of some comeager set.

We can start by defining u from x as before, so that u is “sufficiently generic”. We may hope to define v from y in the same way, so that v is also “sufficiently generic”. The problem can be seen from the group action presentation in Section 1.2. At the second level, we have the usual permutation action of S_{∞} (the action a), but also another copy of S_{∞} acting “from behind” via the action b . The construction in Section 2.1, which is invariant under the action a , is *not* invariant under the action b , and therefore the resulting map will not respect F_2 .

This difficulty can also be seen from a set theoretic perspective. Let $A = A^x$ and $B = A^u$, their corresponding classifying F_1 -invariants. We may want to replace the space $\mathcal{P}_{\aleph_0}(\mathbb{N})$ (which is identified with $2^{\mathbb{N}}$) with the space $\mathcal{P}_{\aleph_0}(A)$. Now we may hope to follow the construction of Section 2.1 to find a “definable” map taking some $Y \in \mathcal{P}_{\aleph_0}(A)$ to a “sufficiently generic” member of $\mathcal{P}_{\aleph_0}(B)$. (The quotation marks are intended to mean that once translated in a reasonable way to a map defined on our Polish space X_2 , it will be Borel definable, and land in some comeager set.) This construction should be done independently of the enumerations of A and Y , for the resulting map to be a homomorphism $F_2 \rightarrow_B F_2$. This hope is immediately crushed. Such constructions are common with $A = \omega$, or more generally an ordinal, but impossible for higher rank sets.

The point of this discussion is to mention that our construction of v , towards $f(x, y) = (u, v)$, has to rely on the enumerations coming from x , while ultimately being independent of those, up to F_2 -equivalence. It cannot be done by a direct iteration of the previous construction. The main new construction, which deals with the $n > 1$ case, is presented in Section 4, Definition 4.1.

We also present in Section 4.1 a variation of the above sketched construction for the $n = 1$ case. This variation is needed simply to “align” the two constructions, as in Section 4.3.

3. PERMUTATIONS

In various points below we will want a member of some product space, constructed in a specific way, to land in some comeager set. As in Section 2.1 we will be able

to guarantee this only after applying a group action. The following lemma will be used to deal with the construction for the $n > 1$ case.

Let S, X, Y_1, \dots, Y_k be infinite sets, considered as discrete metric spaces. Consider the space $(2^S)^X \times (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$ with the product topology. Consider the natural diagonal action of $\text{Sym}(X)$ on $(2^S)^X \times (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$, acting on all copies of X simultaneously. For $i = 1, \dots, k$, consider the natural action of and $\text{Sym}(Y_i)$ on $(2^S)^X \times (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$. These actions commute, leading to an action

$$\text{Sym}(X) \times \text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k) \curvearrowright (2^S)^X \times (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}.$$

We consider each $\text{Sym}(Y_i)$, and $\text{Sym}(X)$, as a topological group with the point-wise convergence topology, and $\text{Sym}(X) \times \text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k)$ with the product topology.

Lemma 3.1. Let $D \subseteq (2^S)^X \times (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$ be dense open. Fix $(\zeta, \xi_1, \dots, \xi_k) \in (2^S)^X \times (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$ satisfying the following assumptions:

- (1) For any finite permutation π of X , the set

$$D_{\pi \cdot \zeta} = \{(\delta_1, \dots, \delta_k) : (\pi \cdot \zeta, \delta_1, \dots, \delta_k) \in D\}$$

is dense in $(2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$.

- (2) (a) For any finite partial function $\tau: X \rightarrow \{0, 1\}$ and any $i \in \{1, \dots, k\}$, there are infinitely many $y \in Y_i$ so that $\xi_i(y)(-)$, considered as a function $X \rightarrow \{0, 1\}$, extends τ .
- (b) Given finite partial functions $\tau: S \rightarrow \{0, 1\}$, $\tau_i: Y_i \rightarrow \{0, 1\}$, $i = 1, \dots, k$, there are infinitely many $x \in X$ so that $\xi_i(-)(x)$, considered as a function $Y_i \rightarrow \{0, 1\}$, extends τ_i , for every $i = 1, \dots, k$, and $\zeta(x)(-)$, considered as a function $S \rightarrow \{0, 1\}$, extends τ .

Then the set

$$G = \{(g, g_1, \dots, g_k) \in \text{Sym}(X) \times \text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k) : (g, g_1, \dots, g_k) \cdot (\zeta, \xi_1, \dots, \xi_k) \in D\}$$

is dense open in $\text{Sym}(X) \times \text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k)$. In particular, if D is assumed to be comeager, then G is concluded to be comeager.

Proof. First, since the map $\text{Sym}(X) \times \text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k) \rightarrow (2^S)^X \times (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$, $(g, g_1, \dots, g_k) \mapsto (g, g_1, \dots, g_k) \cdot (\zeta, \xi_1, \dots, \xi_k)$, is continuous, then G is open as the pre-image of D .

Next we prove that G is dense. Fix finite partial permutations π, π_1, \dots, π_k of X, Y_1, \dots, Y_k respectively. We need to find an extension of these in G . Let $\bar{X}, \bar{Y}_1, \dots, \bar{Y}_k$ be the finite supports of π, π_1, \dots, π_k , respectively.

By assumption (1), $D_{\pi \cdot \zeta}$ is dense. Fix $(\delta_1, \dots, \delta_k) \in D_{\pi \cdot \zeta}$ which agree with $(\pi_1 \cdot \xi_1, \dots, \pi_k \cdot \xi_k)$ on $(2^{\bar{X}})^{\bar{Y}_1} \times \dots \times (2^{\bar{X}})^{\bar{Y}_k}$. Since D is open, we may find finite $\hat{X}, \hat{Y}_1, \dots, \hat{Y}_k$, extending $\bar{X}, \bar{Y}_1, \dots, \bar{Y}_k$, and a finite set \hat{S} , so that if $(\zeta', \xi'_1, \dots, \xi'_k)$ agree with $(\pi \cdot \zeta, \delta_1, \dots, \delta_k)$ on $(2^{\hat{S}})^{\hat{X}} \times (2^{\hat{X}})^{\hat{Y}_1} \times \dots \times (2^{\hat{X}})^{\hat{Y}_k}$, then $(\zeta', \xi'_1, \dots, \xi'_k) \in D$.

It remains to find (g, g_1, \dots, g_k) , extending π, π_1, \dots, π_k , so that $(g, g_1, \dots, g_k) \cdot (\zeta, \xi_1, \dots, \xi_k)$ and $(\pi \cdot \zeta, \delta_1, \dots, \delta_k)$ agree on $(2^{\hat{S}})^{\hat{X}} \times (2^{\hat{X}})^{\hat{Y}_1} \times \dots \times (2^{\hat{X}})^{\hat{Y}_k}$.

For each $i = 1, \dots, k$, for each $y \in \hat{Y}_i \setminus \bar{Y}_i$, consider the function $\tau_y: \bar{X} \rightarrow \{0, 1\}$, $\tau_y(x) = \delta_i(y)(x)$. By assumption (2)(a), there are infinitely many $y' \in Y_i$ so that $\xi_i(y')(-)$ and $\tau_y(-)$ agree on \bar{X} . It follows that there is a finite permutation g_i extending π_i so that $(g_i \cdot \xi_i)(y)(-)$ and $\delta_i(y)(-)$ agree on \bar{X} , for all $y \in \hat{Y}_i \setminus \bar{Y}_i$. It follows that $(\pi, g_1, \dots, g_k) \cdot (\zeta, \xi_1, \dots, \xi_k)$ and $(\pi \cdot \zeta, \delta_1, \dots, \delta_k)$ agree on $(2^{\hat{S}})^{\hat{X}} \times (2^{\bar{X}})^{\hat{Y}_1} \times \dots \times (2^{\bar{X}})^{\hat{Y}_k}$.

Note that conditions (2)(a) and (2)(b) of the lemma are invariant under the group action. We apply condition (2)(b) to $(\pi, g_1, \dots, g_k) \cdot (\zeta, \xi_1, \dots, \xi_k)$. We may write $(\pi, g_1, \dots, g_k) \cdot (\zeta, \xi_1, \dots, \xi_k)$ as $(\pi \cdot \zeta, (\pi, g_1) \cdot \xi_1, \dots, (\pi, g_k) \cdot \xi_k)$, where $(\pi, g_i) \cdot \xi_i$ refers to the action $\text{Sym}(X) \times \text{Sym}(Y_i) \curvearrowright (2^S)^X \times (2^X)^{Y_i}$.

For each $i = 1, \dots, k$, for each $x \in \hat{X} \setminus \bar{X}$, consider the function $\tau_i^x: \hat{Y}_i \rightarrow \{0, 1\}$, $\tau_i^x(y) = \delta_i(y)(x)$. Define also $\tau^x: \hat{S} \rightarrow \{0, 1\}$ by $\tau^x(s) = (\pi \cdot \zeta)(x)(s)$. For each $x \in \hat{X} \setminus \bar{X}$ there are infinitely many $x' \in X$ so that $((\pi, g_i) \cdot \xi_i)(-)(x')$ extends τ_i^x and $(\pi \cdot \zeta)(x')(-)$ extends τ^x . It follows that there is a finite permutation g extending π so that $((g, g_i) \cdot \xi_i)(-)(x)$ and $\delta_i(-)(x)$ agree on \hat{Y}_i , for all i and any $x \in \hat{X} \setminus \bar{X}$, and $(g \cdot \zeta)(-)(x)$ agrees with $\pi \cdot \zeta(-)(x)$ on \hat{S} for any $x \in \hat{X} \setminus \bar{X}$. We conclude that $(g, g_1, \dots, g_k) \cdot (\zeta, \xi_1, \dots, \xi_k)$ and $(\pi \cdot \zeta, \delta_1, \dots, \delta_k)$ agree on $(2^{\hat{S}})^{\hat{X}} \times (2^{\hat{X}})^{\hat{Y}_1} \times \dots \times (2^{\hat{X}})^{\hat{Y}_k}$, as required. \square

3.1. The $n = 1$ case. When dealing with the first coordinate of F_n we only have the action $a: S_\infty \curvearrowright (2^{\mathbb{N}})^{\mathbb{N}}$. More generally, we deal with product actions of the form

$$\text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k) \curvearrowright (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}.$$

Lemma 3.2. Fix a dense open $D \subseteq (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$ and countable infinite sets M_1, \dots, M_k . Then there is a dense open set $D' \subseteq (2^X)^{M_1} \times \dots \times (2^X)^{M_k}$ so that for any $\zeta \in (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$, if ζ satisfies the following property: given any finite partial injective functions $\tau_i: M_i \rightarrow Y_i$, $i = 1, \dots, k$, there are extensions $\alpha_i: M_i \rightarrow Y_i$ so that $(\zeta(i) \circ \alpha_i : i < k)$, a member of the space $(2^X)^{M_1} \times \dots \times (2^X)^{M_k}$, is in D' , then the set

$$G = \{(g_1, \dots, g_k) \in \text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k) : (g_1, \dots, g_k) \cdot \zeta \in D\}$$

is dense open in $\text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k)$. In particular, if D is assumed to be comeager, then there is a comeager D' so that G is concluded to be comeager.

Proof. First, since the map $\text{Sym}(Y_1) \times \dots \times \text{Sym}(Y_k) \rightarrow (2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$, $(g_1, \dots, g_k) \mapsto (g_1, \dots, g_k) \cdot \zeta$, is continuous, then G is open as the pre-image of D .

Next, we describe the set D' so that if ζ satisfies the assumption in the lemma, then the set G is dense. First assume that $M_i = Y_i$ and take D' to be D . The assumption tells us that for any finite partial permutations $\tau_i: M_i \rightarrow Y_i$ there are extensions to total injective maps $\alpha_i: Y_i \rightarrow Y_i$ so that $(\zeta(i) \circ \alpha_i : i < k) \in D$. If α_i were all bijections, so in $\text{Sym}(Y_i)$, we would be done. Nevertheless, since D

is open, we may find bijections σ_i of Y_i which extend τ_i and are sufficiently close to α_i so that $(\zeta(i) \circ \sigma_i : i < k) \in D$ as well. Finally, for any infinite countable M_1, \dots, M_k , we may fix bijections $M_i \rightarrow Y_i$, resulting in a homeomorphism between $(2^X)^{Y_1} \times \dots \times (2^X)^{Y_k}$ and $(2^X)^{M_1} \times \dots \times (2^X)^{M_k}$. We let D' be the image of D . \square

4. THE MAIN CONSTRUCTION

Definition 4.1. Fix a function

$$\alpha: (2^{<\mathbb{N}})^{<\mathbb{N}} \times \mathbb{N} \times \mathbb{N} \rightarrow 2$$

(which will be chosen to be “sufficiently generic”). Define

$$\beta: (2^{\mathbb{N}})^{<\mathbb{N}} \rightarrow (2^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}})^{\mathbb{N}}$$
 by

$$\beta(x_1, \dots, x_l)(m)(t, k) = \alpha(x_1 \circ t, \dots, x_l \circ t, k, m), \text{ for } t \in \mathbb{N}^{<\mathbb{N}}; k, m \in \mathbb{N}.$$

Each x_i is considered a function $\mathbb{N} \rightarrow 2$, and so $x_i \circ t$ is a member of $2^{<\mathbb{N}}$. Define

$$\gamma: (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow (2^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}})^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}}$$
 by

$$\gamma(x)(t, k) = \beta(x \circ t)(k).$$

Remark 4.2.

- The function γ is continuous.
- The function γ is a homomorphism from the orbit equivalence relations

$$\text{Sym}(\mathbb{N}) \curvearrowright (2^{\mathbb{N}})^{\mathbb{N}} \text{ to } \text{Sym}(\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}) \curvearrowright (2^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}})^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}}, \text{ and}$$

$$\text{Sym}(\mathbb{N}) \curvearrowright (2^{\mathbb{N}})^{\mathbb{N}} \text{ to } \text{Sym}(\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}) \curvearrowright (2^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}})^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}}$$

- The definition of γ relies on a choice of α . We will show that there is some α for which γ satisfies the properties which we need. This will happen for α chosen generically, with respect to the product topology $2^{(2^{<\mathbb{N}})^{<\mathbb{N}} \times \mathbb{N} \times \mathbb{N}}$.

Notation 4.3. Let $N = \mathbb{N}^{<\mathbb{N}} \times \mathbb{N}$.

It will be convenient, to utilize the construction above, to work with the space $(2^{\mathbb{N}})^N$ instead of $(2^{\mathbb{N}})^{\mathbb{N}}$. To illustrate the construction, consider the following lemma.

Say that $a \in (2^{\mathbb{N}})^{\mathbb{N}}$ is **injective** if it is a sequence of distinct reals: $a(i) \neq a(j)$ for $i \neq j$. Say that $a \in (2^{\mathbb{N}})^{\mathbb{N}}$ is **separated** if any distinct $n, k \in \mathbb{N}$ are separated by one of the members of a : there is some $i \in \mathbb{N}$ so that $a(i)(n) \neq a(i)(k)$.

Lemma 4.4. For a generic α the following holds. Suppose $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ is separated and $y \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective and separated. Then the pair $(\zeta, \xi_1) = (\gamma(x), \gamma(y)) \in (2^S)^X \times (2^X)^{Y_1}$, where $S = N$, $X = N$, and $Y_1 = N$, satisfies the assumptions (2) in Lemma 3.1. That is:

- for any finite $\tau: N \rightarrow \{0, 1\}$ there are infinitely many $(t, k) \in N$ so that $\gamma(y)(t, k)(-)$ extends τ , and
- for any finite $\tau_1: N \rightarrow \{0, 1\}$ and $\tau_2: N \rightarrow \{0, 1\}$ there are infinitely many $(s, d) \in N$ so that $\gamma(y)(-)(s, d)$ extends τ_1 and $\gamma(x)(s, d)(-)$ extends τ_2 .

Proof. Consider the second bullet, which corresponds to condition (b) Lemma 3.1. Fix finite partial function $\tau_1: Y \rightarrow \{0, 1\}$, $\tau_2: S \rightarrow \{0, 1\}$. Recall the definitions. Fix $d, k \in \mathbb{N}$ and $s, t \in \mathbb{N}^{<\mathbb{N}}$, $s = (s_1, \dots, s_m)$, $t = (t_1, \dots, t_l)$ for some m, l . We abbreviate

$$\begin{aligned} x[s, t] &= (x(s_1) \circ t, \dots, x(s_m) \circ t) \in (2^{<\mathbb{N}})^m, \\ y[t, s] &= (y(t_1) \circ s, \dots, y(t_l) \circ s) \in (2^{<\mathbb{N}})^l. \end{aligned}$$

Then

$$\begin{aligned} \gamma(x)(s, d)(t, k) &= \alpha(x[s, t], k, d), \\ \gamma(y)(t, k)(s, d) &= \alpha(y[t, s], d, k). \end{aligned}$$

So we need to find infinitely many (s, d) for which

$$(\star) \quad \begin{aligned} \alpha(x[s, t], k, d) &= \tau_2(t, k), \text{ for all } (t, k) \text{ in the domain of } \tau_2. \\ \alpha(y[t, s], d, k) &= \tau_1(t, k), \text{ for all } (t, k) \text{ in the domain of } \tau_1. \end{aligned}$$

Since x is separated, given $t_1 \neq t_2 \in \mathbb{N}^{<\mathbb{N}}$ there is some $i \in \mathbb{N}$ so that $x(i) \circ t_1 \neq x(i) \circ t_2$. Construct a sequence $s^* \in \mathbb{N}^{<\mathbb{N}}$ so that for any two distinct t_1, t_2 in the domain of τ_2 there is some i so that $x(s_i^*) \circ t_1 \neq x(s_i^*) \circ t_2$. Note that if s is a sequence which contains s^* then $x[s, t_1] \neq x[s, t_2]$, for any distinct t_1, t_2 in the domain of τ_2 .

Since y is injective, for any $t_1 \neq t_2$ the finite sequences of reals $y \circ t_1$ and $y \circ s_1$ are not equal. Then for any s whose range contains a long enough initial segment of \mathbb{N} (depending on t_1, t_2), $y[t_1, s] \neq y[t_2, s]$. We may find some s^{**} so that for any s which contains s^{**} , for any distinct t_1, t_2 in the domain of τ_1 , $y[t_1, s] \neq y[t_2, s]$.

Fix s which contains both s^* and s^{**} . Then for any $(t_1, k_1), (t_2, k_2)$ from the domain of τ_2 or τ_1 , if the tuples $(t_1, k_1), (t_2, k_2)$ are distinct, then

- the tuples $(x[s, t_1], k_1), (x[s, t_2], k_2)$ are distinct, and
- the tuples $(y[t_1, s], k_1), (y[t_2, s], k_2)$ are distinct.

We claim that, for a generic choice of α , there are infinitely many $d \in \mathbb{N}$ for which (\star) holds with (s, d) . It suffices to prove that for any finite number M there is a dense open set of $\alpha \in 2^{(2^{<\mathbb{N}})^{<\mathbb{N}} \times \mathbb{N} \times \mathbb{N}}$ for which there are at least M many values of d so that (\star) holds for (s, d) . Indeed, given any finite amount of information about α , for a large enough d we have that

- $\alpha(x[s, t], k, d), \alpha(y[t, s], d, k)$ is not yet defined,
- and the tuples $(y[t, s], d, k), (x[s, t], k, d)$ respectively are distinct for distinct (t, k) in the domains of either τ_1, τ_2 .

We may therefore extend α by adding arbitrarily many values of d for which (\star) holds for (s, d) . Note that there are only countably many dense open sets for $\alpha \in 2^{(2^{<\mathbb{N}})^{<\mathbb{N}} \times \mathbb{N} \times \mathbb{N}}$ involved in the argument above, independently of x, y .

Finally, note that the property for $\gamma(y)$ in the first bullet is the same as the property for $\gamma(x)$ in the second bullet. Since y is separated, which is the only assumption we made on x in the proof, we conclude the first bullet as well. \square

4.1. A variation for the $n = 1$ construction. We will also modify the construction in the $n = 1$ case, as outlined in Section 2.1, to be compatible with the above. Fix a continuous function $\beta_0: (2^{\mathbb{N}})^{<\mathbb{N}} \rightarrow (2^N)^{\mathbb{N}}$ (which will be chosen to be “sufficiently generic”). Define $\gamma_0: (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow (2^N)^N$ by

$$\gamma_0(x)(t, k) = \beta_0(x \circ t)(k), \text{ for } t \in \mathbb{N}^{<\mathbb{N}}, k \in \mathbb{N}.$$

Remark 4.5.

- The function γ_0 is continuous.
- The function γ_0 is a homomorphism from the orbit equivalent relation

$$\text{Sym}(\mathbb{N}) \curvearrowright (2^{\mathbb{N}})^{\mathbb{N}} \text{ to } \text{Sym}(\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}) \curvearrowright (2^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}})^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}}.$$

The function β_0 is constructed as follows. Fix comeager sets $C_n \subseteq ((2^N)^{\mathbb{N}})^n$. Fix a continuous $\alpha_0: 2^{\mathbb{N}} \rightarrow (2^N)^{\mathbb{N}}$ with the property that for any pairwise distinct $x_1, \dots, x_n \in 2^{\mathbb{N}}$, $(\alpha_0(x_1), \dots, \alpha_0(x_n)) \in C_n$ (see [Kec95, Theorem 19.1]). Define $\beta_0: (2^{\mathbb{N}})^{<\mathbb{N}} \rightarrow (2^N)^{\mathbb{N}}$ by $\beta_0 = \alpha_0 \circ \iota$, where $\iota: (2^{\mathbb{N}})^{<\mathbb{N}} \hookrightarrow 2^{\mathbb{N}}$ is a continuous injective map.

Notation 4.6. We will say “for almost any β_0 ” or “for a generic β_0 ” to mean β_0 constructed with an appropriate choice of the comeager sets C_n .

As in Lemma 4.4, consider the following example.

Lemma 4.7. Suppose $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective. Then for any comeager $C \subseteq (2^N)^N$, for a generic β_0 , $\gamma_0(x) \in (2^N)^N$ satisfies the assumption in Lemma 3.2, with $X = N$, $Y_1 = N$, and $M_1 = \mathbb{N}$. That is, for $C' \subseteq (2^N)^N$ given by Lemma 3.2, for any finite partial injective $\tau: \mathbb{N} \rightarrow N$ there is an extension $\sigma: \mathbb{N} \rightarrow N$ of τ so that $\gamma_0(x) \circ \sigma \in C'$.

Proof. We describe how to choose the sets $C_n \subseteq ((2^{\mathbb{N}})^N)^n$ in the definition of β_0 . Fix a bijection $e: \mathbb{N} \rightarrow \mathbb{N}^{<\mathbb{N}}$ and let $N_n = \{e(i) : i < n\}$, for $0 < n$. For each $n \in \mathbb{N}$, fix a bijection $s_n: \mathbb{N} \rightarrow N_n \times \mathbb{N}$, inducing a homeomorphism $\hat{s}_n: (2^N)^{N_n \times \mathbb{N}} \rightarrow (2^N)^{\mathbb{N}}$. In particular, $\hat{s}_n^{-1}D'$ is dense open in $(2^N)^{N_n \times \mathbb{N}}$, which we identify with $((2^N)^{\mathbb{N}})^{N_n}$, and in turn with $((2^N)^{\mathbb{N}})^n$. Choose comeager C_n which is a subset of $\hat{s}_n^{-1}D'$ for any bijection $s: \mathbb{N} \rightarrow N_n \times \mathbb{N}$ which agrees with s_n on all but finitely many values. Since x is injective, $x \circ t_1 \neq x \circ t_2$ for distinct $t_1, t_2 \in \mathbb{N}^{<\mathbb{N}}$. By the choice of β_0 , $(\beta_0(x \circ e(0)), \dots, \beta_0(x \circ e(n-1))) \in C_n$ for all $0 < n$.

Let $\tau: \mathbb{N} \rightarrow N$ be a finite partial injection. Recall that $N = \mathbb{N}^{<\mathbb{N}} \times \mathbb{N}$. Choose n so that the image of τ is included in $N_n \times \mathbb{N}$. Let $s: \mathbb{N} \rightarrow N_n \times \mathbb{N}$ be a bijection which extends τ and agrees with s_n on all but finitely many values. Note that $\hat{s}_n^{-1}(\gamma_0(x) \circ s) \in (2^N)^{N_n \times \mathbb{N}}$, which is identified with $(\beta_0(x \circ e(0)), \dots, \beta_0(x \circ e(n-1)))$, is in C_n , and therefore $\gamma_0 \circ s \in D'$, as required. \square

We will also need the following variation of Lemma 4.4.

Lemma 4.8. For generic α and β_0 the following holds. Suppose $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective, and $y \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective and separated. The pair $(\zeta, \xi_1) = (\gamma_0(x), \gamma(y)) \in (2^S)^X \times (2^X)^{Y_1}$, where $S = N$, $X = N$, and $Y_1 = N$, satisfies the assumptions (2) in Lemma 3.1. That is:

- for any finite $\tau: N \rightarrow \{0, 1\}$ there are infinitely many $(t, k) \in N$ so that $\gamma(y)(t, k)(-)$ extends τ , and
- for any finite $\tau_1: N \rightarrow \{0, 1\}$ and $\tau_2: N \rightarrow \{0, 1\}$ there are infinitely many $(s, d) \in N$ so that $\gamma(y)(-)(s, d)$ extends τ_1 and $\gamma(x)(s, d)(-)$ extends τ_2 .

Most aspects of the proof are similar to Lemma 4.4

4.2. A change of countable base set. Recall Notation 4.3: $N = \mathbb{N}^{<\mathbb{N}} \times \mathbb{N}$.

Definition 4.9 (Definition of F_n^* on X_n^*). Consider the space $((2^{\mathbb{N}})^{\mathbb{N}})^{\omega}$, which is naturally homeomorphic to $((2^{\mathbb{N}})^{\mathbb{N}})^{\omega}$. We may define F_n^* on $X_n^* \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^n$, as F_n was defined on $X_n \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^n$ in Section 1.1. See Section 5.1 for more details.

The equivalence relation F_n^* is isomorphic to F_n via a homeomorphism of their domains. To prove that F_n retains its complexity on comeager sets, it suffices to prove that F_n^* retains its complexity on comeager sets.

4.3. The homomorphism $f: F_n \rightarrow F_n^*$.

Definition 4.10. Define $f: ((2^{\mathbb{N}})^{\mathbb{N}})^n \rightarrow ((2^{\mathbb{N}})^{\mathbb{N}})^n$ by $f = \gamma_0 \times \gamma^{n \setminus \{0\}}$. That is,

$$f((\xi_i : i < n)) = (\gamma_0(\xi_0), \gamma(\xi_i) : 0 < i < n).$$

Remark 4.11. For a generic choice of β_0 and α ,

- f sends X_n to X_n^* , the domains of F_n and F_n^* respectively.
- f is a continuous reduction of F_n to F_n^* .

5. COMPLEXITY ON COMEAGER SETS

In this section we prove the following corollary of the main theorem (see Section 1.3). This will illustrate the construction of the map f , and the techniques developed so far.

Theorem 5.1. For $1 \leq n \leq \omega$, F_n retains its complexity on comeager sets. That is, $F_n \leq_B F_n \upharpoonright C$ for any comeager $C \subseteq X_n$.

Recall that our main construction is useful to deal with injective and separated sequences (see Lemma 4.4). The following lemma gives a reduction to the case of injective and separated sequences. Its proof is deferred to Section 5.1.

Lemma 5.2. There is a Borel reduction $\psi: F_n \rightarrow_B F_n$ so that for any $x \in X_n$, if $x \in \text{Image}(\psi)$ then $x(i) \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective for $i < n$ and separated for $0 < i < n$.

As discussed in Section 4.2, it suffices to prove that F_n^* retains its complexity on comeager sets. Fix a comeager $C \subseteq X_n^*$. We will find a Borel reduction f from F_n to F_n^* sending injective and separated sequences into C . This will show that $f \circ \psi$ is a reduction from F_n to $F_n^* \upharpoonright C$.

Proposition 5.3. For a generic choice of β_0 and α , if $x \in X_n$ is such that $x(i) \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective for all $i < n$ and separated for $0 < i < n$, then

$$(\forall^* g \in \text{Sym}(N)^n) g \cdot f(x) \in C.$$

Proof. We prove by induction on $0 \leq k < n$ that

$$(\star_k) \quad \forall^* h \in \text{Sym}(N)^{k+1} \text{ the fiber } C_{h \cdot f(x)|_{k+1}} \text{ is comeager in } ((2^N)^N)^{n \setminus k+1}.$$

First, assume that (\star_k) holds for $0 < k, k+1 < n$, and prove that (\star_{k+1}) holds.

Claim 5.4. $\forall^* h = (h_0, \dots, h_k) \in \text{Sym}(N)^{k+1}$, the pair

$$((h_{k-1}, h_k) \cdot f(x)(k), (h_k, \text{id}) \cdot f(x)(k+1)) \in (2^N)^N \times (2^N)^N$$

satisfies the conditions of Lemma 3.1, with $S = N$, $X = N$ and $Y_1 = N$, with respect to the comeager set

$$\{(a, b) \in (2^N)^N \times (2^N)^N : \text{the fiber } D = C_{(h|_k) \cdot (f(x)|_k), a, b} \subseteq ((2^N)^N)^{n \setminus k+2} \text{ is comeager}\}.$$

Proof of the claim. By assumption (\star_k) , $\forall^* h = (h_0, \dots, h_k)$, the set $D_{(h_{k-1}, h_k) \cdot f(x)(k)} \subseteq (2^N)^N$ is comeager, for D as above. It follows that $\forall^* h = (h_0, \dots, h_k)$, the set $D_{(h_{k-1}, \pi \circ h_k) \cdot f(x)(k)} \subseteq (2^N)^N$ is comeager, for any finite permutation π of N . This concludes condition (1) in Lemma 3.1.

Recall that $f(x)(k) = \gamma(x(k))$ and $f(x)(k+1) = \gamma(x(k+1))$. Since $x(k)$ is separated and $x(k+1)$ is injective and separated, it follows from Lemma 4.4 that the pair $(f(x)(k), f(x)(k+1))$ satisfies condition (2) in Lemma 3.1. Note that both parts (a) and (b) of condition (2) are invariant under the group action and so true for the pair

$$((h_{k-1}, h_k) \cdot f(x)(k), (h_k, \text{id}) \cdot f(x)(k+1))$$

as well. This concludes the proof of the claim. \square

We conclude from Lemma 4.4 that

$$(\forall^* (h_0, \dots, h_k) \in \text{Sym}(N)^{k+1}) (\forall^* (h'_k, h_{k+1}) \in \text{Sym}(N)^2) \text{ the fiber}$$

$C_{(h_0, \dots, h_{k-1}) \cdot f(x)|_k, (h_{k-1}, h'_k \cdot h_k) \cdot f(x)(k), (h'_k \cdot h_k, h_{k+1}) \cdot f(x)(k+1)}$ is comeager in $((2^N)^N)^{n \setminus k+2}$, and therefore

$$(\forall^* h \in \text{Sym}(N)^{k+2}) \text{ the fiber } C_{h \cdot f(x)|_{k+2}} \text{ is comeager in } ((2^N)^N)^{n \setminus k+2}.$$

Finally, we prove the base case of the induction. Let C_1 be the set of all $x \in (2^N)^N$ so that the fiber $C_x \subseteq ((2^N)^N)^{n \setminus 1}$ is comeager. Then $C_1 \subseteq (2^N)^N$ is comeager. Recall that $f(x)(0) = \gamma_0(x(0))$, and $x(0) \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective by assumption. By Lemma 4.7 and Lemma 3.2, for a generic choice of β_0 ,

$$(\forall^* h_0 \in \text{Sym}(N)) h_0 \cdot (f(x)(0)) \in C_1.$$

Let C_2 be the set of all $(a, b) \in (2^N)^N \times (2^N)^N$ so that the fiber $C_{a,b} \subseteq ((2^N)^N)^{n \setminus 2}$ is comeager. We claim that, $\forall^* h_0 \in \text{Sym}(N)$, the pair

$$(h_0 \cdot f(x)(0), (h_0, \text{id}) \cdot f(x)(1))$$

satisfies the conditions of Lemma 3.1. For part (1), note that as $(C_2)_{h_0 \cdot f(x)(0)}$ is comeager $\forall^* h_0 \in \text{Sym}(N)$, then the same conclusion holds for $\pi \circ h_0$, $\forall^* h_0 \in \text{Sym}(N)$, for any finite permutation π of N . For part (2), the pair $(f(x)(0), f(x)(1))$ satisfies the conditions (a) and (b), for a generic choice of α and β_0 , by Lemma 4.8,

as $x(0)$ is injective and $x(1)$ is injective and separated. As before, since these conditions are invariant under the group action, they are also satisfied by the pair $(h_0 \cdot f(x)(0), (h_0, \text{id}) \cdot f(x)(1))$. We then conclude from Lemma 3.1 that

$$(\forall^* h_0 \in \text{Sym}(N))(\forall^*(h'_0, h_1) \in \text{Sym}(N)^2)(h'_0 \cdot h_0 \cdot f(x)(0), (h'_0 \cdot h_0, h_1) \cdot f(x)(1)) \in C_2,$$

and therefore

$$(\forall^*(h_0, h_1) \in \text{Sym}(N)^2)(h_0 \cdot f(x)(0), (h_0, h_1) \cdot f(x)(1)) \in C_2,$$

concluding that (\star_1) holds.

Finally, we conclude the proof of the proposition. In case $n < \omega$, we conclude at stage $n - 1$ that $(\forall^* h \in \text{Sym}(N)^n) [h \cdot f(x) \in C]$, as required. Assume now that $n = \omega$. Let $C_k \subseteq ((2^N)^N)^k$ be the set of all $a \in ((2^N)^N)^k$ for which the fiber $C_a \subseteq ((2^N)^N)^{\omega \setminus k}$ is comeager. We have that $\forall^* h \in \text{Sym}(N)^\omega$, for any $k < \omega$, $h \cdot f(x) \upharpoonright k \in C_k$. It follows that for any sequence of finite permutations $\pi \in \text{Sym}(N)^k$, $\forall^* h \in \text{Sym}(N)^\omega$, $(\pi \circ h) \cdot f(x) \upharpoonright k \in C_k$.

Note that the action $a: \text{Sym}(N) \curvearrowright (2^N)^N$ is generically ergodic, that is, there is a comeager subset of $(2^N)^N$ in which every orbit is dense. Similarly, the action $\text{Sym}(N)^\omega \curvearrowright ((2^N)^N)^\omega$ is generically ergodic, and for any $k < \omega$ the action $\text{Sym}(N)^{\omega \setminus k} \curvearrowright ((2^N)^N)^{\omega \setminus k}$ is generically ergodic. Note that the orbit of some $b \in ((2^N)^N)^{\omega \setminus k}$ is dense if and only if for every $k < l < \omega$ the orbit of b restricted to $((2^N)^N)^{[k, l]}$ is dense. By thinning out the comeager set C , we may assume that if $\xi \in ((2^N)^N)^\omega$ is such that $\xi \upharpoonright k \in C_k$ for all $k < \omega$, then the orbit of $\xi \upharpoonright [k, \omega)$ is dense for $\text{Sym}(N)^{\omega \setminus k} \curvearrowright ((2^N)^N)^{\omega \setminus k}$, for every $k < \omega$.

Claim 5.5. Assume that $\xi \in ((2^N)^N)^\omega$ is such that $\pi \cdot (\xi \upharpoonright k) \in C_k$, for any sequence of finite permutations $\pi \in \text{Sym}(N)^k$, for all $k < \omega$. Then

$$(\forall^* g \in \text{Sym}(N)^\omega) g \cdot \xi \in C.$$

Proof. It suffices to prove that for any dense open set $C \subseteq D \subseteq ((2^N)^N)^\omega$, the set $\{g \in \text{Sym}(N)^\omega : g \cdot \xi \in D\}$ is dense open in $\text{Sym}(N)^\omega$. Fix a sequence of finite permutations $\pi \in \text{Sym}(N)^k$. We need to find an extension of it, $g \in \text{Sym}(N)^\omega$, so that $g \cdot \xi \in D$. Since $(\pi, \text{id}) \cdot (\xi \upharpoonright [k+1, \omega)) \in C_{k+1}$, the fiber $D_{(\pi, \text{id}) \cdot (\xi \upharpoonright [k+1, \omega))} \subseteq ((2^N)^N)^{\omega \setminus [k+1, \omega)}$ is dense open. Since the orbit of $\xi \upharpoonright [k+1, \omega)$ is dense, we may find some $h \in \text{Sym}(N)^{\omega \setminus [k+1, \omega)}$ so that $h \cdot \xi \upharpoonright [k+1, \omega) \in D_{(\pi, \text{id}) \cdot (\xi \upharpoonright [k+1, \omega))}$. Then $g = (\pi, \text{id}, h)$ is the desired extension of π so that $g \cdot \xi \in D$. \square

It follows now that

$$(\forall^* h \in \text{Sym}(N)^\omega)(\forall^* g \in \text{Sym}(N)^\omega) g \cdot (h \cdot f(x)) \in C,$$

and so $(\forall^* h \in \text{Sym}(N)^\omega) [h \cdot f(x) \in C]$, as required. \square

We now fix sufficiently generic α and β_0 , so that the conclusion of the proposition holds, and so that f is a reduction from F_n to F_n^* . For any $x \in (2^{\mathbb{N}})^{\mathbb{N}}$, $(\forall^* g \in \text{Sym}(N)^n) [g \cdot f \circ \psi(x) \in C]$. We may now construct the reduction using a large section uniformization theorem. By [Kec95, 18.6] there is a Borel map $\rho: X_n \rightarrow$

$\text{Sym}(N)^n$ so that $\rho(x) \cdot f \circ \psi(x) \in C$ for any $x \in X_n$. Since ψ and f are reductions, and F_n^* is invariant under the action, we conclude that

$$x \mapsto \rho(x) \cdot f \circ \psi(x)$$

is a Borel reduction of F_n to $F_n^* \upharpoonright C$.

5.1. Different base sets for F_n . Recall the definition of the equivalence relation F_n on the domain $X_n \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^n$ in Section 1.1. Given a sequence of countable infinite sets $\vec{N} = (N_i : i < n)$, define analogously an equivalence relation $F_n(\vec{N})$ on a domain $X_n(\vec{N}) \subseteq \prod_{i < n} (2^{N_i})^{N_{i+1}}$. Fix bijections $\pi_i : \mathbb{N} \rightarrow N_i$. These lead naturally to a homeomorphism

$$\prod_{i < n} (2^{N_i})^{N_{i+1}} \rightarrow \prod_{i < n} (2^{\mathbb{N}})^{\mathbb{N}} = ((2^{\mathbb{N}})^{\mathbb{N}})^n.$$

Define $X_n(\vec{N})$ to be the preimage of X_n , and $F_n(\vec{N})$ to be the pullback of F_n .

Equivalently, these objects can be defined directly as in Section 1.1. For example, given $x \in \prod_{i < n} (2^{N_i})^{N_{i+1}}$, we may define $A_1^x = \{x(0)(t) : t \in N_1\}$, a countable subset of 2^{N_0} , $A_2^x = \{a_1^{x,s} : s \in N_2\}$, where $a_1^{x,s} = \{x(0)(t) : x(1)(s)(t) = 1\} \subseteq A_1^x$, and define A_3^x, \dots, A_n^x analogously. We then have that $x F_n(\vec{N}) y$ if and only if $A_{k+1}^x = A_{k+1}^y$ for all $k < n$.

5.2. Proof of Lemma 5.2. We will construct a Borel reduction $\psi : F_n \rightarrow F_n$ so that for any $x \in \text{Image}(\psi)$, $x(i) \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective for $i < n$ and separated for $0 < i < n$.

First we describe the map in terms of the classifying invariants A_n^x associated to $x \in X_n$. Note that if A_1^x, \dots, A_n^x are all infinite, then there is some $x' \in X_n$ which is injective and is F_n -equivalent to x (that is, $A_i^x = A_i^{x'}$ for $1 \leq i \leq n$). For an injective $x \in X_n$, the condition of being separated corresponds to: for any $1 \leq i < n$, for any $u \neq v \in A_i$ there is some $Z \in A_{i+1}$ so that Z contains exactly one of $\{u, v\}$.

Given some $A_1, \dots, A_n = A_1^x, \dots, A_n^x$ we construct new sets B_1, \dots, B_n , which will correspond to A_1^y, \dots, A_n^y for some $y \in X_n$ which will be defined as $y = \psi(x)$. We may assume that there are infinitely many reals $S = \{\star_1, \star_2, \dots\}$ which are not in A_1^x for any x . Define inductively

- $B_1 = A_1 \cup S$;
- $B_{k+1} = A_{k+1} \cup \{\{a, b\} : a \in A_k, b \in B_k \setminus A_k\}$.

Note that $B_i \setminus A_i$ is infinite for $i = 1, \dots, n$. For $1 \leq i < n$, given $u, v \in B_k$, we find some set $Z \in B_{k+1}$ separating them. If u, v are both in A_k , let $a \in B_k \setminus A_k$, then $\{u, a\} \in B_{k+1}$ separates $\{u, v\}$. If u, v are both in $B_k \setminus A_k$, let $a \in A_k$, then $\{u, a\} \in B_{k+1}$ separates $\{u, v\}$. If $u \in A_k$ and $v \in B_k \setminus A_k$, let $a \in B_k \setminus A_k$ be different than v , then $\{u, a\} \in B_{k+1}$ separates $\{u, v\}$.

Finally, note that the sets (A_1, \dots, A_n) are uniquely defined from (B_1, \dots, B_n) , where the members of A_i are the members of B_i which do not include any member of

S in their transitive closure. This fact corresponds to the map we are constructing being a reduction.

Remark 5.6. We will later use the following fact. Suppose x, x' are such that $A_i^x = A_i^{x'}$ for $i = 1, \dots, k, k < n$. Let B'_i be the result of the construction applied to x' . Then $B_k = B'_k$, and for any distinct $u, v \in B_k$ there is Z which is both in B_{k+1} and B'_{k+1} so that Z separates $\{u, v\}$.

It is left to find a Borel map sending x to y so that A_i^y are as B_i to A_i^x as above. Given $x \in X_n$, we define

$$\psi_0(x) = y = (y(k) : k < n),$$

so that $y(0) \in (2^{\mathbb{N}})^{\mathbb{N} \sqcup \mathbb{N}}$, and $y(k) \in (2^{\mathbb{N} \sqcup \mathbb{N}^k})^{\mathbb{N} \sqcup \mathbb{N}^{k+1}}$ for $k \geq 1$, as follows. First we define $y(0)$. For $n \in \mathbb{N}$ (in the left copy of $\mathbb{N} \sqcup \mathbb{N}$), $y(0)(n) = x(0)(n)$. For $n \in \mathbb{N}$ (in the right copy of $\mathbb{N} \sqcup \mathbb{N}$), $y(0)(n) = \star_n$. That is, $y(0)$ comprises of two sequences, one is x and the other is the sequence of new reals S as above.

Given $y(k) \in (2^{\mathbb{N} \sqcup \mathbb{N}^k})^{\mathbb{N} \sqcup \mathbb{N}^{k+1}}$, define $y(k+1) \in (2^{\mathbb{N} \sqcup \mathbb{N}^{k+1}})^{\mathbb{N} \sqcup \mathbb{N}^{k+2}}$:

- for $n \in \mathbb{N}$, for $m \in \mathbb{N}$, $y(k+1)(n)(m) = x(n)(m)$, and for $t \in \mathbb{N}^{k+1}$, $y(k+1)(n)(t) = 0$.
- for $(m, t) \in \mathbb{N}^{k+2} = \mathbb{N} \times \mathbb{N}^{k+1}$, $y(k+1)((m, t))(a) = 1$ if and only if $a = m \in \mathbb{N}$ or $a = t \in \mathbb{N}^{k+1}$.

Define $\vec{N} = (N_i : i < n)$ by $N_0 = \mathbb{N} \sqcup \mathbb{N}$, $N_k = \mathbb{N} \sqcup \mathbb{N}^{k+1}$. Then ψ_0 is a Borel reduction of F_n to $F_n(\vec{N})$ so that if $y = \psi_0(x)$ then the sets A_1^y, A_2^y, \dots are constructed from A_1^x, A_2^x, \dots as above.

Let π be the homeomorphism which is a reduction of $F_n(\vec{N})$ to F_n . If $y = \pi \circ \psi_0(x)$ then the sets A_1^y, A_2^y, \dots are infinite and separated. The only issue at this point is that $y(k)$ may be a non-injective enumeration of the set A_k^y . We invoke a cleanup function eliminating multiplicities, to end up with injective sequences. Recall from Section 2.1 the Borel map $C : (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}}$ so that for $x \in (2^{\mathbb{N}})^{\mathbb{N}}$,

- if the set enumerated by x is finite, $C(x) = x$,
- if the set enumerated by x is infinite, $C(x)$ is an injective enumeration of the same set.

Another property of this map is that $C(x)$ does not depend on the reals appearing in the sequence x , but only on whether $x(n), x(m)$ are equal, for $n, m \in \mathbb{N}$.

Next, extend this to a map $\hat{C}_0 : (2^{\mathbb{N}})^{\mathbb{N}} \times 2^{\mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}} \times 2^{\mathbb{N}}$ which, after applying C to the first coordinate, corrects the second coordinate to carve out the same subset. That is, given $(x, v) \in (2^{\mathbb{N}})^{\mathbb{N}} \times 2^{\mathbb{N}}$ and $\hat{C}_0(x, v) = (x', v')$, $\{x(i) : v(i) = 1\} = \{x'(i) : v'(i) = 1\}$. This in turn extends to a map

$$\hat{C} : (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}},$$

defined so that if $\hat{C}(x, y) = (x', y')$, then $\hat{C}_0(x, y(i)) = (x', y'(i))$, for all $i \in \mathbb{N}$. For $k+1 < n$, define $C_k : ((2^{\mathbb{N}})^{\mathbb{N}})^n \rightarrow ((2^{\mathbb{N}})^{\mathbb{N}})^n$ by

$$C_k(x(0), \dots, x(k), x(k+1), x(k+2), \dots) = (x(0), \dots, \hat{C}(x(k), x(k+1)), x(k+2), \dots).$$

If $n < \omega$ define $C_{n-1}(x(0), \dots, x(n-1)) = (x(0), \dots, C(x_{n-1}))$. Note that the maps C_k , $k < n$, commute with one another on the domain X_n . Finally, define

$$C^*: ((2^{\mathbb{N}})^{\mathbb{N}})^n \rightarrow ((2^{\mathbb{N}})^{\mathbb{N}})^n \text{ by } C^* = C_0 \circ C_1 \circ \dots$$

Note that C^* is well defined also in the case $n = \omega$, as the n 'th coordinate of $C^*(x)$ is fixed by $C_{n+1} \circ C_{n+2} \circ \dots$. The map

$$\psi = C^* \circ \pi \circ \psi_0: X_n \rightarrow X_n$$

is now the desired reduction, concluding the proof of Lemma 5.2. \square

6. SOME IDEAS AND OBSTACLES TOWARDS THE MAIN THEOREM

We begin working towards a proof of Theorem 1.5. First we make a slight reformulation. Then we briefly sketch the ideas for the case $n = 1$, and emphasize some difficulties for extending these to $n \geq 2$.

6.1. A reformulation. We will prove Theorem 1.5 in the following equivalent formulation. For equivalence relations F and E on the same domain, say that E **extends** F if $F \subseteq E$. For $n < m \leq \omega$ we may view F_n as an equivalence relation on X_m , defined by $x F_n y \iff x \upharpoonright n F_n y \upharpoonright n$, for $x, y \in X_m$. In this case F_n extends F_m , for $n < m$.

Theorem 6.1. Fix $1 \leq n \leq \omega$. For any equivalence relation E , classifiable by countable structures, which extends F_n , either

- F_n is Borel reducible to E , or
- For some $k < n$, E extends F_k on a comeager subset of X_n .

Given a Borel homomorphism $f: F \rightarrow_B E$, define E^* , on the same domain as F , as the pullback of E : $x E^* y \iff f(x) E f(y)$. Then (1) E^* extends F , and (2) E^* is Borel reducible to E (witnessed by f). The definition above, of F_n as an equivalence relation on X_m , is the pullback of F_n by the homomorphism u_n^m .

Proof of Theorem 1.5 from Theorem 6.1. Given an analytic equivalence relation E and a Borel homomorphism $f: F_n \rightarrow_B E$, as in Theorem 1.5, apply Theorem 6.1 to E^* . In the first case, we conclude that F_n is Borel reducible to E^* , and is therefore Borel reducible to E as well. Otherwise, there is $k < n$ and a comeager $C \subseteq X_n$ on which E^* extends F_k . We may find a Borel partial function $g^*: X_k \rightarrow X_n$, defined on a comeager subset of X_k , so that $g^*(x)$ extends x , and $g^*(x) \in C$, for any x in the domain of g^* . In particular, g^* is a homomorphism from F_k to E^* . Define $g = f \circ g^*$. Then $g: F_k \rightarrow_B E$ is a Borel homomorphism.

Finally, we see that f factors through u_k^n , via g , on the comeager set C . Given $x \in C$, since E^* extends F_k on C , x and $g^* \circ u_k^n(x)$ are E^* -related, and so $f(x)$ and $f(g^* \circ u_k^n(x)) = g \circ u_k^n(x)$ are E -related. \square

6.2. **The case $n = 1$.** Fix an analytic equivalence relation E on $(2^{\mathbb{N}})^{\mathbb{N}}$ so that $F_1 \subseteq E$. Recall that F_0 is a trivial equivalence relation, with just one class, so the second clause in Theorem 6.1, stating that E extends F_0 on a comeager set, states that E has a comeager equivalence class. Assume that E does not have a comeager class. [KSZ13, Theorem 6.24] then proves that F_1 is Borel reducible to E as follows. Recall the homomorphism $f: F_1 \rightarrow_B F_1$ defined in Section 2.1. We claim that it reduces F_1 to E . Since E extends F_1 , it remains to show that $x \not F_1 y \implies f(x) \not E f(y)$. Note that there are three different ways for $x, y \in (2^{\mathbb{N}})^{\mathbb{N}}$ to be not F_1 -related:

- a. x and y enumerate disjoint sets;
- b. one of the two sets is contained in the other;
- c. non of the above.

For example, if $x \not F_1 y$ as in case a. above, one can show that then $(f(x), f(y)) \in (E^c)^* \subseteq E^c$, and so $f(x) \not E f(y)$, as required.

Notation 6.2. For $a, b \in (2^{\mathbb{N}})^{\mathbb{N}}$, write $a \cup b$ for some member of $(2^{\mathbb{N}})^{\mathbb{N}}$ enumeration the union of the sets enumerated by a and b . We will ask questions about whether such sequence is equivalent to another, according to F_1 or E . As both extend F_1 , the answer does not depend on the enumeration of $a \cup b$. We will similarly use the notations $a \setminus b$ and $a \cap b$, whenever these are not empty, for some member of $(2^{\mathbb{N}})^{\mathbb{N}}$ enumerating the corresponding sets.

Whenever $a \cup b$, $a \cap b$, or $a \setminus b$ are infinite, **we always take the notation to be an injective enumeration** of the corresponding set. One important aspect of the definition of f in Section 2.1 is that for any a, b in the image of f , both are infinite, and the sets $a \cap b$ and $a \setminus b$ are either empty or infinite.

Assume now that $x \not F_1 y$ according to case c. Following the definition of f , we may write $f(x)$ and $f(y)$ as $a \cup c$ and $b \cup c$, where $a = f(x \setminus y)$, $b = f(y \setminus x)$, and $c = f(x \cap y)$.

It follows from E being meager that $\forall^*(a, b, c) \in (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}} [a \cup c \not E b \cup c]$. Let $C \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^3$ be the corresponding comeager set. As before, for a ‘‘sufficiently generic’’ choice of the function $g: 2^{\mathbb{N}} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}}$ in Section 2.1 it can be verified that for disjoint non-empty z_1, z_2, z_3 , for almost all $(g_1, g_2, g_3) \in (S_{\infty})^3$, $(g_1 \cdot f(z_1), g_2 \cdot f(z_2), g_3 \cdot f(z_3)) \in C$. Applying this to the disjoint non-empty sets $x \setminus y, y \setminus x, x \cap y$, it then follows that $f(x) \not E f(y)$, as required.

6.3. **The case $n \geq 2$.** Our efforts so far were to find a Borel homomorphism $f: F_n \rightarrow_B F_n$, landing in comeager sets (after a Vaught transform). Using this homomorphism we will be able to extend the ideas in Section 6.2 to prove the following: if $x \not F_n y$ differ *only at the last coordinate* (so their restrictions to X_{n-1} are F_{n-1} -equivalent), then $f(x) \not E f(y)$. See Lemma 7.3.

A new difficulty arising in the $n \geq 2$ case is dealing with $x \not F_n y$ which are already F_k -inequivalent for some $k < n$. The issue with trying to extend these arguments directly is a proliferation of cases to consider. For example, suppose $x(0)$ and $y(0)$

are already F_1 -inequivalent. There are three cases a., b., and c. as above. Consider case c., so we have that $x(0) \cap y(0), x(0) \setminus y(0), y(0) \setminus x(0)$ are not empty. We would now need to view each set in $x(1)$, which is considered a subset of $x(0)$, as a union of two sets, a subset of $x(0) \cap y(0)$ and a subset of $x(0) \setminus y(0)$. Given two different members of $x(1)$, we would need to worry about whether they may agree on either their restrictions to $x(0) \cap y(0)$ or $x(0) \setminus y(0)$. Similarly, we would have to keep track on which members of $x(1)$ and $y(1)$ agree on the intersection $x(0) \cap y(0)$. While this may be handled for $n = 2$, when $n \gg 2$ there are ever more cases to consider and divisions to keep track of.

The solution will be to “decompose” the equivalence relation E (which extends F_n) to a sequence of equivalence relations E_k so that E_k extends both F_k and E . While this is not generally possible, it is possible generically. This is the content of Lemma 7.4.

The proof of Lemma 7.3 relies on the Baire-categoric techniques developed above, and will be a natural extension of the arguments in Section 5. The proof of Lemma 7.4 will involve higher set theoretic techniques as well.

7. PROOF OF THE MAIN THEOREM

We now prove the main result, Theorem 1.5, in its equivalent formulation, Theorem 6.1. As above, it will be convenient to work with F_n^* instead of F_n (see Section 4.2). We prove the following, which is equivalent to Theorem 6.1.

Theorem 7.1. Fix $1 \leq n \leq \omega$. For any E which is classifiable by countable structures and which extends F_n^* , either

- F_n is Borel reducible to E , or
- For some $k < n$, E extends F_k^* on a comeager subset of X_n^* .

Towards that end, fix $1 \leq n \leq \omega$, and E as above which extends F_n^* . Assume that E does not extend F_k^* on a comeager set, for any $k < n$. We must prove that F_n is Borel reducible to E . Recall the definition of $f: F_n \rightarrow_B F_n$, Definition 4.10 in Section 4, and the definition of $\psi: F_n \rightarrow F_n$ from Lemma 5.2. Since E extends F_n^* , $f \circ \psi: F_n \rightarrow_B E$ is a homomorphism.

Claim 7.2. There are maps α and β_0 so that $f \circ \psi$ is a reduction of F_n to E .

Towards that end, fix $x, y \in X_n$ in the image of ψ so that $x \not\#_n y$. We need to prove that $f(x) \not\# f(y)$. We will split into countably many cases. In each case we will show that $f(x) \not\# f(y)$ for generically chosen α and β_0 . This will conclude the proof of the claim.

Lemma 7.3. Fix $0 \leq k < \omega$. Let E_{k+1} be an analytic equivalence relation, defined on a comeager subset of X_{k+1}^* , extending F_{k+1}^* on this domain. Assume that E_{k+1} does not extend F_k^* on any comeager set. Let $x, y \in X_{k+1}$ be in the image of ψ . Assume that $x \not\#_{k+1} y$ yet $(x \upharpoonright k) \#_k (y \upharpoonright k)$. Then $f_{k+1}(x) \not\#_{k+1} f_{k+1}(y)$, where $f_k: F_k \rightarrow_B F_k$ is the homomorphism $\gamma_0 \times \gamma^{k \setminus \{0\}}$ as in Definition 4.10.

Lemma 7.4. There are analytic equivalence relations E_k for $k < n$, defined on comeager subsets of X_k^* , so that

- (1) E_k extends F_k^* , on a comeager set;
- (2) E_{k+1} does not extend F_k^* on any comeager set;
- (3) $E \subseteq E_k$, on a comeager set, for each $k < n$. That is, on a comeager set, if $x E y$ then $x \upharpoonright k E_k y \upharpoonright k$.

In fact we get the following picture, on comeager sets, with $E_n = E$:

$$\begin{array}{ccccccc} F_1^* & \supseteq & F_2^* & \supseteq & F_3^* & \supseteq & \dots & \supseteq & F_n^* \\ \cap & & \cap & & \cap & & & & \cap \\ E_1 & \supseteq & E_2 & \supseteq & E_3 & \supseteq & \dots & \supseteq & E_n \end{array}$$

Remark 7.5. The lemma is equivalent to Lemma 1.8. See Section 6.1.

Finally, the proof of the main theorem terminates as follows. Let k be minimal so that $x \upharpoonright k \not E_k y \upharpoonright k$. It follows from Lemma 7.3, and that $f_k(x \upharpoonright k) = f_n(x) \upharpoonright k$, that $f(x) \upharpoonright k \not E_k f(y) \upharpoonright k$, and therefore $f(x) \not E f(y)$, as required.

We note that, without the assumption that E is classifiable by countable structures, the proof works if we assume that E can be decomposed as in Lemma 7.4 above. In the terminology of Theorem 1.5, we get the following variation.

Theorem 7.6. Given analytic equivalence relations E_k for $k \leq n$ and a diagram of Borel homomorphisms which commute on comeager sets as in Lemma 1.8, so that f_k does not factor through u_l^k for $l < k \leq n$, then $F_n \leq_B E_n$.

We conclude the paper by proving Lemma 7.3 and Lemma 7.4.

7.1. Proof of Lemma 7.3. For this subsection we fix $f = f_{k+1}: F_{k+1} \rightarrow F_{k+1}$. Since f is a homomorphism and $x \upharpoonright k E_k y \upharpoonright k$, we may assume that $x \upharpoonright k = y \upharpoonright k$. Now $x(k), y(k) \in (2^{\mathbb{N}})^{\mathbb{N}}$ are F_1 -inequivalent. There are three options:

- a. The two subsets of $2^{\mathbb{N}}$ enumerated by $x(k), y(k)$ are disjoint;
- b. one of the two sets is contained in the other;
- c. neither of the above.

When we use set notation, such as $x(k) \cap y(k)$, or $x(k) \setminus y(k)$, we refer to the sets enumerated by $x(k), y(k)$, respectively. We assume that $k \geq 1$. For $k = 0$ the arguments are similar to Section 6.2.

Case a. Assume that $x(k)$ and $y(k)$ enumerate disjoint subsets of $2^{\mathbb{N}}$. Since $x \upharpoonright k = y \upharpoonright k$, and by the definition of f , we may write $f(x) = (a, r)$ and $f(y) = (a, s)$, where $a \in ((2^{\mathbb{N}})^{\mathbb{N}})^k$ and $r, s \in (2^{\mathbb{N}})^{\mathbb{N}}$. We view (a, r, s) as a member of the space $((2^{\mathbb{N}})^{\mathbb{N}})^k \times (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$. The following is a consequence of our assumption that E_{k+1} does not extend F_k on any comeager set.

Lemma 7.7. $\forall^*(x, y, z) \in ((2^{\mathbb{N}})^{\mathbb{N}})^k \times (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}} [(x, y) \not E_{k+1} (x, z)]$.

Proof. Given $x \in ((2^N)^N)^k$, consider the equivalence relation E_x on $(2^N)^N$, $y E_x z \iff (x, y) E_{k+1} (x, z)$. The lemma is equivalent to the statement: for almost every $x \in ((2^N)^N)^k$ every E_x class is meager. If this fails, then for almost every $x \in ((2^N)^N)^k$ there is a comeager equivalence class in E_x . In this case, we conclude that E_{k+1} extends F_k on the comeager set of all $(x, y) \in ((2^N)^N)^k \times (2^N)^N$ so that y is in the comeager equivalence class of E_x . \square

Cosmetic modifications. As in Section 1.2, there is a natural action

$$\text{Sym}(N)^k \times \text{Sym}(N) \times \text{Sym}(N) \curvearrowright ((2^N)^N)^k \times (2^N)^N \times (2^N)^N.$$

More specifically, expressing the group as

$$\text{Sym}(N)^{k-1} \times \text{Sym}(N) \times \text{Sym}(N) \times \text{Sym}(N)$$

and the space as

$$((2^N)^N)^{k-1} \times (2^N)^N \times (2^N)^N \times (2^N)^N,$$

the blue copy of $\text{Sym}(N)$ acts on the three blue copies of N diagonally, and the two copies of $\text{Sym}(N)$ act separately on the two copies of $(2^N)^N$, as in Section 1.2. The point is that the two projections $((2^N)^N)^k \times (2^N)^N \times (2^N)^N \rightarrow ((2^N)^N)^{k+1}$ (the maps $(a, b, c) \mapsto (a, b)$ and $(a, b, c) \mapsto (a, c)$) are equivariant.

Claim 7.8. Given a comeager set $C \subseteq ((2^N)^N)^k \times (2^N)^N \times (2^N)^N$, for almost any β_0 and α ,

$$\forall^*(g, h_1, h_2) \in \text{Sym}(N)^k \times \text{Sym}(N) \times \text{Sym}(N) [(g, h_1, h_2) \cdot (a, r, s) \in C].$$

Applying this claim to the comeager set C we get from Lemma 7.7, we conclude that for some (g, h_1, h_2) , $(g, h_1) \cdot (a, r) \notin E_{k+1} (g, h_2) \cdot (a, s)$, and so $f(x) \notin E_{k+1} f(y)$, as required for Lemma 7.3. We finish Case a. by proving the claim.

Proof of the claim. We will use the following variation of Lemma 4.4, in the case that $k > 1$.

Lemma 7.9. For a generic α the following holds. Suppose $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ is separated and $y_1, y_2 \in (2^{\mathbb{N}})^{\mathbb{N}}$ are injective and separated. Assume further that y_1, y_2 are disjoint. Then the triplet $(\zeta, \xi_1, \xi_2) = (\gamma(x), \gamma(y_1), \gamma(y_2)) \in (2^S)^X \times (2^X)^{Y_1} \times (2^X)^{Y_2}$, where $S = X = Y_1 = Y_2 = N$, satisfies the conditions in part (2) of Lemma 3.1.

If $k = 1$, the following variation of Lemma 4.8 will be used:

Lemma 7.10. For generic α and β_0 the following holds. Suppose $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ is injective, and $y_1, y_2 \in (2^{\mathbb{N}})^{\mathbb{N}}$ are injective and separated. Then the triplet $(\zeta, \xi_1, \xi_2) = (\gamma_0(x), \gamma(y_1), \gamma(y_2)) \in (2^S)^X \times (2^X)^{Y_1} \times (2^X)^{Y_2}$, where $S = X = Y_1 = Y_2 = N$, satisfies the conditions in part (2) of Lemma 3.1.

As in Claim 5.4 we get: $\forall^*(h_0, \dots, h_{k-1}) \in \text{Sym}(N)^k$ the triplet

$$((h_{k-2}, h_{k-1}) \cdot f(x(k-1)), (h_{k-1}, \text{id}) \cdot f(x(k)), (h_{k-1}, \text{id}) \cdot f(y(k))) \in (2^N)^N \times (2^N)^N \times (2^N)^N$$

satisfies the conditions of Lemma 3.1 with $k = 2$, $S = X = Y_1 = Y_2 = N$, with respect to the comeager set

$$\{(a, b, c) \in ((2^N)^N)^3 : ((h \upharpoonright k - 1) \cdot (f(x) \upharpoonright k - 1), a, b, c) \in C\}$$

We conclude from Lemma 3.1 that

$$(\forall^*(h_0, \dots, h_{k-1}) \in \text{Sym}(N)^k)(\forall^*(h'_{k-1}, h_k^1, h_k^2) \in \text{Sym}(N)^3)$$

$$((h \upharpoonright k - 1) \cdot (f(x) \upharpoonright k - 1), (h_{k-2}, h'_{k-1} \cdot h_{k-1}) \cdot f(x(k-1)), (h'_{k-1} \cdot h_{k-1}, h_k^1) \cdot f(x(k)), (h'_{k-1} \cdot h_{k-1}, h_k^2) \cdot f(y(k)))$$

is in C , and therefore $\forall^*(h_0, \dots, h_{k-1}, h_k^1, h_k^2) \in \text{Sym}(N)^k \times \text{Sym}(N) \times \text{Sym}(N)$

$$((h \upharpoonright k - 1) \cdot (f(x) \upharpoonright k - 1), (h_{k-2}, h_{k-1}) \cdot f(x(k-1)), (h_{k-1}, h_k^1) \cdot f(x(k)), (h_{k-1}, h_k^2) \cdot f(y(k))) \in C,$$

concluding the proof of the claim. \square

We remark that if $k = 0$, the proof is similar, using the following variation of Lemma 4.7.

Lemma 7.11. Suppose $x, y \in (2^{\mathbb{N}})^{\mathbb{N}}$ are injective enumerations of two disjoint subsets of $2^{\mathbb{N}}$. Then for any comeager $C \subseteq (2^N)^N \times (2^N)^N$, for a generic β_0 , the pair $(\gamma_0(x), \gamma_0(y)) \in (2^N)^N \times (2^N)^N$ satisfies the assumption in Lemma 3.2, with $X = N$, $Y_1 = Y_2 = N$, and $M_1 = M_2 = \mathbb{N}$.

Case b. Assume that $x(k)$ enumerates a subsets of $y(k)$. We skip the details of this case, as they are similar and slightly simpler than Case c.

Case c. Assume that $x(k)$ and $y(k)$ enumerate two sets so that $x(k) \cap y(k)$, $x(k) \setminus y(k)$, and $y(k) \setminus x(k)$ are not empty. As before we focus on the case $k \geq 1$. We may assume that any member of 2^N which appears both in $x(k)$ and $y(k)$ appears in the same coordinate $x(k)(i), y(k)(i)$. Recall the definition of $\gamma(x(k)) \in (2^N)^N$. We may identify it with a member of $(2^N)^{S_1 \sqcup S_2}$, where S_1 is the set of all $(t, k) \in N$ so that the image of t is contained in $x(k) \cap y(k)$, and $S_2 = N \setminus S_1$. Note that both S_1 and S_2 are infinite. The space $(2^N)^{S_1 \sqcup S_2}$ is naturally identified with $(2^N)^{S_1} \times (2^N)^{S_2}$, giving a homeomorphism

$$\iota: ((2^N)^N)^k \times (2^N)^{S_1} \times (2^N)^{S_2} \rightarrow ((2^N)^N)^k \times (2^N)^N.$$

Note that we may write $f(x) = \iota(a, \xi_0, \xi_1)$ and $f(y) = \iota(a, \xi_0, \xi_2)$ for some $a \in ((2^N)^N)^k$, $\xi_0 \in (2^N)^{S_1}$, $\xi_1, \xi_2 \in (2^N)^{S_2}$. The following is a consequence of our assumption that E_{k+1} does not extend F_k on a comeager set.

Lemma 7.12. $\forall^*(a, \xi_0, \xi_1, \xi_2) \in ((2^N)^N)^k \times (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_2} [\iota(a, \xi_0, \xi_1) \notin E_{k+1} \iota(a, \xi_0, \xi_2)]$

Proof. Assume for contradiction that the statement fails. Since almost every E_{k+1} class is dense, and ι is a homeomorphism, it follows that

$$\forall^*(a, \xi_0, \xi_1, \xi_2) \in ((2^N)^N)^k \times (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_2} [\iota(a, \xi_0, \xi_1) \in E_{k+1} \iota(a, \xi_0, \xi_2)]$$

Given $a \in ((2^N)^N)^k$, consider the equivalence relation E_a on $(2^N)^{S_1} \times (2^N)^{S_2}$, $(\xi_0, \xi_1) E_a (\zeta_0, \zeta_1) \iff \iota(a, \xi_0, \xi_1) \in E_{k+1} \iota(a, \zeta_0, \zeta_1)$. Then for almost every $a \in ((2^N)^N)^k$, $\forall^*(\xi_0, \xi_1, \xi_2) \in (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_2} [(\xi_0, \xi_1) E_a (\xi_0, \xi_2)]$.

Fix a bijection $S_2 \rightarrow S_1$, giving a homeomorphism $s: (2^N)^{S_1} \rightarrow (2^N)^{S_2}$. Then

$$(\forall^* a \in ((2^N)^N)^k)(\forall^*(\xi_0, \xi_1, \xi_2) \in (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_1}) [(\xi_0, \xi_1) E_a (\xi_0, s(\xi_2))].$$

Note that for any a the map $(\zeta_0, \zeta_1) \mapsto (s^{-1}(\zeta_1), s(\zeta_0))$ is an E_a -invariant homeomorphism $(2^N)^{S_1} \times (2^N)^{S_2} \rightarrow (2^N)^{S_1} \times (2^N)^{S_2}$. We conclude that

$$(\forall^* a \in ((2^N)^N)^k)(\forall^*(\xi_0, \xi_1, \xi_2, \xi_3) \in (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_1} \times (2^N)^{S_2}) \\ (\xi_0, \xi_1) E_a (\xi_0, s(\xi_2)) E_a (\xi_2, s(\xi_0)) E_a (\xi_2, \xi_3)$$

That is, for almost every $a \in ((2^N)^N)^k$ there is a comeager equivalence class for E_a . As in Lemma 7.7 we conclude that E_{k+1} extends F_k on a comeager set, a contradiction. \square

Cosmetic modifications. As in Section 1.2, there is a natural action

$$\text{Sym}(N)^k \times \text{Sym}(S_1) \times \text{Sym}(S_2) \times \text{Sym}(S_2) \curvearrowright ((2^N)^N)^k \times (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_2}.$$

More specifically, expressing the group as

$$\text{Sym}(N)^{k-1} \times \text{Sym}(N) \times \text{Sym}(S_1) \times \text{Sym}(S_2) \times \text{Sym}(S_2)$$

and the space as

$$((2^N)^N)^{k-1} \times (2^N)^N \times (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_2},$$

the blue copy of $\text{Sym}(N)$ acts on the four blue copies of N diagonally, while the groups $\text{Sym}(S_1)$, $\text{Sym}(S_2)$, $\text{Sym}(S_2)$ act separately on the spaces $(2^N)^{S_1}$, $(2^N)^{S_2}$, $(2^N)^{S_2}$. The point is that the projection maps $((2^N)^N)^k \times (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_2} \rightarrow ((2^N)^N)^{k+1}$, $(a, b, c, d) \mapsto \iota(a, b, c)$ and $(a, b, c, d) \mapsto \iota(a, b, d)$, are equivariant.

The following holds for a generic choice of β_0 and α as in Section 4.

Claim 7.13. Given a comeager set

$$C \subseteq ((2^N)^N)^k \times (2^N)^{S_1} \times (2^N)^{S_2} \times (2^N)^{S_2},$$

$$\forall^*(g, \delta_0, \delta_1, \delta_2) \in \text{Sym}(N)^k \times \text{Sym}(S_1) \times \text{Sym}(S_2) \times \text{Sym}(S_2),$$

$$(g \cdot a, (g(k-1), \delta_0) \cdot \xi_0, (g(k-1), \delta_1) \cdot \xi_1, (g(k-1), \delta_1) \cdot \xi_2) \in C.$$

Applying this claim to the comeager set C we get from Lemma 7.12, we conclude that for some $(g, \delta_0, \delta_1, \delta_2)$,

$$\iota((g, \delta_0, \delta_1) \cdot (a, \xi_0, \xi_1)) \not\equiv_{k+1} \iota((g, \delta_0, \delta_2) \cdot (a, \xi_0, \xi_2)),$$

and so $\iota((a, \xi_0, \xi_1)) \not\equiv_{k+1} \iota((a, \xi_0, \xi_2))$, that is, $f(x) \not\equiv_{k+1} f(y)$, as required for Lemma 7.3.

The proof of the claim is similar to that in Case a., where the following variations of Lemma 4.4 and Lemma 4.8 are used. Recall Remark 5.6. Since x, y are in the image of ψ , then $x(k), y(k)$ are **jointly separated**, that is, for any distinct $n, k \in \mathbb{N}$ there is i in $x(k) \cap y(k)$ so that $x(k)(i) = y(k)(i)$ separates n, k .

Lemma 7.14. For generic α and β_0 the following holds. Fix $x \in (2^{\mathbb{N}})^{\mathbb{N}}$, and $y_1, y_2 \in (2^{\mathbb{N}})^{\mathbb{N}}$ so that the sets $y_1 \cap y_2$, $y_1 \setminus y_2$, $y_2 \setminus y_1$, are non-empty. Assume that y_1, y_2 are injective and jointly separated. Let $S_1 \subseteq N$ be the set of all (t, k) for which the image of t is contained in $y_1 \cap y_2$, $S_2 = N \setminus S_1$. Let $\xi_1 \in (2^N)^{S_1}$, $\xi_2, \xi_3 \in (2^N)^{S_2}$ be so that (ξ_1, ξ_2) and (ξ_1, ξ_3) correspond to $\gamma(y_1)$ and $\gamma(y_2)$ via the identification of $(2^N)^{S_1} \times (2^N)^{S_2}$ with $(2^N)^N$. Then the conditions in part (2) of Lemma 3.1 are satisfied for $(\zeta, \xi_1, \xi_2, \xi_3)$, where $k = 3$, $S = X = N$, $Y_1 = S_1$, $Y_2 = Y_3 = S_2$, for either

- $\zeta = \gamma(x)$, assuming x is separated,
- $\zeta = \gamma_0(x)$, assuming x is injective.

Proof. The new aspect here is in part (2)(a) of Lemma 3.1. Fix a finite partial function $\tau: N \rightarrow \{0, 1\}$. For each $i = 1, 2, 3$, we need to find infinitely many $(s, d) \in Y_i$ so that $\xi_i(s, d)(-)$ extends τ . This is analogous to the arguments about $\gamma(x)$ in the proof of Lemma 4.4, just that now we must find $(s, d) \in S_1$, when $i = 1$, and $(s, d) \in S_2$, when $i = 2, 3$.

By assumption, given $t_1 \neq t_2 \in \mathbb{N}^{<\mathbb{N}}$, there is $i \in y_1 \cap y_2$ so that $y_1 \circ t_1(i) = y_2 \circ t_1(i) \neq y_2 \circ t_2(i) = y_1 \circ t_2(i)$. We can therefore find $s^* \in \mathbb{N}^{<\mathbb{N}}$ whose image is in $y_1 \cap y_2$, so that for any s which contains s^* and any distinct t_1, t_2 in the domain of τ , $y_1[s, t_1] \neq y_1[s, t_2]$ and $y_2[s, t_1] \neq y_2[s, t_2]$. (Using the notation from the proof of Lemma 4.4.)

Now for any $d \in \mathbb{N}$, $(s^*, d) \in S_1$. As before, for a generic α , there are infinitely many d for which $\xi_1(s^*, d)(-) = \gamma(y_1)(s^*, d)(-)$ extends τ . Next, fix some s extending s^* so that the image of s is not contained in $y_1 \cap y_2$. Then $(s, d) \in S_2$ for any $d \in \mathbb{N}$. Again we may find infinitely many d for which $\xi_2(s, d)(-) = \gamma(y_1)(s, d)(-)$ extends τ , and infinitely many d for which $\xi_3(s, d)(-) = \gamma(y_2)(s, d)(-)$ extends τ . \square

We remark that for $k = 0$ the following variation of Lemma 4.7 is used.

Lemma 7.15. Suppose $x, y, z \in (2^{\mathbb{N}})^{\mathbb{N}}$ are injective enumerations of pairwise disjoint subsets of $2^{\mathbb{N}}$. Then for any comeager $C \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^3$, for a generic β_0 , the triplet $(\gamma_0(x), \gamma_0(y), \gamma_0(z)) \in (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$ satisfies the assumption in Lemma 3.2, with $X = N$, $Y_1 = Y_2 = Y_3 = N$, and $M_1 = M_2 = M_3 = \mathbb{N}$.

7.2. Proof of Lemma 7.4. First we recall some background.

7.2.1. E -Pins. Let E be an analytic equivalence relation on a Polish space X . Assume that \mathbb{P} is a forcing poset and τ is a \mathbb{P} -name which is forced to be a member of the Polish space X , as interpreted in the generic extension. The pair (\mathbb{P}, τ) is an **E-pin** if

$$\mathbb{P} \times \mathbb{P} \Vdash \tau_l E \tau_r,$$

where τ_l, τ_r are the interpretation of τ according to the left and right generics respectively.

Lemma 7.16 (see [LZ20, Proposition 2.1.2]). For E, \mathbb{P}, τ as above, (\mathbb{P}, τ) is an E -pin if and only if in any extension of V , given two filters G_1, G_2 which are separately \mathbb{P} -generic over V , $\tau[G_1] \cap \tau[G_2]$.

The reader is referred to [LZ20, Chapter 2] for more on pins. Below we will use specifically-designed pins in certain symmetric ZF models, following [Sha21].

7.2.2. Symmetric models. Let \mathbb{P}_n be Cohen forcing for producing a generic member of $((2^{\mathbb{N}})^{\mathbb{N}})^n$. This can be defined as the poset of all Borel sets up to inclusion mod meager (see [Zap08]). An important fact we will use is that given a sufficiently large countable model M , the set of $x \in ((2^{\mathbb{N}})^{\mathbb{N}})^n$ which are \mathbb{P}_n -generic over M is comeager. We will consider the equivalent combinatorial presentation of \mathbb{P}_n as the poset of all finite approximations, ordered by extensions.

Let $G \subseteq \mathbb{P}_n$ be generic over V and $x \in ((2^{\mathbb{N}})^{\mathbb{N}})^n$ in $V[G]$ be the corresponding generic member. Note that $x \in X_n$, since $X_n \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^n$ is comeager. Let

$$A_k = A_k^x,$$

for $k \leq n$, as in Section 1.1. Consider the models $V(A_n)$, the minimal extension of V which contains A_n and satisfies ZF . Such models were studied by Monro [Mon73]. Their relationship to the Friedman-Stanley jumps was introduced in [Sha21].

Remark 7.17. For $k < n$, the poset \mathbb{P}_n can be naturally presented as a product $\mathbb{P}_k \times \mathbb{P}_n^k$, where \mathbb{P}_n^k adds a member of $((2^{\mathbb{N}})^{\mathbb{N}})^{n \setminus k}$ by finite approximations.

Fact 7.18. There is a poset \mathbb{Q}_n in $V(A_n)$, definable from A_n over V , and a \mathbb{Q}_n -name σ_n for a member of X_n , so that it is forced that $A_n^{\sigma_n} = A_n$.

Proof. Take \mathbb{Q}_n to be the poset to add, by finite approximations, a countable enumeration of the hereditary closure of A_n . The sequence $(A_k : k < n)$ may be viewed as a member of the space $\prod_{k < n} (2^{A_k})^{A_{k+1}} = (2^{\mathbb{N}})^{A_1} \times (2^{A_1})^{A_2} \times (2^{A_2})^{A_3} \times \dots$. After forcing with \mathbb{Q}_n , given enumerations of the sets A_1, A_2, \dots , we may naturally translate this to a member of $((2^{\mathbb{N}})^{\mathbb{N}})^n$, as in Section 5.1, and this will be our σ_n . \square

Remark 7.19. (\mathbb{Q}_n, σ_n) is an F_n -pin in the model $V(A_n)$. Given an equivalence relation E which extends F_n , then (\mathbb{Q}_n, σ_n) is an E -pin as well.

Definition 7.20. Let \mathbb{R}_n^k to be the product $\mathbb{Q}_k \times \mathbb{P}_n^k$, and let ρ_n^k be an \mathbb{R}_n^k -name for the member of $((2^{\mathbb{N}})^{\mathbb{N}})^n$ whose restriction to $((2^{\mathbb{N}})^{\mathbb{N}})^k$ is σ_k , and its restriction to $((2^{\mathbb{N}})^{\mathbb{N}})^{n \setminus k}$ is added by \mathbb{P}_n^k .

The useful property of \mathbb{R}_n^k is that it allows us to add the set A_n over the model $V(A_k)$ in a sufficiently homogeneous way.

Fact 7.21. There is an \mathbb{R}_n^k -generic R over $V(A_k)$ so that $A_n^{\rho_n^k[R]} = A_n$.

Fact 7.22. For any two conditions $p, q \in \mathbb{R}_n^k$ there is an automorphism of \mathbb{R}_n^k (in $V(A_k)$) sending p to q and fixing the name for $A_n^{\rho_n^k}$.

We will use the following property of the models $V(A_n)$ (see [Sha21, Lemma 4.5]).

Lemma 7.23. If $B \in V(A_n)$ is definable from A_n over V , and $B \subseteq V(A_k)$ for $k < n$, then $B \in V(A_k)$ and is definable from A_k over V .

For example, if r is a real in $V(A_1)$ which is definable from A_1 , then $r \in V$.

Proof. Let \dot{B} be the \mathbb{R}_n^k -name for the set which is defined from $A_n^{\rho_n^k}$ according to the definition of B from A_n . It follows from Remark 7.22 that for $b \in V(A_k)$, if there is some condition in \mathbb{R}_n^k forcing that $\check{b} \in \dot{B}$, then every condition in \mathbb{R}_n^k forces that $\check{b} \in \dot{B}$. We may now define B in $V(A_k)$ as the set of all $b \in V(A_k)$ for which it is forced that $\check{b} \in \dot{B}$. \square

7.2.3. *From pins to sets.* Let E be an equivalence relation which is classifiable by countable structures. Using the Scott analysis, we have a complete classification of E , $x \mapsto B_x$, assigning a hereditarily countable set to each x in the domain of E . This map is absolute, so that B_x is always the same set, no matter in which model (containing x) we perform the calculation.

Lemma 7.24. Let E and $x \mapsto B_x$ be as above and assume that (\mathbb{Q}_n, σ_n) is an E -pin in $V(A_n)$. Then there is a set $B \in V(A_n)$, definable from A_n , so that $\mathbb{Q}_n \Vdash \check{B} = B_{\sigma_n}$.

Proof. For any two \mathbb{Q}_n -generics G_1, G_2 over $V(A_n)$, $B_{\sigma_n[G_1]} = B_{\sigma_n[G_2]}$. So the set $B = B_{\sigma_n[G_1]}$ is in $V(A_n)$, definable as the unique set which is forced to be equal to B_{σ_n} . Note that B , just like A_n , is likely not hereditarily countable in $V(A_n)$. \square

Lemma 7.25. Let $B_n \in V(A_n)$ be a set which is definable from A_n over V . Let M be a sufficiently large countable substructure, $C_n \subseteq ((2^{\mathbb{N}})^{\mathbb{N}})^n$ the set of all \mathbb{P}_n -generics over M . Note that C_n is comeager and $C_n \subseteq X_n$. Define E_n on C_n by

$$x E_n y \iff B_n^x = B_n^y,$$

where B_n^y is the set defined in $M(A_n^y)$ from A_n^y according to the definition of B_n from A_n . Then E_n extends F_n on C_n . Moreover, for $k < n$, E_n extends F_k on a comeager set if and only if $B_n \in V(A_k)$ is definable from A_k over V .

Proof. If $x F_n y$ then $A_n^y = A_n^x$ and so $B_n^x = B_n^y$. Therefore E_n extends F_n on C_n .

Assume that $B_n \in V(A_k)$ and is definable from A_k over V . For $x, y \in C_n$, if $x \upharpoonright k = y \upharpoonright k$ then $M(A_k^x) = M(A_k^y)$ and therefore $B_n^x = B_n^y$, as both are definable using the same definition from A_k^x in the model $M(A_k^x)$, and so $x E_n y$ by definition. We conclude that E_n extends F_k on C_n .

Next, assume that E_n extends F_k , $k < n$ on a comeager set. Then for any \mathbb{P}_n -generics x, y over V , if $A_k^x = A_k^y$ then $B_n^x = B_n^y$, as calculated in $V(A_n^x), V(A_n^y)$ respectively. It follows that for any two \mathbb{R}_n^k -generics R_1, R_2 over $V(A_k)$, $B_n^{\rho_n^k[R_1]} = B_n^{\rho_n^k[R_2]}$, so B_n can be defined in $V(A_k)$ as the unique set which is forced to be equal to $B_n^{\rho_n^k}$. \square

7.2.4. *Concluding Lemma 7.4.* It was convenient to use the equivalence relations F_k^* before. For now let us return to the usual presentation and prove the equivalent:

Lemma 7.26. Let E be an equivalence relation, classifiable by countable structures, which extends F_n but does not extend F_k on any comeager set, for $k < n$. Then there are equivalence relations E_k for $k < n$, defined on comeager subsets of X_k , so that

- (1) E_k extends F_k on a comeager set;
- (2) E_{k+1} does not extend F_k on any comeager set;
- (3) $E \subseteq E_k$, on a comeager set, for each $k < n$. That is, on a comeager set, if $x E y$ then $x \upharpoonright k E_k y \upharpoonright k$.

Fix E as in the statement of the lemma. Apply Lemma 7.24 to the E -pin (\mathbb{Q}_n, σ_n) in $V(A_n)$, and get a set $B \in V(A_n)$ as in Lemma 7.24. B is definable from (\mathbb{Q}_n, σ_n) and therefore definable from A_n . It follows from Lemma 7.25 that $B \notin V(A_k)$ for $k < n$.

Fix ordinals η, β so that $B \in \mathcal{P}^\beta(\eta)$. For $1 \leq k < n$ define the ordinal α_k to be the least so that $\text{t.c.}(B) \cap \mathcal{P}^{\alpha_k}(\eta) \notin V(A_{k-1})$, and let $B_k = \text{t.c.}(B) \cap \mathcal{P}^{\alpha_k}(\eta)$. Since B_k is a subset of $V(A_k)$ which is definable from A_n , it follows from Lemma 7.23 that $B_k \in V(A_k)$ is definable from A_k over V . Let $B_n = B$.

We now define equivalence relations E_k from B_k , as in Lemma 7.25, so that E_k extends F_k on a comeager subset of X_k , and does not extend F_{k-1} on a comeager set. Note also that for $k < n$, since B_k is defined as the intersection of B_{k+1} with $\mathcal{P}^{\alpha_k}(\eta)$, then E_k extends E_{k+1} on a comeager set. Moreover, E_n and E agree on a comeager set. This concludes the proof of the lemma.

REFERENCES

- [CC22] John D. Clemens and Samuel Coskey, *New jump operators on equivalence relations*, Journal of Mathematical Logic **22** (2022), no. 3.
- [Cle22] John D. Clemens, *Relative primeness and borel partition properties for equivalence relations*, Trans. Amer. Math. Soc. **375** (2022), no. 1, 111–149.
- [CMRS23] F. Calderoni, D. Marker, L. Motto Ros, and A. Shani, *An anti-classification result for groups acting freely on the line*, Advances in Mathematics **418** (2023).
- [For00] Matthew Foreman, *A descriptive view of ergodic theory*, Descriptive set theory and dynamical systems (Marseille-Luminy, 1996), 2000, pp. 87–171. MR1774425
- [Fri00] Harvey M. Friedman, *Borel and Baire reducibility*, Fund. Math. **164** (2000), no. 1, 61–69. MR1784655
- [Gao09] S. Gao, *Invariant descriptive set theory*, CRC Press, 2009.
- [Hjo00] Greg Hjorth, *Classification and orbit equivalence relations*, Mathematical Surveys and Monographs, vol. 75, American Mathematical Society, Providence, RI, 2000. MR1725642
- [HK01] Greg Hjorth and Alexander S. Kechris, *Recent developments in the theory of Borel reducibility*, Fund. Math. **170** (2001), no. 1-2, 21–52. MR1881047
- [HKL98] Greg Hjorth, Alexander S. Kechris, and Alain Louveau, *Borel equivalence relations induced by actions of the symmetric group*, Ann. Pure Appl. Logic **92** (1998), no. 1, 63–112. MR1624736

- [Kan08] Vladimir Kanovei, *Borel equivalence relations*, University Lecture Series, vol. 44, American Mathematical Society, Providence, RI, 2008. Structure and classification. MR2441635
- [Kay17] Burak Kaya, *The complexity of topological conjugacy of pointed cantor minimal systems*, Archive for Mathematical Logic **56** (2017), no. 3-4, 215–235.
- [Kec95] Alexander S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995. MR1321597
- [KSZ13] Vladimir Kanovei, Marcin Sabok, and Jindřich Zapletal, *Canonical ramsey theory on polish spaces*, Cambridge Tracts in Mathematics, Cambridge University Press, 2013.
- [Lou94] Alain Louveau, *On the reducibility order between Borel equivalence relations*, Logic, methodology and philosophy of science, IX (Uppsala, 1991), 1994, pp. 151–155. MR1327979
- [LZ20] Paul B. Larson and Jindřich Zapletal, *Geometric set theory*, Mathematical Surveys and Monographs, vol. 248, American Mathematical Society, Providence, RI, 2020. MR4249448
- [Mar07] David Marker, *The Borel complexity of isomorphism for theories with many types*, Notre Dame J. Formal Logic **48** (2007), no. 1, 93–97. MR2289899
- [Mon73] G. P. Monro, *Models of ZF with the same sets of sets of ordinals*, Fund. Math. **80** (1973), no. 2, 105–110. MR0347602
- [Sha21] Assaf Shani, *Borel reducibility and symmetric models*, Trans. Amer. Math. Soc. **374** (2021), no. 1, 453–485. MR4188189
- [URL17] Douglas Ulrich, Richard Rast, and Michael C. Laskowski, *Borel complexity and potential canonical Scott sentences*, Fund. Math. **239** (2017), no. 2, 101–147. MR3681584
- [Zap08] Jindřich Zapletal, *Forcing idealized*, Cambridge Tracts in Mathematics, Cambridge University Press, 2008.

DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY UNIVERSITY,
MONTREAL, QC H3G 1M8, CANADA

Email address: assaf.shani@concordia.ca

URL: <https://sites.google.com/view/assaf-shani/>