

A COUNTEREXAMPLE REGARDING AN EQUIVALENCE RELATION ON A PRODUCT SPACE

ASSAF SHANI

ABSTRACT. We present a counterexample to Conjecture 14.1.6 from [Kan08], regarding Borel equivalence relations on product spaces.

This paper concerns the study of equivalence relations on Polish spaces up to Borel reducibility. Given equivalence relations E and F on Polish spaces X and Y respectively, a map $f: X \rightarrow Y$ is said to be a **reduction** of E to F if for any $x_1, x_2 \in X$,

$$x_1 E x_2 \iff f(x_1) F f(x_2).$$

We say that E is **Borel reducible** to F , denoted $E \leq_B F$ if there is a Borel measurable function which is a reduction of E to F . In this case, we think of E as no more complicated than F . Borel reducibility is the most central concept in the study of equivalence relations on Polish spaces. Say that E and F are **Borel bireducible**, denoted $E \sim_B F$, if $E \leq_B F$ and $F \leq_B E$. We write $E <_B F$ (a strict reduction) if a Borel reduction exists in only that direction. An equivalence relation E on a Polish space X is **Borel** if E is a Borel subset of $X \times X$, with the product topology.

A central motivation for the field is to study the complexity of various classification problems in mathematics. Generally speaking, separable mathematical objects can be coded as members of some Polish space. Natural notions of equivalence, such as isomorphism between countable graphs, isometry between separable metric spaces, or homeomorphism between compact metric spaces, can then be seen as equivalence relations on Polish spaces. These are generally analytic, and sometimes Borel. The focus on Borel measurable reductions is in attempt to consider only maps coming from “natural mathematical practice”. The particular choice of “Borel measurability” is strongly supported by theoretical and experimental evidence, as well as the success of the field since its inception with the papers [HKL90, FS89]. For more background the reader is referred to the expository books [Gao09, Kan08], and the surveys [HK01, Kec99, Kec02, KTD12, For18, MR21].

Given equivalence relations F_1, F_2 on spaces Y_1, Y_2 respectively, define $F_1 \times F_2$ on $Y_1 \times Y_2$ as the pointwise relation: $(y_1, y_2), (y'_1, y'_2) \in Y_1 \times Y_2$ are $F_1 \times F_2$ -related if $y_1 F_1 y'_1$ and $y_2 F_2 y'_2$.

In this note we provide a counterexample to the following conjecture, attributed to Zapletal in [Kan08].

Date: January 30, 2025.

Conjecture 1 (Conjecture 14.1.6 in [Kan08]). Let X, Y be Polish spaces, $P \subseteq X \times Y$ a Borel set, F a Borel equivalence relation on X and E a Borel equivalence relation on P so that for any $(x, y), (x', y') \in P$,

$$(x, y) E (x', y') \implies x F x'.$$

Assume that G is a Borel equivalence relation so that for any $x_0 \in X$,

$$E \upharpoonright \{(x, y) \in P : x F x_0\} \leq_B G.$$

Then $E \leq_B F \times G$.

Roughly speaking, the conjecture says that if E extends F on the first coordinate, and on every fiber E is no more complicated than G , then E is no more complicated than $F \times G$.

Two particularly important families of Borel equivalence relations are the smooth equivalence relations and the countable equivalence relations. (See Chapters 5.4 & 7 in [Gao09].)

An equivalence relation G is said to be *smooth* if G is Borel reducible to $=_{\mathbb{R}}$, where $=_{\mathbb{R}}$ is the equality relation on the Polish space \mathbb{R} . A Borel equivalence relation F on a Polish space Y is *countable* if each F -class is countable.

The motivation for this conjecture is as follows. The conjecture is true in the case where F is a countable Borel equivalence relation and G is smooth (see [Kan08, Theorem 14.1.1]). This fact is essential in the proof of a dichotomy, due to Hjorth and KeCHRIS [HK01], for the equivalence relation E_3 (see [Kan08, Chapter 14]).

Our counterexample will use the Friedman-Stanley jumps, defined as follows. Given a space X , let $X^{\mathbb{N}}$ be the space of all sequence $x = (x(n) : n \in \mathbb{N})$, where $x(n) \in X$, equipped with the product topology. If X is Polish, then $X^{\mathbb{N}}$ is Polish.

Definition 2 (Friedman-Stanley [FS89]). Let E be an equivalence relation on a Polish space X . Define E^+ on the space $X^{\mathbb{N}}$ by

$$x E^+ y \iff \forall n \exists m (x(n) E y(m)) \ \& \ \forall n \exists m (y(n) E x(m)),$$

that is, $\{[x(n)]_E : n \in \mathbb{N}\} = \{[y(n)]_E : n \in \mathbb{N}\}$.

Note that if E is Borel, then E^+ is Borel. Friedman and Stanley proved the following (see [Gao09, 8.3.6]).

Fact 3 (Friedman-Stanley [FS89]). For a Borel equivalence relation E , $E <_B E^+$.

Define F on $X = \mathbb{R}^{\mathbb{N}}$ by $F = (=_{\mathbb{R}})^+$. That is, $x F y$ if x and y enumerate the same sets of reals.

Next we define the equivalence relation E . Let $2^{\mathbb{N}}$ be the space of all binary sequences, given the product topology, where the set $2 = \{0, 1\}$ is given the discrete topology. Consider the Polish space $\mathbb{R}^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$ with the product topology. Given $(x, y) \in \mathbb{R}^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$, define $A_n^{(x,y)} = \{x(n) : y(n) = 1\}$ for $n \in \mathbb{N}$. That is, we view each $y(n)$ is a binary sequence “carving out” a subset of the set enumerated x .

$$\begin{array}{cccccccc}
 \vdots & \vdots & \vdots & \vdots & & \vdots & \vdots & \vdots \\
 * & 1 & 0 & 1 & \dots & * & - & * & \dots \\
 * & 1 & 1 & 0 & \dots & * & * & - & \dots \\
 * & 0 & 1 & 1 & \dots & - & * & * & \dots \\
 * & 0 & 1 & 0 & \dots & - & * & - & \dots \\
 x & y(0) & y(1) & y(2) & & A_0^{(x,y)} & A_1^{(x,y)} & A_2^{(x,y)} &
 \end{array} \mapsto$$

Define $P \subseteq \mathbb{R}^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$ as the set of all (x, y) such that:

- (1) $\forall m \exists k (y(k)(m) = 1)$;
- (2) $\forall k \exists m (y(k)(m) = 1)$;
- (3) $\forall k, l_1, l_2 (x(l_1) = x(l_2) \rightarrow y(k)(l_1) = y(k)(l_2))$.

That is: (1) each $x(m)$ appears in $A_k^{(x,y)}$ for some k , (2) each $A_k^{(x,y)}$ is not empty, and (3) if x enumerates the same item twice, on indices l_1 and l_2 , then each binary sequence $y(k)$ agrees on l_1 and l_2 . Note that P is a Borel set. In fact it is a dense G_δ subset of $\mathbb{R}^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$.

Finally, we define an equivalence relation E on P by

$$(x, y) E (x', y') \iff \{A_n^{(x,y)} : n \in \mathbb{N}\} = \{A_n^{(x',y')} : n \in \mathbb{N}\},$$

that is, (x, y) and (x', y') code the same sets of subsets of \mathbb{R} . Note that E is Borel.

Let $=_{2^{\mathbb{N}}}$ be the equality relation on $2^{\mathbb{N}}$, and let $G = (=_{2^{\mathbb{N}}})^+$. Note that G and F are Borel bireducible (as there is a Borel bijection between \mathbb{R} and $2^{\mathbb{N}}$, see [Gao09, 1.4.3]).

Claim 4. For any $x_0 \in \mathbb{R}^{\mathbb{N}}$,

$$E \upharpoonright \{(x, y) \in P : x F x_0\} \leq_B G.$$

Proof. Fix $x_0 \in \mathbb{R}^{\mathbb{N}}$. Define $f: \{(x, y) \in P : x F x_0\} \rightarrow (2^{\mathbb{N}})^{\mathbb{N}}$ by

$$\begin{aligned}
 f(x, y)(n)(k) = i &\iff (\forall k' \in \mathbb{N})(x(k') = x_0(k) \implies y(n)(k') = i) \\
 &\iff (\exists k' \in \mathbb{N})(x(k') = x_0(k) \wedge y(n)(k') = i).
 \end{aligned}$$

The equivalence between the definitions above follows from part (3) in the definition of P . Then f is a reduction of $E \upharpoonright \{(x, y) \in P : x F x_0\}$ to F . \square

What f does is the following. The F -class of x_0 are all sequences which enumerate the same set as x_0 . Using a fixed enumeration of this set, via x_0 , we code subsets of it, $A_n^{(x,y)}$, as binary sequences by identifying k with $x_0(k)$. In particular, $f(x_0, y) = y$ for any y .

Theorem 5. The equivalence relations E, F, G provide a counterexample to Conjecture 1.

Remark 6. The equivalence relation E is Borel bireducible with the ‘second Friedman-Stanley jump’, $((=_{\mathbb{R}})^+)^+$. The equivalence relation E was studied in [Sha], particularly with respect to Baire category techniques. Note that $((=_{\mathbb{R}})^+)^+$ is defined on

the space $(\mathbb{R}^{\mathbb{N}})^{\mathbb{N}}$ and therefore does not seem to provide a counterexample to Conjecture 1. The different presentation of $((=\mathbb{R})^+)^+$, as E on $\mathbb{R}^{\mathbb{N}} \times (2^{\mathbb{N}})^{\mathbb{N}}$, is therefore crucial here.

We include a proof that $((=\mathbb{R})^+)^+ \leq_B E$, which we use below. The idea is simple. Given $z \in (\mathbb{R}^{\mathbb{N}})^{\mathbb{N}}$, each $z(n) \in \mathbb{R}^{\mathbb{N}}$ codes a subset of \mathbb{R} . We will send z to a pair $(x, y) \in P$ so that x enumerates the set of all reals appearing in the sets $z(n)$, and $y(n)$ carves out the subset $z(n)$ of x .

Fix a bijection $e: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$. Define $x(e(i, j)) = z(i)(j)$. Define $y(n)(e(i, j)) = 1$ if there is some k so that $z(n)(k) = z(i, j)$, and $y(n)(e(i, j)) = 0$ otherwise. The map $z \mapsto (x, y)$ is a Borel reduction of $((=\mathbb{R})^+)^+$ to E .

We will also use the well known fact that $F \times F \sim_B F$. For the non-trivial direction, consider the map sending $(x, y) \in \mathbb{R}^{\mathbb{N}} \times \mathbb{R}^{\mathbb{N}}$ to $z \in \mathbb{R}^{\mathbb{N}}$ so that $z(2n) = x(n)$ and $z(2n + 1) = y(n)$. Then $(x, y) \mapsto z$ is a Borel reduction of $F \times F$ to F .

Now Conjecture 1 would imply that $E \leq_B F \times G$, and so

$$((=\mathbb{R})^+)^+ \leq_B E \leq_B F \times G \leq_B F \times F \leq_B F \leq_B (=\mathbb{R})^+,$$

a contradiction, as $((=\mathbb{R})^+)^+ \not\leq_B (=\mathbb{R})^+$ by Fact 3.

REFERENCES

- [For18] Matthew D. Foreman, *What is ... a borel reduction?*, Notices of the American Mathematical Society (2018).
- [FS89] Harvey Friedman and Lee Stanley, *A borel reducibility theory for classes of countable structures*, J. Symbolic Logic **54** (1989), no. 3, 894–914.
- [Gao09] S. Gao, *Invariant descriptive set theory*, CRC Press, 2009.
- [HK01] Greg Hjorth and Alexander S. Kechris, *Recent developments in the theory of Borel reducibility*, Fund. Math. **170** (2001), no. 1-2, 21–52. MR1881047
- [HKL90] L. A. Harrington, A. S. Kechris, and A. Louveau, *A Glimm-Effros dichotomy for Borel equivalence relations*, J. Amer. Math. Soc. **3** (1990), no. 4, 903–928. MR1057041
- [Kan08] Vladimir Kanovei, *Borel equivalence relations*, University Lecture Series, vol. 44, American Mathematical Society, Providence, RI, 2008. Structure and classification. MR2441635
- [Kec02] Alexander S. Kechris, *Actions of Polish groups and classification problems*, Analysis and logic (Mons, 1997), 2002, pp. 115–187. MR1967835
- [Kec99] Alexander S. Kechris, *New directions in descriptive set theory*, The Bulletin of Symbolic Logic **5** (1999), no. 2, 161–174.
- [KTD12] Alexander S. Kechris and Robin D. Tucker-Drob, *The complexity of classification problems in ergodic theory* (2012), 265–300.
- [MR21] Luca Motto Ros, *Classification problems from the descriptive set theoretical perspective* (2021).
- [Sha] Assaf Shani, *The finite friedman-stanley jumps: generic dichotomies for borel homomorphisms*. *arxiv:2405.18360*.

DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY UNIVERSITY,
MONTREAL, QC H3G 1M8, CANADA

Email address: assaf.shani@concordia.ca

URL: <https://sites.google.com/view/assaf-shani/>