

BOREL REDUCIBILITY AND VIRTUAL CLASSES
(DRAFT - UNDER PREPARATION)

ASSAF SHANI

In set theory, a *virtual* object is one that exists in a generic extension. This note aims to introduce the study of virtual equivalence classes, and its applicability, particularly towards questions about Borel reducibility and Vaught’s conjecture. The emphasis will be on the unifying aspect of this approach. For example, we will present proofs of the topological Vaught’s conjecture for CLI groups (due to Becker) and Harrington’s theorem on models of size \aleph_1 for counterexamples to the $\mathcal{L}_{\omega_1, \omega}$ -Vaught conjecture, as well as a result on abelian torsion groups of size \aleph_1 due to Fuchs and Kulikov, and the well known existence of a sequence of Turing degrees satisfying $x'_{n+1} \leq_T x_n$. With respect to the last two results especially, this note is very much in the spirit of [Mil17], proving things “the hard way”. Nevertheless, the given treatment of the topological Vaught conjecture is quite simple, and for several application to equivalence relations the only known proofs use virtual classes.

This note will be most attractive to a reader with some background and an interest in set theory. Such reader will find the details reasonable, and the general approach quite pleasing. I hope that any reader interested in Borel reducibility or Vaught’s conjecture will also benefit from this note, in particular to get an intuition for what kind of questions may be susceptible to the techniques presented here.

The reader is referred to [Mil17] for background on the interaction between descriptive set theory and forcing. Particularly, we assume familiarity with Cohen forcing (in both the combinatorial and topological presentations), the Levy collapse, as well as Mostowski and Shoenfield absoluteness.

The paper [Hjo98] is also a survey of applications of ‘virtual Borel sets’ to descriptive set theory. In a sense the present note can be seen as a continuation of that line of work. The focus is a little different and most applications presented here are not present in [Hjo98]. When there is overlap, the current presentation is cleaner thanks to recent developments such as [Kan08b, LZ20]. Chapter 2 of [LZ20, Chapter 2] is a systematic study of virtual equivalence classes, the most comprehensive to date. Virtual classes were also studied recently from various perspectives [KMS16, KS16, URL17, Sha21]. This note, while far from comprehensive, attempts to present a more unified storyline. Many of the proofs presented here are different than in other sources.

For an introduction to Borel reducibility the reader is referred to the books [Gao09, Kan08b] and the surveys [MR21, For18, KTD12, HK01, Kec99]. The book [Kan08b] particularly emphasizes forcing techniques. With regards to applications to Borel reducibility, we focus here on techniques rather than state-of-the-art results. In particular we use virtual classes to study well known benchmark equivalence relations such as $E_0, E_0^{\mathbb{N}}, =^+, E_{\omega_1}$. Some recent results are presented, in particular, a result from [CT19] on the isomorphism and bi-embeddability relations

for abelian torsion groups, and a result from [CMRS23] on the isomorphism relation for Archimedean ordered groups.

1. VIRTUAL CLASSES AND VAUGHT’S CONJECTURE

Let X be a Polish space and E an analytic equivalence relation on X . An important fact which we use below is that the space X and the equivalence relation E can be reasonably interpreted in any generic extension. We will often rely on Mowstowski’s Absoluteness Theorem (see [Jec03, Theorem 25.4]) to conclude that the statement $x E y$ agrees between various models which contain both x and y . The discussion below naturally extends beyond analytic equivalence relations, as long as E can be interpreted in a sufficiently absolute manner.

Definition 1.1. Let \mathbb{P} be a forcing poset and τ a \mathbb{P} -name so that $\mathbb{P} \Vdash \tau \in X$. We may think of τ as a *virtual member* of X , or of $[\tau]_E$ as a *virtual equivalence class*.¹ Assume further that in any generic extension, given two filters $G_1, G_2 \subseteq \mathbb{P}$ which are generic over the ground model, we have that

$$\tau[G_1] E \tau[G_2].$$

In this case we say that the pair (\mathbb{P}, τ) is a *stable virtual E -class* [Kan08b, 17.1.2], or an *E -pin* [LZ20, 2.1.1].

In this case $[\tau]_E$ is an equivalence class in a generic extension, which is “stable” or “pinned” in the sense that its interpretation does not depend on the generic filter. We will often omit the adjective ‘stable’ and call (\mathbb{P}, τ) a virtual E -class.

Given $x \in X$ and a poset \mathbb{P} , let \check{x} be the canonical \mathbb{P} -name for x . Then (\mathbb{P}, \check{x}) is a stable virtual E -class.

Definition 1.2. Say that a stable virtual E -class (\mathbb{P}, τ) is *pinned* [Kan08b, 17.1.2], or *trivial* [LZ20, 2.3.1], if there is some $x \in X$ so that

$$\mathbb{P} \Vdash \check{x} E \tau.$$

In the descriptive set theoretic context, the continuum hypothesis is often identified with a perfect set property: a set is either countable, or admits a definable injective image of $2^{\mathbb{N}}$. In this sense, a definable set of size \aleph_1 can be seen as a violation of the continuum hypothesis.

By classical descriptive set theoretic wisdom, all analytic sets satisfy the perfect set property (see [Kec95, 29.1]). The modern wisdom is that any reasonably definable subset of a Polish space satisfies the perfect set property (see [Kan03], page 145 as well as Theorem 27.9 and the following discussion).

Next, we consider sets defined as quotients of Polish spaces by definable equivalence relations. We have *Silver’s dichotomy* (see [Gao09, 5.3.5]): for any co-analytic equivalence relation E (in particular, any Borel equivalence relation), either there are countably many E -classes or there is a perfect set of E -inequivalent elements. (That is, the quotient X/E satisfies the continuum hypothesis.)

Example 1.3. Let X be the Polish space of all linear orderings of \mathbb{N} . Define E_{ω_1} on X by $x E_{\omega_1} y$ if x, y are isomorphic, or if both are ill-founded. E_{ω_1} is an analytic equivalence relation with \aleph_1 many equivalence classes.

Fix an uncountable ordinal α , let $\mathbb{P}_\alpha = \text{Col}(\omega, |\alpha|)$, and let τ_α be a \mathbb{P}_α -name for a member of X of order-type α . Then $(\mathbb{P}_\alpha, \tau_\alpha)$ is a non-trivial virtual E_{ω_1} -class.

¹A closely related terminology is that of *virtual Borel sets* [Ste84, Hjo98].

For a countable first order language \mathcal{L} consider $\text{Mod}(L)$, the Polish space of all \mathcal{L} -structures with universe \mathbb{N} , and \cong_L the isomorphism relation on $\text{Mod}(L)$ (see [Gao09, 3.6]). Given a theory T we let $\text{Mod}(T)$ be the subspace of all \mathcal{L} -structures which are models of T , and \cong_T the isomorphism relation restricted to $\text{Mod}(T)$.

The equivalence relation \cong_T is generally analytic, so is not covered by Silver's dichotomy. *Vaught's Conjecture* asserts that a violation of CH cannot be defined in this way, that is, either \cong_T has countably many countable models, up to isomorphism, or there is a perfect set of pairwise non-isomorphic models of T (see [Gao09, p.261]).

We begin this note by showing how non-trivial virtual classes arise naturally from definable violations of the continuum hypothesis.

Proposition 1.4. Suppose T is a counterexample to Vaught's Conjecture. That is

- (1) $\text{Mod}(T)$ has uncountably many equivalence classes;
- (2) there is no perfect set of \cong_T -in-equivalent elements.

Then there is an unpinned stable virtual \cong_T -class.

First, we make some remarks about product forcing (see [Jec03, p. 229], [Kun80, VIII §1]), and prove the key Lemma 1.6 towards the proof of Proposition 1.4.

Given posets \mathbb{P}, \mathbb{Q} , we may force with the product poset $\mathbb{P} \times \mathbb{Q}$. In this case, a generic filter may be identified as $G \times H$, where G, H and \mathbb{P}, \mathbb{Q} -generics respectively, and we say that G and H are *mutually generic*. Given a poset \mathbb{P} and a \mathbb{P} -name τ , let τ_l, τ_r be $\mathbb{P} \times \mathbb{P}$ -names so that given a generic filter $G_l \times G_r \subseteq \mathbb{P} \times \mathbb{P}$, $\tau_l[G_l \times G_r] = \tau_l[G_l]$ and $\tau_r[G_l \times G_r] = \tau_r[G_r]$.

Lemma 1.5 ([LZ20, 2.1.2]). Let \mathbb{P} be a forcing poset and τ and \mathbb{P} -name so that $\mathbb{P} \Vdash \tau \in X$. Then (\mathbb{P}, τ) is stable if and only if

$$\mathbb{P} \times \mathbb{P} \Vdash \tau_l E \tau_r.$$

Proof. For the non-trivial direction, we assume that $\mathbb{P} \times \mathbb{P} \Vdash \tau_l E \tau_r$ and need to show that in any generic extension $V[H]$, given two filters $G_1, G_2 \subseteq \mathbb{P}$ which are generic over V (but not necessarily mutually generic), $\tau[G_1] E \tau[G_2]$. Let G be \mathbb{P} -generic over $V[H]$. In this case, $G_1 \times G$ and $G \times G_2$ are $\mathbb{P} \times \mathbb{P}$ -generic over V (see [Kun80, Theorem 1.4]). We conclude that

$$\tau[G_1] E \tau[G] E \tau[G_2].$$

We remark that the latter argument was carried in the model $V[H][G]$, and we use the absoluteness of the statement $\tau[G_1] E \tau[G_2]$ to conclude that it is true in $V[H]$ as well. \square

Lemma 1.6. Let \mathbb{P} be a forcing poset and τ and \mathbb{P} -name so that $\mathbb{P} \Vdash \tau \in X$. Given $p \in \mathbb{P}$, let $\mathbb{P} \upharpoonright p$ be the poset of all conditions stronger than p . Assume that there is no $p \in \mathbb{P}$ for which $(\mathbb{P} \upharpoonright p, \tau)$ is a stable virtual E -class. Then there is a perfect set $Y \subseteq X$ of E -inequivalent elements.

Proof. By assumption, for any $p \in \mathbb{P}$, the condition $(p, p) \in \mathbb{P} \times \mathbb{P}$ does not force that $\tau_l E \tau_r$. It follows that for any $p \in \mathbb{P}$ we may find q_1, q_2 extending p so that $(q_1, q_2) \Vdash \tau_l \not E \tau_r$. We can iterate this fact to build a 'binary tree of conditions'.

Let $(D_n : n \in \omega)$, $(C_n : n \in \omega)$ be sequences of dense open subsets of \mathbb{P} and $\mathbb{P} \times \mathbb{P}$ respectively. We may recursively define conditions $p_s \in \mathbb{P}$ for each s in the full binary tree $2^{<\omega}$ so that for each $s \in 2^{<\omega}$,

- $p_s \in D_{|s|}$, where $|s|$ is the length of s ,
- $p_{s \smallfrown 0}, p_{s \smallfrown 1}$ both extend p_s ;
- $(p_{s \smallfrown 0}, p_{s \smallfrown 1})$ force that $\tau_l \not\equiv \tau_r$.
- $(p_{s \smallfrown 0}, p_{s \smallfrown 1}) \in C_{|s|+1}$

A branch in the binary tree $b \in 2^\omega$ corresponds to a sequence of conditions $(p_{b \upharpoonright n} : n \in \omega)$. The idea is that this sequence of conditions decide enough information about τ to determine a unique point in $x_b \in X$. This can be achieved by a sufficiently rich family $(D_n : n \in \omega)$. Furthermore, a sufficiently rich family $(C_n : n \in \omega)$ will ensure that for distinct branches b, b' , $x_b \not\equiv x_{b'}$.

One way to do this is as follows. Fix a ‘sufficiently elementary’ countable model M and let $(D_n : n \in \omega)$, $(C_n : n \in \omega)$ be an enumeration of all dense open subsets in M of \mathbb{P} , $\mathbb{P} \times \mathbb{P}$ respectively. \square

Proof of Proposition 1.4. We use the fact that item (1) above is absolute, that is, true in any generic extension. Work in a $\mathbb{P} = \text{Col}(\omega, \mathbb{R})$ generic extension. As there are uncountably many $\text{Mod}(T)$ -equivalence classes, and only countably many from the ground model, there is some member of $\text{Mod}(T)$ which is not equivalent to any ground model member. That is, we may find a \mathbb{P} -name τ so that for any $x \in \text{Mod}(T)$ in the ground model, $\mathbb{P} \Vdash \tau \not\equiv_T \dot{x}$. By Lemma 1.6 and item (2), there is some $p \in \mathbb{P}$ so that $(\mathbb{P} \upharpoonright p, \tau)$ is a stable virtual \cong_T -class, which is not trivial by the choice of τ .

The absoluteness of (1) follows from Shoenfield’s Absoluteness Theorem (see [Jec03, Theorem 25.20]) as it can be presented as a $\mathbf{\Pi}_2^1$ statement. The most obvious attempt to phrase it does not quite work: ‘for any sequence $x \in (\text{Mod}(T))^\mathbb{N}$ there is $y \in \text{Mod}(T)$ so that for all $n \in \mathbb{N}$, y is not isomorphic to $x(n)$ ’, is a $\mathbf{\Pi}_3^1$ formula. An equivalent $\mathbf{\Pi}_2^1$ statement is ‘for any countable list of $\mathcal{L}_{\omega_1, \omega}$ sentences (ψ_1, ψ_2, \dots) there is $y \in \text{Mod}(T)$ with Scott sentence different from ψ_i for each $i \in \mathbb{N}$ ’. See also [Lar17, Remark 10.4] \square

1.0.1. *The topological Vaught’s Conjecture.* We briefly mention here that the arguments above work for the $\mathcal{L}_{\omega_1, \omega}$ -Vaught’s Conjecture as well as the more general Topological Vaught’s Conjecture.

Given an $\mathcal{L}_{\omega_1, \omega}$ sentence ϕ , let $\text{Mod}(\phi)$ the Polish space of all \mathcal{L} -structures with universe \mathbb{N} , which satisfy ϕ , and \cong_ϕ the isomorphism relation on $\text{Mod}(\phi)$. The $\mathcal{L}_{\omega_1, \omega}$ -Vaught conjecture asserts that \cong_ϕ either has countably many equivalence classes, or else there is a perfect set of $\text{Mod}(\phi)$ -inequivalent members of $\text{Mod}(\phi)$.

The isomorphism relation on models with universe \mathbb{N} may be viewed as induced by an action of S_∞ , the group of all permutations of \mathbb{N} . Equipped with the pointwise convergence topology, this is a Polish group (see [Gao09, Section 2.4]). Given a Polish group G and a continuous action $a: G \curvearrowright X$ on a Polish space X , let E_a be the induced *orbit equivalence relation* on X ,

$$x E y \iff \exists g \in G (g \cdot x = y).$$

The *topological Vaught conjecture* asserts that E_a either has countably many equivalence classes, or else there is a perfect set of E_a -inequivalent members of X . Say

that a group G satisfies *Vaught's conjecture* if this holds for any continuous action of G on a Polish space.²

Proposition 1.7. Suppose $a: G \curvearrowright X$ is a counterexample to the topological Vaught conjecture. Then there is an unpinned stable virtual E_a -class.

The proof proceeds as Proposition 1.4, after proving the absoluteness of “ E_a is a counterexample to the topological Vaught conjecture”. See [Hjo01, Claim (2) on p.133].

1.1. Pinned equivalence relations.

Definition 1.8. Say that E is *pinned* if every stable virtual E -class is pinned.

We begin with a few simple examples. We will make use of following ‘mutual genericity’ lemma.

Lemma 1.9. If G, H are mutually generic over V then $V[G] \cap V[H] = V$.

Recall that if $G \times H$ is $\mathbb{P} \times \mathbb{Q}$ -generic over V , then G is \mathbb{P} -generic over $V[H]$. The lemma therefore is a corollary of the following (applied with $N = V$ and $M = V[H]$).

Lemma 1.10. Suppose $N \subseteq M$ are transitive models of ZF, $\mathbb{P} \in N$ is a poset. If G is \mathbb{P} -generic over M , then $N[G] \cap M = N$.

Proof sketch. Suppose $\tau \in N$ is a \mathbb{P} -name and $\tau[G] = X \in M$. Using an inductive argument on the rank, it suffices to assume that $X \subseteq N$, and conclude that $X \in N$. Since G is generic over M , we have $\tau[G] = X$ in $M[G]$ as well. Working in M , we may find $p \in \mathbb{P}$ forcing that $\tau = \check{X}$. Now the set X can be defined in N as the set of all x for which $p \Vdash \check{x} \in \tau$. The point is that for any $q \in \mathbb{P}$ and $x \in N$, the statement $q \Vdash \check{x} \in \tau$ is absolute between M and N . \square

Example 1.11. The equality relation on \mathbb{R} , $=_{\mathbb{R}}$, is pinned.

Proof. Assume that (\mathbb{P}, τ) is a stable virtual $=_{\mathbb{R}}$ -class. Take G_1, G_2 mutually \mathbb{P} -generics over V , and let $x = \tau[G_1]$. Then

$$x := \tau[G_1] = \tau[G_2] \in V[G_1] \cap V[G_2] = V.$$

We see now that $\mathbb{P} \Vdash \tau E \check{x}$, and so (\mathbb{P}, τ) is trivial. \square

Example 1.12. Let Γ be a countable group, $a: \Gamma \curvearrowright X$ a Borel action, and $E = E_a$ the induced orbit equivalence relation. Then E is pinned.

Proof. Assume that (\mathbb{P}, τ) is a stable virtual E -class. Take G_1, G_2 mutually \mathbb{P} -generics over V . Then

$$A := [\tau[G_1]]_E = [\tau[G_2]]_E \in V[G_1] \cap V[G_2] = V.$$

In this calculation we used the fact that, as Γ is a countable group, the orbit $[\tau[G_1]]_E = \Gamma \cdot \tau[G_1]$ is the same whether calculated in $V[G_1]$ or $V[G_1 \times G_2]$. Fix some $x \in A$. Now $x \in V$ (since $A \in V$), and $\mathbb{P} \Vdash \tau E \check{x}$. \square

²By results of Becker and Kechris [BK96], ‘continuous action’ may be replaced by ‘Borel action’, see [Gao09, Proposition 9.5.1].

Example 1.13. Let $\Gamma_n, n \in \mathbb{N}$, be countable groups, $a_n: \Gamma_n \curvearrowright X_n$ Borel actions. Consider the pointwise product action $a: \prod_{n \in \mathbb{N}} \Gamma_n \curvearrowright \prod_{n \in \mathbb{N}} X_n$, and let $E = E_a$ be the induced orbit equivalence relation on $\prod_{n \in \mathbb{N}} X_n$. Then E is pinned.

Proof. Let (\mathbb{P}, τ) be a stable virtual E -class. Fix G_1, G_2 mutually \mathbb{P} -generics over V . Let $\tau[G_i]_n \in X_n$ be the n 'th coordinate of the sequence $\tau[G_i]$. Since $\tau[G_1] E \tau[G_2]$, we have that $\tau[G_1]_n, \tau[G_2]_n$ are in the same orbit of the action a_n . Define

$$A_n := \Gamma_n \cdot \tau[G_1]_n = \Gamma_n \cdot \tau[G_2]_n.$$

Then

$$A := (A_n : n \in \mathbb{N}) \in V[G_1] \cap V[G_2] = V.$$

Working in V , fix a sequence $x \in \prod_n A_n$. Then $\mathbb{P} \Vdash \tau E \check{x}$. \square

We have seen in Example 1.3 that E_{ω_1} is not pinned. The most central example of an unpinned *Borel* equivalence relation is the following.

Example 1.14. Consider the natural action $a: S_\infty \curvearrowright \mathbb{R}^\mathbb{N}$, defined by $g \cdot x = x \circ g^{-1}$. Let $E = E_a$ be the induced orbit equivalence relation on $\mathbb{R}^\mathbb{N}$. Then E is not pinned.

Proof. Let $\mathbb{P} = \text{Col}(\omega, \mathbb{R})$, and let τ be a \mathbb{P} -name which is forced to be a member of $\mathbb{R}^\mathbb{N}$ enumerating the ground model reals. Working in a generic extension, if G_1, G_2 are \mathbb{P} -generics over V , then $\tau[G_1], \tau[G_2]$ are two enumerations of the same set, so there is a permutation $g \in S_\infty$ so that $\tau[G_1] \circ g = \tau[G_2]$. Therefore (\mathbb{P}, τ) is a stable virtual E -class. For any $x \in \mathbb{R}^\mathbb{N}$ in the ground model, a countable set of reals, it is forced that $\{x_n : n \in \mathbb{N}\}$ is a strict subset of $\{\tau_n : n \in \mathbb{N}\}$, and so $\tau \not E \check{x}$. So (\mathbb{P}, τ) is not trivial. \square

A Polish group G is *CLI* if it *admits a complete left invariant metric*. See [Gao09, Chapter 2] for a thorough discussion on CLI Polish groups. We mention here that:

- G is CLI if and only if it admits a complete *right* invariant metric;
- any abelian Polish group is CLI;
- given countable groups $\Gamma_n, n \in \mathbb{N}$, the product group $\prod_{n \in \mathbb{N}} \Gamma_n$ is CLI.
- the group S_∞ is not CLI.

A big success towards the topological approach to Vaught's Conjecture is a theorem of Sami [Sam94] that all abelian Polish groups satisfy Vaught's Conjecture. This was strengthened by Becker, who proved that all CLI groups satisfy Vaught's conjecture (see [Gao09, Theorem 9.5.2]). Becker's result follows from the following theorem, together with Proposition 1.6.

Theorem 1.15 (Hjorth [Hjo99], see [Kan08b, 17.4]). Let G be a CLI Polish group, $a: G \curvearrowright X$ a continuous action. Then E_a is pinned.

Corollary 1.16 (Becker). Every CLI group satisfies the Vaught conjecture.

Proof of Theorem 1.15. Fix a complete right invariant metric d for G and denote the identity of G by e . Let (\mathbb{P}, τ) be a stable virtual E -class. We will construct, in V , the following objects.

- A descending sequence of closed subsets of X , $C_0 \supseteq C_1 \supseteq \dots$ with diameters converging to 0;
- a descending sequence of conditions in \mathbb{P} , $p_0 \geq p_1 \geq \dots$ so that
 - (1) p_n forces (w.r.t. \mathbb{P}) $\tau \in C_n$;

- (2) (p_n, p_n) forces (w.r.t. $\mathbb{P} \times \mathbb{P}$) that there is $g \in G$ with $d(g, e) < \frac{1}{2^n}$ so that $g \cdot \tau_l = \tau_r$.

Given these, let $x \in X$ be the unique point in the intersection $\bigcap_n C_n$.

Claim 1.17. $\mathbb{P} \Vdash \tau E \check{x}$.

Proof. Since (\mathbb{P}, τ) is a stable virtual class, it suffices to prove that $p_0 \Vdash \tau E \check{x}$. Fix a \mathbb{P} -generic H over V which contains p_0 . We need to show that $\tau[H] E x$, that is, we need to find $g \in G$ (in $V[H]$) so that $g \cdot \tau[H] = x$. By absoluteness of the statement ' $\tau[H] E x$ ', it suffices to show this in a further generic extension.

We force further³, to find a sequence H_1, H_2, \dots of \mathbb{P} -generic filters over V so that, letting $H = H_0$,

- $p_n \in H_n$ and
- H_{n-1}, H_n are mutually generics over V .

Let $y_n = \tau[H_n]$. Notice that $y_n \in C_n$, and so $\lim_{n \rightarrow \infty} y_n = x$. For $n \in \mathbb{N}$, we have that $(p_n, p_n) \in H_n \times H_{n+1}$, and so there is $g_n \in G$ with $d(g_n, e) < \frac{1}{2^n}$ so that $g_n \cdot y_n = y_{n+1}$. We now have that

$$g_n \cdots g_0 \cdot y_0 = y_{n+1} \xrightarrow{n \rightarrow \infty} x.$$

We will show that there is a limit $\lim_{n \rightarrow \infty} g_n \cdots g_0 = g \in G$. In this case, $g \cdot y_0 = x$, concluding the proof of the claim. For the existence of the limit, we see that the sequence $(g_n \cdots g_0 : n \in \mathbb{N})$ is Cauchy, as

$$d(g_{n+1} \cdots g_n \cdots g_0, g_n \cdots g_0) = d(g_{n+1}, e) < \frac{1}{2^{n+1}}.$$

□

We note that the equality in the last equation is the only place in the proof where the invariance of d is used. We now proceed to describe the recursive construction. Given p_n and C_n , first we find a closed ball B of radius $< \frac{1}{2^n}$ and an extension q of p_n forcing that $\tau \in B$, and set $C_{n+1} = C_n \cap B$. Then, we extend q to a condition p_{n+1} satisfying condition (2). The following two lemmas show that this is possible.

Lemma 1.18. For any $p \in \mathbb{P}$ and $\epsilon > 0$ there is an extension $q \leq p$ and a ball B of radius $\leq \epsilon$ so that $q \Vdash \tau \in B$.

Proof. Fix a dense open subset of X , $D \subseteq X$ in the ground model. Let $H \subseteq \mathbb{P}$ be a generic filter containing p . Find $d \in D$ so that the distance between $\tau[H]$ and d is $< \epsilon/2$ and let B be the $\epsilon/2$ -ball around d . Since $\tau[H] \in B$, there is some condition in H , extending p , forcing that $\tau \in B$. □

Lemma 1.19 ([Kan08b, Lemma 17.4.1]). For any $p \in \mathbb{P}$ and $\epsilon > 0$ there is an extension $q \leq p$ so that (q, q) forces that there is $g \in G$ with $d(g, e) < \epsilon$ so that $g \cdot \tau_l = \tau_r$.

Proof. Fix p and ϵ , and assume for contradiction that no such q exists. Then for any q extending p there are extensions q_1, q_2 of q so that (q_1, q_2) forces that:

$$(\star) \text{ for any } g, \text{ if } g \cdot \tau_l = \tau_r \text{ then } d(g, e) \geq \epsilon.$$

³A naive hope would be to find a generic filter H so that $p_n \in H$ for all n , which would imply that $\tau[H] = x$. This is not generally possible.

Fix a sufficiently elementary countable elementary model M , and let $(D_n : n \in \omega)$, $(C_n : n \in \omega)$ enumerating all dense open subset in M of \mathbb{P} , $\mathbb{P} \times \mathbb{P}$, respectively. We may now construct a binary tree of conditions $\{p_s : s \in 2^{<\omega}\}$, as in Lemma 1.6, with $p_\emptyset = p$.

Any branch $b \in 2^\omega$ corresponds to a \mathbb{P} -generic filter $H_b = \{p_s : s \in 2^{<\omega}\}$ over M , and therefore to a point $x_b = \tau[H_n] \in X$. Moreover, for distinct $b, b' \in 2^\omega$, $H_b \times H_{b'}$ is $\mathbb{P} \times \mathbb{P}$ -generic over M , containing some condition forcing (\star) . Note that $x_b \mathop{E} x_{b'}$, since (\mathbb{P}, τ) is a stable virtual class.

Fix $b_0 \in 2^\omega$. For any $b, b' \neq b_0$,

$$g_{b_0, b'} \cdot g_{b_0, b}^{-1} \cdot x_b = x_{b'},$$

and so $d(g_{b_0, b'}, g_{b_0, b}) = d(g_{b_0, b'} \cdot g_{b_0, b}^{-1}, e) \geq \epsilon$. We conclude that $\{x_b : b_0 \neq b \in 2^\omega\}$ is an uncountable ϵ -separated family of members of G , a contradiction to G being separable. \square

\square

2. DIFFERENT VIRTUAL CLASSES

So far we focused on whether unpinned stable virtual classes exist or not. Next, following [LZ20], we study the space of all stable virtual classes: *the virtual realm*.

Definition 2.1 (Larson-Zapletal [LZ20, 2.1.4]). Let (\mathbb{P}, τ) and (\mathbb{Q}, σ) be stable virtual E -classes. Assume further that in any generic extension, given two filters $G \subseteq \mathbb{P}$, $H \subseteq \mathbb{Q}$ which are generic over the ground model

$$\tau[G] \mathop{E} \sigma[H].$$

In this case say that (\mathbb{P}, τ) and (\mathbb{Q}, σ) are *E -equivalent*⁴, denoted $(\mathbb{P}, \tau) \mathop{E} (\mathbb{Q}, \sigma)$.

A product forcing argument shows that $(\mathbb{P}, \tau) \mathop{E} (\mathbb{Q}, \sigma)$ if and only if

$$\mathbb{P} \times \mathbb{Q} \Vdash \tau \mathop{E} \sigma,$$

where we identify τ, σ with the corresponding $\mathbb{P} \times \mathbb{Q}$ -names which interpret the left and right generics, respectively. See [LZ20, 2.1.5].

Example 2.2. If (\mathbb{P}, τ) is pinned, $x \in X$ in the ground model is such that $\mathbb{P} \Vdash \tau \mathop{E} \check{x}$, then $(\mathbb{P}, \tau) \mathop{E} (\mathbb{Q}, \check{x})$, where \mathbb{Q} is the trivial (or any) forcing.

Example 2.3. In Example 1.3, $(\mathbb{P}_\alpha, \tau_\alpha), (\mathbb{P}_\beta, \tau_\beta)$ are not equivalent for $\alpha \neq \beta$. On the other hand, if $\mathbb{P} = \text{Col}(\omega, \mathbb{R})$, and τ is a \mathbb{P} -name for linear order of \mathbb{N} whose order type is ω_1 of the ground model, then $(\mathbb{P}_{\omega_1}, \tau_{\omega_1}) \mathop{E}_{\omega_1} (\mathbb{P}, \tau)$.

We now revisit Proposition 1.4. By collapsing more and more, we may get many different virtual classes, similar to Example 1.3.

Lemma 2.4. Suppose $E = E_a$ is a counterexample to the topological Vaught conjecture. Let \mathcal{S} be a set of stable virtual E -classes. Then there is a stable virtual E -class (\mathbb{P}, τ) which is not equivalent to any $(\mathbb{Q}, \sigma) \in \mathcal{S}$.

⁴In [LZ20] the extension of E to virtual classes is denoted by \bar{E} . Furthermore, there the pairs (\mathbb{P}, τ) are called *pins*, and a *virtual E -class* refers to the \bar{E} -equivalence class of (\mathbb{P}, τ) .

Proof. Fix a cardinal κ larger than $|\mathcal{S}|$ and $|\mathcal{P}(\mathbb{Q})|$ for any $\mathbb{Q} \in \mathcal{S}$, and let $\mathbb{P} = \text{Col}(\omega, \kappa)$. Working in a \mathbb{P} -generic extension, for each $(\mathbb{Q}, \sigma) \in \mathcal{S}$ we may find a filter $\dot{G}_{(\mathbb{Q}, \sigma)} \subseteq \mathbb{Q}$ which is generic over the ground model. Since E has uncountably many classes, in the generic extension, we may find $x \in X$ so that $x \notin \sigma[\dot{G}_{(\mathbb{Q}, \sigma)}]$ for any $(\mathbb{Q}, \sigma) \in \mathcal{S}$.

In the ground model, we may find a name τ so that

$$\mathbb{P} \Vdash \tau \notin \sigma[\dot{G}_{(\mathbb{Q}, \sigma)}],$$

where, for $(\mathbb{Q}, \sigma) \in \mathcal{S}$, $\dot{G}_{(\mathbb{Q}, \sigma)}$ is a \mathbb{P} -name for a filter in \mathbb{Q} generic over the ground model. It follows that $\mathbb{P} \times \mathbb{Q} \Vdash \tau \notin \sigma$. Finally, as in Proposition 1.4, there must be some $p \in \mathbb{P}$ so that $(\mathbb{P} \upharpoonright p, \tau)$ is a stable virtual class. $(\mathbb{P} \upharpoonright p, \tau)$ is not equivalent to any (\mathbb{Q}, σ) from \mathcal{S} , as required. \square

Given a virtual E -class (\mathbb{P}, τ) , say that \mathbb{P} is the *support* of (\mathbb{P}, τ) .

Definition 2.5 ([LZ20, 2.5.1]). Define $\kappa(E)$ to be the smallest cardinal κ so that every stable virtual E -class is equivalent to one supported by a poset of cardinality $< \kappa$. If no such κ exists, $\kappa(E) = \infty$. $\kappa(E)$ is called the *pinned cardinal* of E .

We may rephrase Lemma 2.4 as follows.

Corollary 2.6. If $E = E_a$ is a counterexample to the topological Vaught conjecture, then $\kappa(E) = \infty$.

For Borel equivalence relations, the pinned cardinal is bounded.

Theorem 2.7 ([LZ20, 2.5.6]). If E is a Borel equivalence relation, then $\kappa(E) < \beth_{\omega_1}$. More precisely, if E is $\mathbf{\Pi}_\alpha^0$, then $\kappa(E) \leq \beth_\alpha^+$.

Together with Proposition 1.7, we recover a corollary of Silver’s dichotomy.

Corollary 2.8. If $E = E_a$ is a Borel orbit equivalence relation, then E satisfies Vaught’s conjecture.

An ultimate extension of the above discussion is the following theorem of Hjorth.

Theorem 2.9 (Hjorth [Hjo97]). Suppose $E = E_a$ is a counterexample to the topological Vaught conjecture, $a: G \curvearrowright X$. Then there is a closed subgroup G' of G and a continuous homomorphism from G' onto S_∞ .

It is known that if G is CLI then it does not satisfy the conclusion of the theorem. Hjorth took this further and characterized when the topological Vaught conjecture fails on analytic sets. The proof of the theorem relies heavily on the existence of many inequivalent stable virtual equivalence classes (Corollary 2.6).

2.1. Some set theoretic questions about pinned cardinals.

Pinned cardinals. From a set theoretic point of view, understanding which cardinals are “definable”, in some reasonable sense, is of interest. As equivalence relations on Polish spaces are of particular interest, and are considered natural objects, we consider $\kappa(E)$ as interesting cardinals.

Question 2.10. Which cardinals κ are equal to $\kappa(E)$ for some analytic equivalence relation E ?

For example, for $0 < \alpha < \omega_1$, the cardinals \aleph_α and \beth_α^+ are realized as pinned cardinals of Borel equivalence relations. See [LZ20, 2.5.15] and [LZ20, 2.5.18].

Larson and Zapletal [LZ20, 2.5.10] proved that a measurable cardinal κ reflects the statement “ $\kappa(E) < \infty$ ” for analytic equivalence relations E . That is, if E is analytic and $\kappa(E) < \infty$, then $\kappa(E) < \kappa$.

Question 2.11. What is the least cardinal κ so that for any analytic equivalence relation E , if $\kappa(E) < \infty$ then $\kappa(E) < \kappa$?

Larson and Zapletal proved that the answer is between the first ω_1 -Erdos and the first measurable cardinal. Moreover, if one restricts to orbit equivalence relations E , the answer is precisely the first ω_1 -Erdos cardinal. See [LZ20, 2.5.8] and [LZ20, 2.5.9].

Supports for stable virtual classes. All the virtual classes above came from collapsing some cardinal to be countable.

Question 2.12. Which forcing notions \mathbb{P} can support a non-trivial stable virtual class for an analytic equivalence relation E ?

For orbit equivalence relations, such supporting poset must collapse.

Theorem 2.13 ([LZ20, 2.6.6]). If (\mathbb{P}, τ) is an unpinned stable virtual E_a -class, for an orbit equivalence relation E_a , then \mathbb{P} collapses \aleph_1 to be countable.

For non-orbit equivalence relations the situation is more subtle. For example, Namba forcing supports a non-trivial stable virtual class for a (necessarily non-orbit) analytic equivalence relation [LZ20, 2.6.8]. On the other hand, no *reasonable*⁵ forcing, in particular no proper forcing, can support a non-trivial stable virtual class for any analytic equivalence relation [LZ20, 2.6.2].

3. VIRTUAL CLASSES FOR ISOMORPHISM RELATIONS

Let us focus on isomorphism relations for countable structures: \cong_ϕ on $\text{Mod}(\phi)$ where ϕ is an $\mathcal{L}_{\omega_1, \omega}$ sentence (or a countable first order theory). These are the orbit equivalence relations which are induced by S_∞ . Much of the early work on analytic equivalence relations was focused on isomorphism relations, and is deeply connected to model theory.

Example 3.1. Let \mathcal{M} be an \mathcal{L} -structure, on some domain M . Let $\mathbb{P}_{\mathcal{M}} = \text{Col}(\omega, M)$ and let $\tau_{\mathcal{M}}$ be a $\mathbb{P}_{\mathcal{M}}$ -name for a structure on \mathbb{N} isomorphic to \mathcal{M} . Then $(\mathbb{P}_{\mathcal{M}}, \tau_{\mathcal{M}})$ is a stable virtual $\cong_{\mathcal{L}}$ -equivalence class.

If the set M is countable, we may find $x \in \text{Mod}(\mathcal{L})$ which is isomorphic to \mathcal{M} , and therefore $(\mathbb{P}_{\mathcal{M}}, \tau_{\mathcal{M}})$ is pinned. The converse is not necessarily true. Let \mathcal{L} be $\{=\}$ and M some uncountable set. Then $(\mathbb{P}_{\mathcal{M}}, \tau_{\mathcal{M}})$ is pinned, as τ is forced to be isomorphic to the structure \mathbb{N} .

Example 3.2. In the language of linear orders, the stable virtual classes from Example 1.3 are of the form $(\mathbb{P}_{\mathcal{M}}, \tau_{\mathcal{M}})$ where $\mathcal{M} = (\alpha, \in)$, for an ordinal α . For uncountable α , $(\mathbb{P}_{\mathcal{M}}, \tau_{\mathcal{M}})$ is unpinned.

⁵*Reasonable* is a technical notion, introduced by Foreman and Magidor [FM95], while studying definable counterexamples to the continuum hypothesis.

Example 3.3. Let $\mathcal{L} = \{U_n : n \in \mathbb{N}\}$, where each U_n is an unary predicate. Given a subset $A \subseteq 2^{\mathbb{N}}$, consider the structure $\mathcal{M}(A)$ with domain A so that $\mathcal{M}(A) \models U_n(x) \iff x(n) = 1$. Then $(\mathbb{P}_{\mathcal{M}(A)}, \tau_{\mathcal{M}(A)})$ is a stable virtual $\cong_{\mathcal{L}}$ -class, which is pinned if and only if A is countable.

An \mathcal{L} -structure \mathcal{M} with universe \mathbb{N} may be identified with a sequence $x \in (2^{\mathbb{N}})^{\mathbb{N}}$, where $x(n)(m) = 1 \iff \mathcal{M} \models U_m(n)$. The isomorphism relation $\cong_{\mathcal{L}}$ is induced by the natural S_{∞} action on $(2^{\mathbb{N}})^{\mathbb{N}}$, permuting the indices, which is identified with Example 1.14.

Definition 3.4 (see [URL17]). Let (\mathbb{P}, τ) be a stable virtual $\cong_{\mathcal{L}}$ -class. Say that (\mathbb{P}, τ) is *grounded* if there is some (possibly uncountable) model \mathcal{M} (in the ground model) so that

$$\mathbb{P} \Vdash \tau \cong \mathcal{M}.$$

Note that in this case (\mathbb{P}, τ) and $(\mathbb{P}_{\mathcal{M}}, \tau_{\mathcal{M}})$ are $\cong_{\mathcal{L}}$ -equivalent. Say that an equivalence relation E is *grounded* if every stable virtual E -class is grounded.

Note that a pinned equivalence relation is grounded.

Example 3.5. In Example 3.3, it can be shown that for any stable virtual $\cong_{\mathcal{L}}$ -class (\mathbb{P}, τ) , there is a set $A \subseteq 2^{\mathbb{N}}$ so that $\mathbb{P} \Vdash \tau \cong \mathcal{M}(A)$. So $\cong_{\mathcal{L}}$ is grounded.

In Example 1.3 it can be shown that for any stable virtual E_{ω_1} -class (\mathbb{P}, τ) , either τ is forced to be ill-founded, or there is an ordinal α so that $\mathbb{P} \Vdash \tau \cong (\alpha, \in)$. So E_{ω_1} is grounded. More generally, isomorphism relations for rigid structures are grounded [LZ20, 2.4.5].

More subtle instances of grounded isomorphism relations are given in [URL17], such as the isomorphism relations for the theories $\text{REF}(\text{bin})$ and $\text{REF}(\text{inf})$.

3.1. A theorem of Kaplan and Shelah and some applications. For an isomorphism relation \cong_{ϕ} , as it is an orbit equivalence relation, a support \mathbb{P} for an unpinned stable virtual E -class must collapse \aleph_1 , by Theorem 2.13.

Theorem 3.6 ([KS16, Corollary 4.3]). Let (\mathbb{P}, τ) be a stable virtual class for an isomorphism relation. If (\mathbb{P}, τ) is not grounded, then \mathbb{P} collapses \aleph_2 to be countable.

The proof goes through a translation between the question of a stable virtual isomorphism class being grounded and the following question: let T be a first order theory with dense isolated types, must T have an atomic model. The answer is positive if $|T| \leq \aleph_1$. This was proven in the 1970's independently by Knight, Kueker, and Shelah (see [KS16, Proposition 4.2]). Laskowski and Shelah [LS93] constructed a theory of size \aleph_2 with dense isolated types but no atomic model.

We present a few applications of Theorem 3.6. The theorem will be applied in the following form.

Corollary 3.7. If $(\text{Col}(\omega, \omega_1), \tau)$ is a stable virtual isomorphism class then there is a structure M (of size $\leq \aleph_1$) so that $\text{Col}(\omega, \omega_1) \Vdash \tau \cong M$.

3.1.1. Harrington's theorem. Going back to Vaught's conjecture, we recover the following theorem due to Harrington. See also [Lar17] and [KMS16].

Theorem 3.8 (Harrington). Suppose \cong_{ϕ} is a counterexample to the $\mathcal{L}_{\omega_1, \omega}$ -Vaught conjecture. Then ϕ has models of size \aleph_1 with arbitrary high Scott rank below ω_2 .

Proof. Fix an ordinal $\alpha < \omega_2$. Let $\mathcal{P} = \text{Col}(\omega, \omega_1)$. We repeat the arguments from Proposition 1.4, relying on the absoluteness of the statements involved.

In a \mathbb{P} -generic extension, α is countable, and ϕ has uncountably many models up to isomorphism, so we may find one with Scott rank greater than α . Let τ be a \mathbb{P} -name for such model. By Lemma 1.6, there is some $p \in \mathbb{P}$ so that $(\mathbb{P} \upharpoonright p, \tau)$ is a stable virtual \cong_ϕ -class.

By Theorem 3.6, $(\mathbb{P} \upharpoonright p, \tau)$ is grounded, so there is a structure \mathcal{M} so that $p \Vdash \tau \simeq \mathcal{M}$. As \mathcal{M} is forced to be countable by \mathbb{P} , then $|\mathcal{M}| \leq \aleph_1$. Finally, the Scott rank of \mathcal{M} is greater than α after forcing with \mathbb{P} , and therefore also in the ground model. \square

3.1.2. Abelian torsion groups and Ulm invariants. Countable Abelian torsion groups are classified by their Ulm-invariants. We focus here on reduced torsion abelian p -groups for a fixed p . A group is *reduced* if it has no non-trivial divisible subgroups. We briefly recall Ulm's classification result.

Given an abelian p -group C , define the transfinite sequence of its *Ulm subgroups* as follows. $C^0 = C$. Given C^α , for an ordinal α , $C^{\alpha+1} = \bigcap_{n \in \mathbb{N}} p^n C^\alpha$. For a limit ordinal δ , $C^\delta = \bigcap_{\alpha < \delta} C^\alpha$. This process eventually stabilizes at some stage $\kappa(C)$. The final subgroup $C^{\kappa(C)}$ is divisible, and is therefore trivial under our assumption. Let $\sigma(\lambda, k) \in \omega \cup \{\omega\}$ be the number of groups of order p^k appearing in a decomposition of the *Ulm factor* $C^\lambda / C^{\lambda+1}$ as a direct sum of cyclic groups.

Fact 3.9 (Ulm classification. See [Zip35, 7], [Fuc15, Theorem 1.6]). The map

$$C \mapsto \sigma(C) = (\sigma(\lambda, k) : \lambda < \kappa(C), k \in \mathbb{N}),$$

is a complete classification of countable reduced abelian p -groups.

Observation 3.10. The map $C \mapsto \sigma(C)$ is absolute for generic extensions. That is, $\sigma(C)$ remains the same when calculated in a forcing extension.

Given a sequence $A = (\sigma(\lambda, k) : \lambda < \kappa, k \in \mathbb{N})$, where κ is an ordinal and $\sigma(\lambda, k) \leq \omega$, we may ask if A is a classifying invariant for the classification above, that is, if there is a reduced abelian p -group C so that $\sigma(C) = A$. The countable case was answered by Zippin.

Theorem 3.11 (Zippin [Zip35]). Assume that

- κ is countable;
- for any $\lambda < \kappa$ there is some k so that $\sigma(\lambda, k) \neq 0$;
- for $\lambda + 1 < \kappa$ there are infinitely many k so that $\sigma(\lambda, k) \neq 0$.

Then there is a countable reduced abelian p -group C so that $\sigma(C) = (\sigma(\lambda, k) : \lambda < \kappa, k \in \mathbb{N})$.

Zippin's result was extended by Kulikov and Fuchs to uncountable groups [Fuc73, Theorem 76.1]. For cardinality \aleph_1 we get the following (see [CT19, Theorem 5.10]).

Theorem 3.12. Assume that

- $\kappa < \omega_2$;
- for any $\lambda < \kappa$ there is some k so that $\sigma(\lambda, k) \neq 0$;
- for $\lambda + 1 < \kappa$ there are infinitely many k so that $\sigma(\lambda, k) \neq 0$.

Then there is a reduced abelian p -group C of size $\leq \aleph_1$ so that $\sigma(C) = (\sigma(\lambda, k) : \lambda < \kappa, k \in \mathbb{N})$.

We recover this result directly from Zippin's theorem and Theorem 3.6.

Proof. Let $\mathbb{P} = \text{Col}(\omega, \omega_1)$. Since κ is countable after forcing with \mathbb{P} , it follows from Zippin's theorem that there is a \mathbb{P} -name τ for a countable reduced abelian p -group so that $\mathbb{P} \Vdash \sigma(\tau) = (\sigma(\lambda, k) : \lambda < \kappa, k \in \mathbb{N})$. That is, (\mathbb{P}, τ) is a stable virtual \cong_p -class. By Theorem 3.6, there is a reduced abelian group C so that $\mathbb{P} \Vdash \tau \cong C$. By the absoluteness of the Ulm invariants, it follows that $\sigma(C) = (\sigma(\lambda, k) : \lambda < \kappa, k \in \mathbb{N})$, as desired. \square

3.1.3. *Scott sets and models of PA.* Given a model M of Peano Arithmetic, seen as an end extension of the standard model, its *standard system* $\text{SSy}(M)$ is the collection of sets of the form $Y \cap \mathbb{N}$ where Y is definable in M (with parameters). The standard system of a model of PA always satisfies the following properties.

Definition 3.13. A set $S \subseteq \mathcal{P}(\mathbb{N})$ is a *Scott set* if

- If $A \in S$ and B is Turing reducible to A then $B \in S$;
- S is a Boolean algebra;
- If $T \in S$ codes an infinite binary tree then there is $b \in S$ coding an infinite branch of T .

It is an open question whether any Scott set can be realized as the standard system of a model of PA. For countable sets, Scott proved that it is the case.

Theorem 3.14 (Scott). Any countable Scott set can be realized as the standard system of a model of PA.

Knight and Nadel extended the result to sets of size \aleph_1 .

Theorem 3.15 (Knight and Nadel [KN82b]). Any Scott set of size \aleph_1 can be realized as the standard system of a model of PA.

See [Git08] for more background on this problem, and results beyond \aleph_1 . Here we provide a quick proof of Knight and Nadel's theorem, using Scott's theorem and Theorem 3.6. First, we introduce a related question.

Fix a countable language. Fix a recursive coding of formulas as natural numbers. We therefore identify theories and types as subsets of \mathbb{N} . Similarly, via a fixed recursive coding of binary sequences, we may identify binary trees as subsets of \mathbb{N} .

Definition 3.16. Given a Scott set S , say that a model M is *S -saturated* if

- For any $a_1, \dots, a_n \in M$, $\text{tp}(a_1, \dots, a_n) \in S$;
- For any $a_1, \dots, a_n \in M$ and any type $p(x, y_1, \dots, y_n) \in S$, if $p(x, a_1, \dots, a_n)$ is consistent then it is realized in M .

Fact 3.17. Given a Scott set S and an S -saturated model M of PA, the standard system of M is precisely S .

Scott's Theorem 3.14 follows from the following.

Theorem 3.18 (Scott, see [Wil80]). If S is a countable Scott set and T is a first order theory coded in S , then there is an S -saturated model of T .

This immediately extends to size \aleph_1 , using Theorem 3.6.

Theorem 3.19. If S is a Scott set of size $\leq \aleph_1$ and T is a first order theory coded in S , then there is an S -saturated model of T .

Proof. We will use the fact that a countable S -saturated model is unique up to isomorphism. This is because S -saturated models are ω -homogeneous. Let $\mathbb{P} = \text{Col}(\omega, \omega_1)$. Since S is a countable Scott set after forcing with \mathbb{P} , there is a \mathbb{P} -name τ for a countable S -saturated model of T . It follows that (\mathbb{P}, τ) is a stable virtual isomorphism class. By Theorem 3.6, there is a model M so that $\mathbb{P} \Vdash \tau \cong M$. Now M is the desired S -saturated model of T . \square

Theorem 3.19 was proved in the context of models of PA by Knight and Nadel [KN82a]. Note that if T is PA, the theorem produces a model of PA whose standard system is S , concluding the proof of Theorem 3.15.

4. BOREL REDUCIBILITY

Given equivalence relations E and F on Polish spaces X and Y , say that E is *Borel reducible* to F , denoted $E \leq_B F$, if there is a Borel measurable $f: X \rightarrow Y$ so that

$$x_1 E x_2 \iff f(x_1) F f(x_2), \text{ for any } x_1, x_2 \in X.$$

In this case, f induces a well defined injective map on the quotient spaces $X/E \hookrightarrow Y/F$. So ' $E \leq_B F$ ' is often thought of as 'the "Borel cardinality" of X/E is less than or equal to the "Borel cardinality" of Y/F '.

For an equivalence relation E so that all E -classes are Borel (this includes all Borel equivalence relations and all orbit equivalence relations), the statement ' E has countably many equivalence classes' is equivalent to ' $E \leq_B =_{\mathbb{N}}$ ', where $=_{\mathbb{N}}$ is the equality relation on \mathbb{N} . The statement ' E has perfectly many classes' is equivalent to ' $=_{\mathbb{R}} \leq_B E$ '. The study of Borel reducibility is an extension of the study of cardinalities of quotients of Polish spaces, beyond questions about the continuum hypothesis.

The celebrated E_0 -dichotomy, due to Harrington, Kechris, and Louveau, shows that for Borel equivalence relations there is a "next cardinality" after $=_{\mathbb{R}}$. A classification problem E is said to be **concretely classifiable** if $E \leq_B =_{\mathbb{R}}$. That is, if there is a Borel measurable complete classification using real numbers as invariants.

Theorem 4.1 ([HKL90]). Let E be a Borel equivalence relation. If E is not concretely classifiable, then $E_0 \leq_B E$.

For an introduction to Borel reducibility, the reader is referred to the books [Gao09, Kan08b], as well as the surveys [MR21, For18, KTD12, HK01, Kec99]. Here we focus on applications of virtual classes to Borel reducibility.

Lemma 4.2. Let E and F be analytic equivalence relations on Polish spaces X and Y respectively. Assume that $f: X \rightarrow Y$ is a Borel reduction of E to F . Let (\mathbb{P}, τ) be a stable virtual E -class. Let σ be a \mathbb{P} -name forced to be equal to $f(\tau)$. Then (\mathbb{P}, σ) is a stable virtual F -class. Moreover, (\mathbb{P}, τ) is pinned if and only if (\mathbb{P}, σ) is pinned.

Corollary 4.3. If F is pinned and $E \leq_B F$ then E is pinned.

More generally:

Corollary 4.4 ([LZ20, 2.5.4]). If $E \leq_B F$ then $\kappa(E) \leq \kappa(F)$.

Remark 4.5. For Lemma 4.2, and the corollaries, the assumption that the reduction is Borel was only used to conclude that this map is well defined, and is still

a reduction, in any generic extension. We will consider also reductions with this absoluteness property which may not be Borel.

4.1. The Friedman-Stanley jump operation. Given an equivalence relation E on X , the **Friedman-Stanley jump** of E is the equivalence relation E^+ on $X^{\mathbb{N}}$ defined by

$$x E^+ y \iff \forall n \exists m (x(n) E y(n)) \text{ and } \forall n \exists m (y(n) E x(m)),$$

equivalently, if $\{[x(n)]_E : n \in \mathbb{N}\} = \{[y(n)]_E : n \in \mathbb{N}\}$. The quotient $E^+/X^{\mathbb{N}}$ may be identified with $\mathcal{P}_{\aleph_0}(E/X)$, the countable powerset of E/X .

Friedman and Stanley [FS89] proved that this is a *jump operator* on Borel equivalence relations. That is, $E <_B E^+$ for any Borel equivalence relation E .

Example 4.6. Let $=^+$ be $(=_{\mathbb{R}})^+$, defined on $\mathbb{R}^{\mathbb{N}}$ by $x =^+ y \iff \{x(n) : n \in \mathbb{N}\} = \{y(n) : n \in \mathbb{N}\}$. $=^+$ is Borel bireducible with orbit equivalence relation from Example 1.14, and the isomorphism relation from Example 3.3. By Theorem 1.15 and Corollary 4.3, we have:

Corollary 4.7. $=^+$ is not Borel reducible to any orbit equivalence relation induced by a CLI group.

The iterated Friedman-Stanley jumps, $=^{+\alpha}$, are defined recursively, where $=^{+1}$ is $=^+$, $=^{+(\alpha+1)}$ is $(=^{+\alpha})^+$, and we take products at limit stages (see [Gao09, 12.2.6]). The Friedman-Stanley jumps play a central role in the theory of equivalence relations. A classification problem is considered *classifiable using countable sets of reals as complete invariants* if it is Borel reducible to $=^+$; “classifiable using countable sets of countable sets of reals as complete invariants” if it is Borel reducible to $=^{+2}$; and so on.

5. POTENTIAL INVARIANTS

For our treatment of equivalence relations which are classifiable by countable structures, their central property is the existence of an absolute complete classification. Recall that, for an equivalence relation E on X , a *complete classification* is a map $c: X \rightarrow I$ so that

$$x_1 E x_2 \iff c(x_1) = c(x_2), \text{ for all } x_1, x_2 \in X.$$

Definition 5.1. Say that c is an **absolute complete classification** if

- (1) The map $c: X \rightarrow I$ is defined in some set theoretic way ($c(x) = A \iff \psi(x, A)$, for some set theoretic formula ψ).
- (2) In any ZF extension $V \subseteq N$, ψ still defines a map which is a complete classification of E .
- (3) Given ZF models $V \subseteq N \subseteq W$, if x, A are in N and $N \models \psi(x, A)$, then $W \models \psi(x, A)$.

Similarly, a map $c: X \rightarrow I$ is an **absolute E -invariant map** if it satisfies the above with ‘complete classification’ replaced by ‘ E -invariant’, that is, $x_1 E x_2 \implies c(x_1) = c(x_2)$.

The third condition says that the calculation of the invariant of x , $A = c(x)$, does not change as we move to a generic extension. This is the crucial aspect making such classifications “reasonable”. For example, it prevents the classification $x \mapsto [x]_E$ from being “reasonable” when E is not a countable equivalence relation. In a sense,

it is a generous way of saying that the invariant $c(x)$ can be *computed from* x . That is, the computation is local, and does not depend on things unrelated to x , like some set theoretic truths in the universe.

For an isomorphism relation $\cong_{\mathcal{L}}$, the map

$$M \mapsto \varphi_M,$$

sending a model to its Scott sentence, is an absolute complete classification as above. Furthermore, in this case the invariants φ_M can be coded by hereditarily countable sets.

In more concrete scenarios, there is often a nice absolute complete classification, using hereditarily countable sets as invariants, of a more simple combinatorial nature. The seemingly technical conditions (2) and (3) above are always trivial to verify.

Example 5.2 (Absolute classification for Friedman-Stanley jumps). The map $\mathbb{R}^{\mathbb{N}} \rightarrow \mathcal{P}_{\aleph_0}(\mathbb{R})$, $x \mapsto \{x(n) : n \in \mathbb{N}\}$, is an absolute complete classification of $=^+$.

Consider the second Friedman-Stanley jump $=^{++}$ defined on the space $(\mathbb{R}^{\mathbb{N}})^{\mathbb{N}}$. The map $(\mathbb{R}^{\mathbb{N}})^{\mathbb{N}} \rightarrow \mathcal{P}_{\aleph_0} \mathcal{P}_{\aleph_0}(\mathbb{R})$,

$$x \mapsto \{\{x(n)(m) : m \in \omega\} : n \in \omega\}$$

is an absolute complete classification of $=^{++}$.

Similarly, there is an absolute complete classification of $=^{+\alpha}$ using invariants in $\mathcal{P}_{\aleph_0}^{\alpha}(\mathbb{R})$. See also [Sha24a, 7.4].

Example 5.3 (Absolute classification for CBERs). Suppose E is a countable Borel equivalence relation (CBER), that is, every E -class is countable. By the Feldman-Moore theorem [Gao09, Theorem 7.1.4], such E is the orbit equivalence relation induced by some Borel action of a countable group (recall Example 1.12). In this case, the map $x \mapsto [x]_E$ is an absolute complete classification. The point is that the E -class $[x]_E$ does not change in generic extensions, and therefore condition (3) holds.

Remark 5.4. Let E and F be analytic equivalence relation on Polish spaces X and Y respectively. Suppose F admits an absolute complete classification $y \mapsto B_y$ as above. Assume further that E is Borel reducible to F . Then E admits an absolute complete classification as well (with the same type of invariants).

Definition 5.5. Let E be an equivalence relation on a Polish space X and $x \mapsto A_x$ an absolute complete classification of E . Say that a set A is a **potential E-invariant** if in some forcing extension there is an x in X such that $A = A_x$. If A is a potential invariant for E , say that A is **trivial** if there is an x in the ground model such that $A = A_x$. In a given model in which there is some x so that $A = A_x$, we will say that A is **realized**.

Example 5.6. In Example 5.2, the potential invariants for $=^+$ (with this fixed absolute classification) are precisely all sets of reals. The potential invariants for $=^{+\alpha}$ are all sets in $\mathcal{P}^{\alpha}(\mathbb{R})$.

Potential invariants directly correspond to stable virtual classes, as explained below. This approach to study virtual classes is essentially equivalent to the one taken by Ulrich, Rast, and Laskowski [URL17]. See [PS24, Section 4].

Lemma 5.7. For E as above, there is a one-to-one correspondence between

- stable virtual E -classes (\mathbb{P}, τ) , and
- potential invariants A ,

such that (\mathbb{P}, τ) is pinned if and only if A is trivial. More specifically, a potential invariant A corresponds to (\mathbb{P}, τ) if and only if $\mathbb{P} \Vdash A_\tau = \dot{A}$.

Proof. Let (\mathbb{P}, τ) be a stable virtual E -class. Let $G_l \times G_r$ be $\mathbb{P} \times \mathbb{P}$ -generic and let x_l, x_r be the interpretations of τ according to G_l, G_r respectively. Since x_l and x_r are E -related, it follows that $A = A_{x_l} = A_{x_r}$. Furthermore, this set A is in the intersection $V[G_l] \cap V[G_r]$, which is equal to V by mutual genericity. If (\mathbb{P}, τ) is pinned there is $x \in V$ such that $x E x_l$. In particular, $A_x = A_{x_l} = A$, so A is a trivial. Conversely, if there is $x \in V$ with $A = A_x$ then x witnesses that (\mathbb{P}, τ) is pinned: given any \mathbb{P} -generic G over V , $A_{\tau[G]} = A = A_x$, so $\tau[G]$ is E -related to x .

Now let A be a potential invariant for E . By assumption there is a poset \mathbb{Q} , a generic G and $x \in V[G]$ such that $A = A_x$. Let τ be a \mathbb{Q} -name such that $\tau[G] = x$. Fix a condition $q \in \mathbb{Q}$ such that q forces that $A_\tau = A$ and define $\mathbb{P} = \mathbb{Q} \upharpoonright p$. Now (\mathbb{P}, τ) is a stable virtual E -class such that $\mathbb{P} \Vdash A_\tau = A$. □

A corollary of the proof is that the map $(\mathbb{P}, \tau) \mapsto A$ described there is a complete classification of all stable virtual E -classes, up to E -equivalence, where the classifying invariants are precisely the potential invariants for E .

Corollary 5.8. For E as above, the pinned cardinal $\kappa(E)$ is the smallest cardinal κ such that any potential E -invariant is trivial in a generic extension by a poset of size $< \kappa$.

Example 5.9. For $\alpha < \omega_1$, $\kappa(=^{+\alpha}) = \beth_\alpha^+$. In particular, for $\alpha < \beta$, $=^{+\beta} \not\leq_B =^{+\alpha}$.

See [LZ20, 2.5.5] for a proof of the full Friedman-Stanley theorem using these ideas. The original proof of Friedman and Stanley used Borel determinacy.

6. CARDINAL ARITHMETIC CONSIDERATIONS

In this section we consider some applications of Corollary 4.4 where the desired inequality between the pinned cardinals is not proved outright, but is consistent. This will suffice to conclude Borel irreducibility, as the latter is absolute.

6.1. Below $=^+$. The first example of an unpinned Borel equivalence relation was $=^+$, specifically, appearing in Hjorth's proof that $=^+$ is not Borel reducible to a CLI group action (see Example 4.6). Kechris asked if $=^+$ is the minimal example.

Question 6.1 (see [Kan08b, Question 17.6.1]). If E is an unpinned Borel equivalence relation, must $=^+$ be Borel reducible to E ?

This was refuted by Zapletal in [Zap11]. We present a counterexample. Consider the graph \mathcal{G} defined on $2^{\mathbb{N}}$ by⁶

$$x \mathcal{G} y \iff x \leq_T y \text{ or } y \leq_T x,$$

where \leq_T is the Turing reducibility relation on $2^{\mathbb{N}}$. One can verify that:

- (1) \mathcal{G} has a clique of size \aleph_1 ;
- (2) \mathcal{G} does not have cliques of size \aleph_2 .

⁶Zapletal used a different graph, see [Zap11, Fact 2.2]. This version was suggested by Clinton Conley.

Let $\mathcal{C} \subseteq (2^{\mathbb{N}})^{\mathbb{N}}$ be the set of all $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ such that $\{x(n) : n \in \mathbb{N}\}$ is a \mathcal{G} -clique. Consider the Borel equivalence relation E defined as $(=_{2^{\mathbb{N}}})^+ \upharpoonright \mathcal{C}$, the restriction of $(=_{2^{\mathbb{N}}})^+$ to the (invariant) Borel set \mathcal{C} .

Note that the map $x \mapsto \{x(n) : n \in \omega\}$ remains an absolute complete classification of E . The potential invariants are precisely all sets of reals $A \subseteq 2^{\mathbb{N}}$ which form a \mathcal{G} -clique. In particular, all the potential invariants are of size $\leq \aleph_1$, and so $\kappa(E) \leq \aleph_1^+$. Also, any uncountable \mathcal{G} -clique $A \subseteq 2^{\mathbb{N}}$ is a non-trivial potential invariant. It follows that E is not pinned, and $\kappa(E) = \aleph_1^+$. We claim that E is a counterexample to Question 6.1.

Claim 6.2. $=^+$ is not Borel reducible to E .

Proof. This follows from Corollary 4.4. More specifically, move to a forcing extension in which CH fails: $\beth_1 > \aleph_1$. In this extension, $\kappa(E) = \aleph_1^+ < \beth_1^+ = \kappa(=^+)$, and therefore $=^+ \not\leq_B E$. Finally, the statement ' $=^+ \leq_B E$ ' is absolute for forcing extensions, and therefore $=^+$ is not Borel reducible to E in the ground model as well. \square

6.2. Analytic equivalence relations. In the context of non-Borel analytic equivalence relation, especially non-orbit equivalence relations, the common notion of reducibility is *absolutely Δ_2^1 -reducibility*, see [Hjo00, Section 9.1].⁷ Corollary 4.4 still holds: $\kappa(E) \leq \kappa(F)$ if there is an absolute Δ_2^1 -reduction from E to F . In fact, this is true if there is a reduction which is an absolute map as in Definition 5.1, where 'complete classification' is replaced with 'reduction'. We will call such map an *absolute reduction*. We prove below a negative absolute reducibility result, which in particular implies a negative absolutely Δ_2^1 -reducibility result.

For analytic equivalence relations, the pinned cardinal may be unbounded, in which case it is not very useful. In this case, the following local notion may provide meaningful information.

Definition 6.3 ([LZ20, 2.5.1]). Let \mathbb{P} be a poset. Consider all stable virtual classes of the form (\mathbb{P}, τ) , up to E -equivalence. Let $\lambda(E, \mathbb{P})$ be the cardinality of this quotient.

From Lemma 4.2 we again see:

Lemma 6.4. If there is an absolute reduction from E to F , then $\lambda(E, \mathbb{P}) \leq \lambda(F, \mathbb{P})$

Suppose now E admits an absolute classification, $x \mapsto A_x$, as in Section 5. Recall Lemma 5.7, that there is a one-to-one correspondence taking a stable virtual E -class (\mathbb{P}, τ) to a potential invariant A so that $\mathbb{P} \Vdash A_\tau = \check{A}$.

Corollary 6.5. $\lambda(E, \mathbb{P})$ is the cardinality of the set of all potential invariants for E which become trivial after forcing with \mathbb{P} .

Example 6.6. For E_{ω_1} , the potential invariants are ordinals, the trivial ones are countable ordinals. The ordinals which become countable after forcing with $\text{Col}(\omega, \kappa)$ are precisely the ordinals below κ^+ , so $\lambda(E_{\omega_1}, \text{Col}(\omega, \kappa)) = \kappa^+$.

Example 6.7. For $=^+$ the potential invariants are sets of reals, the trivial ones are the countable sets of reals. The sets of reals which become countable after forcing with $\text{Col}(\omega, \omega_1)$ are precisely those of size $\leq \aleph_1$, so $\lambda(=^+, \text{Col}(\omega, \omega_1)) = \beth_1^{\aleph_1}$.

⁷See also the discussion in [Gao09, Section 9.2]. Δ_2^1 reductions are sometimes considered as well, in which case results often rely on absoluteness assumptions beyond ZFC.

We present here an application due to Calderoni and Thomas [CT19]. Consider the isomorphism relation \cong_{TA} , and the bi-embeddability relation \equiv_{TA} , for torsion abelian groups. Both equivalence relations have unbounded pinned cardinals. Calderoni and Thomas [CT19] proved that \equiv_{TA} is strictly below \cong_{TA} with respect to absolutely Δ_2^1 -reducibility.⁸

We sketch the proof that there is no absolute reduction from \cong_{TA} to \equiv_{TA} . For a fixed prime p , let \cong_p be the isomorphism relation restricted to torsion abelian p -groups. Following [CT19], we show that there is no absolute reduction from \cong_p to \equiv_{TA} . By Lemma 6.4 it suffices to show that $\lambda(\mathbb{P}, \cong_p) > \lambda(\mathbb{P}, \equiv_{\text{TA}})$, for some poset \mathbb{P} , in some forcing extension. We show below (Propositions 5.8 and 5.9 in [CT19]) that

- (1) $\lambda(\cong_p, \text{Col}(\omega, \omega_1)) \geq 2^{\aleph_1}$;
- (2) $\lambda(\equiv_{\text{TA}}, \text{Col}(\omega, \omega_1)) \leq \aleph_2^{\aleph_0}$.

Corollary 6.8. There is no absolute reduction of \cong_p to \equiv_p .

Proof. Working in a generic extension in which $2^{\aleph_0} = \aleph_1$ and $2^{\aleph_1} > \aleph_2$ we have

$$\lambda(\cong_p, \text{Col}(\omega, \omega_1)) \geq 2^{\aleph_1} > \aleph_2 = \aleph_2^{\aleph_0} \geq \lambda(\equiv_{\text{TA}}, \text{Col}(\omega, \omega_1)),$$

concluding the desired irreducibility. \square

Proof of (1).

Remark 6.9. Suppose $\sigma = (\sigma(\lambda, k) : \lambda < \kappa, k \in \mathbb{N})$ is a sequence satisfying the second two bullet points in Theorem 3.11 (that is, κ is not assumed to be countable). In any forcing extension in which κ is countable, σ satisfies all conditions of Theorem 3.11, and therefore σ is a classifying invariant in that extension. That is, σ is a potential classifying invariant for \cong_p .

For $\xi \in 2^{\omega_1}$, define $\sigma^\xi = (\sigma^\xi(\lambda, k) : \lambda < \omega_1, k \in \mathbb{N})$, $\sigma^\xi(\lambda, k) \in \{0, 1\}$, by

if $\xi(\lambda) = 0$, then $\sigma^\xi(\lambda, k) = 1 \iff k$ is even

if $\xi(\lambda) = 1$, then $\sigma^\xi(\lambda, k) = 1 \iff k$ is odd

Then $\{\sigma^\xi : \xi \in 2^{\omega_1}\}$ are distinct potential invariants for \cong_p which become trivial after collapsing ω_1 . By Corollary 6.5 we conclude that $\lambda(\cong_p, \text{Col}(\omega, \omega_1)) \geq 2^{\aleph_1}$.

Proof sketch of (2). Let \mathcal{A}_p be the domain of \equiv_p . Calderoni and Thomas show that there is an absolute complete classification of \equiv_p , $c_p : \mathcal{A}_p \rightarrow I_p$, with classifying invariants $I_p \subseteq \omega_1 \times \omega \times \omega$.

This in terms gives an absolute complete classification of \equiv_{TA} with classifying invariants $\prod_p I_p$, sending an abelian torsion group C to the sequence $(c_p(C_p) : p)$ where C_p is the p -primary component of C (see the proof of [CT19, Proposition 5.9]).

In this case, the potential invariants for \equiv_{TA} are members of $\prod_p (\text{Ord} \times \omega \times \omega)$, and those which are realized after forcing with $\text{Col}(\omega, \omega_1)$ are members of $\prod_p (\omega_2 \times \omega \times \omega)$. By Corollary 6.5, $\lambda(\equiv_{\text{TA}}, \text{Col}(\omega, \omega_1))$ is bounded by the cardinality of $\prod_p (\omega_2 \times \omega \times \omega)$, which is $\aleph_2^{\aleph_0}$.

Finally, we briefly describe the absolute classification c_p . Fix an enumeration $(R_{p,m} : m \in \omega)$ of representatives of the countably many bi-embeddability classes

⁸In [CT19] they consider Δ_2^1 -reductions, under the assumption that there exists a Ramsey cardinal, which ensures that such maps are absolute.

of countable Σ -cyclic p -groups. The map c_p sends a countable abelian p -group C to a triple (α, m, d) so that

- if the rank of $C^{\kappa(C)}$ is infinite then $\alpha = m = d = 0$;
- if the rank of $C^{\kappa(C)}$ is finite and $\kappa(C)$ is a limit ordinal, then $\alpha = \kappa(C)$, $d = \text{rk}(C^{\kappa(C)})$ and $m = 0$;
- if the rank of $C^{\kappa(C)}$ is finite and $\kappa(C)$ is a successor ordinal $\beta + 1$, then $\alpha = \kappa(C)$, $d = \text{rk}(C^{\kappa(C)})$ and m is chosen so that the Ulm factor $C^\beta/C^{\beta+1}$ is bi-embeddable with $R_{p,m}$.

The absoluteness of the map is evident. The fact that this is a complete classification of \equiv_p follows from [CT19, Theorem 4.11] (see the proof of [CT19, Proposition 5.11]).

7. SYMMETRIC MODELS

We saw that studying virtual equivalence classes in different models of set theory is useful to prove irreducibility results. For example, we were able to distinguish between various unpinned equivalence relations, by working in ZFC models with certain cardinal arithmetic assumptions, distinguishing their pinned cardinals.

Next we will apply similar ideas to distinguish between some equivalence relations which are pinned. The key is to work in choiceless models, where the equivalence relations may become unpinned.

Recall the examples from Section 1.1. The proof that $=_{\mathbb{R}}$ is pinned (Example 1.11), and the proof that a countable equivalence relation is pinned (Example 1.12) can be carried out in ZF. However, the proof in Example 1.13 used the axiom of choice. Similarly Theorem 1.15 used the axiom of choice, specifically DC. Below we use symmetric models in which DC, and various weak fragments of choice, fail.

In Section 7.2 we prove a Borel irreducibility result by considering the axiomatic strength (as a fragment of choice) of the statement ‘ E is pinned’. In Section 8 we consider finer questions of definability in symmetric models.

7.1. Abstract nonsense. Suppose A is a set in some generic extension of V . Let $V(A)$ be the minimal transitive model of ZF extending V and containing the set A . For example, this model can be written as the class directed union of $L(A, x)$ for $x \in V$. We will sometimes call this the model **generated by A** (over V).

There are similar ways of “generating models” from a set A : $L(A)$, $\text{HOD}_{\text{tc}\{A\}}$, $\text{HOD}_{V, \text{tc}\{A\}}$, as well as various symmetric models.⁹ For the A ’s we study here, all these models will have the same relevant properties. Also, in our examples the models $V(A)$ and $\text{HOD}_{V, \text{tc}\{A\}}$ will coincide.

The minimality of $V(A)$ implies the following ‘definable closure’ property.

Fact 7.1. The following holds in $V(A)$. For any set X , there is some formula ψ , parameters \bar{a} from the transitive closure of A and $v \in V$ such that X is the unique set satisfying $\psi(X, A, \bar{a}, v)$. Equivalently, there is a formula φ such that $X = \{x : \varphi(x, A, \bar{a}, v)\}$.

Proof. Work in the model $V(A)$. Consider the model $\text{HOD}_{V, \text{tc}\{A\}}$ of all hereditarily definable sets using parameters which are members of V , the transitive closure of

⁹See [Jec03, Chapter 13] for ordinal definability and [Jec03, Chapter 15], [Hal17, Chapter 17] for symmetric models.

A , as well as the set A itself. (That is, all sets definable as in the statement of the fact.) Then $\text{HOD}_{V, \text{tc}\{A\}}$ is a transitive model of ZF (see [Jec03, Theorem 13.26]), which extends V and contains A . By minimality of $V(A)$, we conclude that it is equal to $\text{HOD}_{V, \text{tc}\{A\}}$, and so every set is definable in this way. \square

We will be particularly interested in sets definable from A and parameters in V alone. We will primarily work with models $V(A)$ in which the axiom of choice fails.

Fact 7.2. If V satisfies AC and $A \subseteq V$ then $V(A)$ satisfies AC.

In particular, if A is a set of ordinals then $V(A)$ satisfies choice. If x is member of a Polish space, then x can be coded as a subset of \mathbb{N} , and therefore $V(x)$ satisfies choice as well.

7.2. $E_0^{\mathbb{N}}$ as an unpinned equivalence relation. First let us see how pinned equivalence relations become unpinned. Let $E_0^{\mathbb{N}}$ on $(2^{\mathbb{N}})^{\mathbb{N}}$ be defined by pointwise-equivalence. Note that the map

$$x \mapsto A_x = ([x(n)]_{E_0} : n \in \mathbb{N})$$

is an absolute complete classification for $E_0^{\mathbb{N}}$ (recall Example 5.3). The invariants are countable sequences of E_0 -classes.

Let \mathbb{P} be the poset of all finite approximations to a member of $(2^{\mathbb{N}})^{\mathbb{N}}$, ordered by extensions. That is, a condition in \mathbb{P} is a finite sequence of finite binary sequences. Fix $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ a \mathbb{P} -generic over V . Let $A_n = [x(n)]_{E_0}$ and $A = (A_n : n \in \mathbb{N})$. We also fix \mathbb{P} -names \dot{A}_n, \dot{A} for these objects.

Proposition 7.3. In $V(A)$, $\prod_{n \in \mathbb{N}} A_n = \emptyset$.

Claim 7.4. In $V(A)$, A is a potential invariant for $E_0^{\mathbb{N}}$.

It follows that in $V(A)$, A is a non-trivial potential invariant for $E_0^{\mathbb{N}}$.

Corollary 7.5. $E_0^{\mathbb{N}}$ is not pinned in $V(A)$.

Recall that if E is a countable Borel equivalence relation then “ E is pinned” is provable in ZF (Example 1.12).

Corollary 7.6. $E_0^{\mathbb{N}}$ is not Borel reducible to any countable Borel equivalence relation.

Proof. In $V(A)$, we see that $E_0^{\mathbb{N}}$ is not Borel reducible to any countable Borel equivalence relation, by Corollary 4.3. By absoluteness, the same is true in V . \square

Proof of Claim 7.4. In any extension of $V(A)$ where $\prod_{n \in \mathbb{N}} A_n$ is not empty, A is a classifying invariant. This is the case, for example, if we collapse $\bigcup_{n \in \mathbb{N}} A_n$ to be countable. More explicitly, we may add a generic choice function $x \in \prod_{n \in \mathbb{N}} A_n$ by finite approximations. \square

We now work towards the proof of Proposition 7.3. Let \mathbb{P}_n be the poset of all finite approximations to a member of $(2^{\mathbb{N}})^n$. We will use the following fact.

Claim 7.7. For each n ,

- the tuple $(x_0, \dots, x_{n-1}) \in (2^{\mathbb{N}})^n$ is \mathbb{P}_n -generic over V ;
- $x_n \in (2^{\mathbb{N}})^1$ is \mathbb{P}_1 -generic over $V[x_0, \dots, x_{n-1}]$.

This can be seen from the fact that the restriction map from \mathbb{P} to \mathbb{P}_n is a forcing projection (see [Cum10, 5.2]).

Lemma 7.8. Suppose $Z \in V(A)$, $Z \subseteq V$, and Z is definable using x_0, \dots, x_n , A , and a parameter in V . Then $Z \in V[x_0, \dots, x_n]$.

Proof. Fix a formula φ and a parameter $v \in V$ such that, in $V(A)$,

$$z \in Z \iff \varphi(z, A, x_0, \dots, x_n, v).$$

Claim 7.9. Suppose $p, q \in \mathbb{P}$ are conditions so that their restrictions to \mathbb{P}_n are compatible. Then p, q cannot force conflicting statements about $\varphi^{V(\dot{A})}(\dot{z}, \dot{A}, \dot{x}_0, \dots, \dot{x}_n, \dot{v})$.

Proof. Assume to the contrary, that $p \Vdash \varphi^{V(\dot{A})}(\dot{z}, \dots)$ and $q \Vdash \neg \varphi^{V(\dot{A})}(\dot{z}, \dots)$. Without loss of generality, assume that our generic $x \in (2^{\mathbb{N}})^{\mathbb{N}}$ extends p . Let $x' \in (2^{\mathbb{N}})^{\mathbb{N}}$ be the result of making finite changes to x_{n+1}, x_{n+2}, \dots so that x' extends q . Note that x' is \mathbb{P} -generic over V . Furthermore, $\dot{A}[x] = \dot{A}[x'] = A$, and $x'_i = x_i$ for $i \leq n$.

Working in $V[x]$, we conclude that

$$\varphi^{V(A)}(z, A, x_0, \dots, x_n, v).$$

However, working in $V[x']$ we conclude that

$$\neg \varphi^{V(A)}(z, A, x_0, \dots, x_n, v),$$

a contradiction. \square

Finally, we can define Z in $V[x_0, \dots, x_n]$ as all $z \in V$ such that there is some $p \in \mathbb{P}$ which agrees with x_0, \dots, x_n and such that $p \Vdash \varphi^{V(\dot{A})}(\dot{z}, \dot{A}, \dot{x}_0, \dots, \dot{x}_n, \dot{v})$. \square

Proof of Proposition 7.3. If $y \in \prod_{n \in \mathbb{N}} A_n$, then $y \in (2^{\mathbb{N}})^{\mathbb{N}}$ may be viewed as a subset of $\mathbb{N} \times \mathbb{N}$, and therefore as a subset of V . If additionally $y \in V(A)$, we conclude from the lemma, and Fact 7.1, that $y \in V[x_0, \dots, x_n]$ for some $n \in \mathbb{N}$. Since $y_{n+1} \in A_{n+1}$, it only differs from x_{n+1} on finitely many coordinates. It follows that $x_{n+1} \in V[y_{n+1}] \subseteq V[y] \subseteq V[x_0, \dots, x_n]$. This is a contradiction, as $x_{n+1} \in 2^{\mathbb{N}}$ is \mathbb{P}_1 -generic over $V[x_0, \dots, x_n]$. \square

This direction is further explored in [Sha24b]. For example, for countable Borel equivalence relations E, F so that E is (μ, F) -ergodic, for some probability measure μ , there is a symmetric model (a submodel of a random real extension) in which $E^{\mathbb{N}}$ is unpinned, yet $F^{\mathbb{N}}$ is pinned (see [Sha24b, Proposition 3.11]). Also, there is a model in which $E^{\mathbb{N}}$ is pinned, for any countable Borel equivalence relation E , yet the axiom ‘countable choice for countable sets of reals’ fails [Sha24b, Theorem 4.10].

8. DEFINABILITY OF POTENTIAL INVARIANTS

Let E and F analytic equivalence relations on Polish spaces X and Y respectively, and $f: X \rightarrow Y$ a Borel homomorphism from E to F , that is,

$$x E y \implies f(x) F f(y).$$

Then for a stable virtual E -class (\mathbb{P}, τ) , $(\mathbb{P}, f(\tau))$ is a stable virtual F -class. In other words, there is a map, definable using f as a parameter, sending stable virtual E -classes to stable virtual F -classes, and this holds uniformly in any generic extension.

Example 8.1. Assume further that E and F admit absolute classifications $x \mapsto A_x$ and $y \mapsto B_y$ respectively. Then stable virtual classes correspond to potential invariants (see Lemma 5.7). In this case, a Borel homomorphism f from E to F corresponds to a definable map, using as parameters f, E, F , and the complete classifications, sending potential E -invariants to potential F -invariants.

For concreteness, let us write it here. The map sends a potential E -invariant A to the unique set B satisfying

‘for some (equivalently, any) $x \in X$ in a generic extension,
if $A = A_x$ then $B = B_{f(x)}$ ’.

Furthermore, if $f: X \rightarrow Y$ is a reduction of E to F , then the map $A \mapsto B$ is injective on potential invariants. Moreover, in this case A can be defined from its image B as follows

‘for some (equivalently, any) $y \in Y$ in a generic extension,
if $B = B_y$ and $x \in X$ is such that $f(x) F y$, then $A = A_x$ ’.

Corollary 8.2. For E and F as above, if $E \leq_B F$ and A is a potential E -invariant in some generic extension, and B is the potential F -invariant corresponding to it, then $V(A) = V(B)$, where $V(A)$ is the minimal transitive extension of V containing A .

Example 8.3 ($E_0^{\mathbb{N}} \not\leq_B$ CBER). Let us prove again that $E_0^{\mathbb{N}}$ is not Borel reducible to a countable Borel equivalence relation, using the symmetric model $V(A)$ from Section 7.2.

Assume for contradiction that there is a Borel reduction from $E_0^{\mathbb{N}}$ to a countable Borel equivalence relation F . Since F is countable, the map $x \mapsto [x]_F$ is an absolute complete classification. By Corollary 8.2 we conclude that $V(A) = V(B)$ where B is an F -class, that is, $B = [y]_F$ for some $y \in V(B)$. It follows that $V(A) = V(y)$, and therefore $V(A)$ satisfies choice, by Fact 7.2, a contradiction to Proposition 7.3.

Similar to Example 8.1, we can extend an absolute E -invariant map to be defined on potential invariants.

Definition 8.4. Assume that E on X admits an absolute complete classification $x \mapsto A_x$ and $c: X \rightarrow I$ is an absolute E -invariant map. We define c on potential E -invariants as follows. Given a potential E -invariant A , define

$c(A)$ to be the unique $B \in I$ so that for some (equivalently, any) $x \in X$ in a generic extension, if $A = A_x$ then $B = c(x)$.

Note that the map on potential invariants $A \mapsto c(A)$ is defined uniformly, using the parameters used in the absolute definitions of $c: X \rightarrow I$ and of $x \mapsto A_x$.

9. ERGODICITY AND UNCLASSIFIABILITY

Recall that E_0 is defined on $2^{\mathbb{N}}$ as eventual equality between binary sequences. E_0 can be seen as the orbit equivalence relation induced by an action of $\bigoplus_{n \in \mathbb{N}} \mathbb{Z}_2$. A fundamental result in the theory of classification is that an E_0 is not *concretely classifiable*, that is, $E_0 \not\leq_B =_{\mathbb{R}}$. The standard argument relies on a basic ergodic theoretic technique.

Fact 9.1. Let Γ be a countable group and X a Polish space. Assume that $a: \Gamma \curvearrowright X$ is a generically ergodic action, that is, any Borel invariant set is either meager or comeager. Then any E_a -invariant Borel map $X \rightarrow \mathbb{R}$ is constant on a comeager set.

Similarly, in the measure theoretic context, if Γ is a countable group, $a: \Gamma \curvearrowright X$ is an ergodic action on a standard measure space (X, ν) (any invariant Borel set is either null or conull), then any E_a -invariant map $X \rightarrow \mathbb{R}$ is constant on a full measure set.

As each orbit is countable, we conclude that there is no Borel reduction from E_a to $=_{\mathbb{R}}$.

The action of $\bigoplus_{n \in \mathbb{N}} \mathbb{Z}_2$ on $2^{\mathbb{N}}$ described above is generically ergodic, and ergodic with respect to the coin-flipping measure on $2^{\mathbb{N}}$.

The goal of this section is to provide a generalized ergodicity criterion which

- implies unclassifiability by more complex invariants, such as sets of real numbers;
- is simple to apply, like the above ergodicity argument.

First, we view the unclassifiability of E_0 through an equivalent forcing point of view. Recall that a real number may be identified as a subset of ω . The unclassifiability of E_0 is true in greater generality. For example, in $L(\mathbb{R})$ there is no injective map from $2^{\mathbb{N}}/E_0$ into $\mathcal{P}(\alpha)$ for any ordinal α . (See [Hjo95], [CK11]. Also [Sha24a, 7.7])

Proposition 9.2. There is no absolute complete classification of E_0 using invariants which are sets of ordinals. In fact, any absolute invariant map from E_0 to sets of ordinals must send a comeager subset of $2^{\mathbb{N}}$ to the same set of ordinals.

The proposition relies on the weak homogeneity of Cohen forcing \mathbb{P} (adding a member of $2^{\mathbb{N}}$ by finite approximations): given any two conditions $p, q \in \mathbb{P}$ there is an automorphism γ of \mathbb{P} so that $\gamma \cdot p$ is compatible with q . Let $x \in 2^{\mathbb{N}}$ be a Cohen generic real over L . Feferman showed that there is no definable well ordering of the reals in $L[x]$. Levy showed that in fact any hereditarily ordinal definable set in $L[x]$ is in L (see [Kan06, Section 7]). The following is a mild generalization, allowing the the parameter $[x]_{E_0}$ in definitions.

Claim 9.3. Let $x \in 2^{\mathbb{N}}$ be a Cohen generic real over V . If $Z \in V[x]$ is a set of ordinals which is definable using $[x]_{E_0}$ and parameters from V alone, then $Z \in V$.

We omit the proof here, which is similar to the proof of Lemma 7.8. In fact, the claim follows from Lemma 7.8, if we view $x = x_0$ as part of a sequence x_0, x_1, \dots as in Section 7.2.

Proof of Proposition 9.2. Note that the map $x \mapsto [x]_{E_0}$ is an absolute complete classification of E_0 (Example 5.3). Assume that $x \mapsto B_x$ is an absolute E_0 -invariant map where B_x is always a set of ordinals. Let $x \in 2^{\mathbb{N}}$ be a Cohen generic real over V , and let $A = [x]_{E_0}$. As in Section 8 conclude that $B = B_x$ is definable from $[x]_{E_0}$ and parameters in V , and therefore $B \in V$, since B is a set of ordinals. We conclude that $B_x = \dot{B}$ is forced, and therefore there is a comeager set of $x \in 2^{\mathbb{N}}$ for which $B_x = B$. In particular, $x \mapsto B_x$ is not a complete classification. \square

We will repeat these ideas in much greater generality below, which will allow us to prove generalized unclassifiability results.

9.1. A comic relief. For $x \in 2^{\mathbb{N}}$, let x' be its Turing jump. Recall the following fact from computability theory.

Theorem 9.4. There exists a sequence of reals x_0, x_1, \dots so that $x_n \geq_T x'_{n+1}$.

Define $x \prec y \iff x' \leq_T y$. Define a graph \mathcal{G} on $2^{\mathbb{N}}$ by

$$x \mathcal{G} y \iff x \prec y \text{ or } y \prec x.$$

Define $\mathcal{C} \subseteq (2^{\mathbb{N}})^{\mathbb{N}}$ as all \mathcal{G} -cliques (a Borel set), and E as $(=_{2^{\mathbb{N}}})^+ \upharpoonright \mathcal{C}$.

Let $(x_\alpha : \alpha < \omega_1)$ be a \leq_T -increasing sequence so that $x_{\alpha+1} = (x_\alpha)'$. Then $\{x_\alpha : \alpha < \omega_1\}$ is a non-trivial potential invariant for E , and so E is not pinned. In particular, $E \not\leq_B =_{\mathbb{R}}$ (recall Example 1.11). Since E is Borel, it follows from Theorem 4.1 that $E_0 \leq_B E$.

Proof of Theorem 9.4. Assume for contradiction that the theorem fails, then the relation \prec is wellfounded. For $x \in X$, let A_x be an enumeration of $\{x_n : n \in \mathbb{N}\}$ of order type $\text{o.t.}(\prec \upharpoonright \{x_n : n \in \mathbb{N}\})$. Then $x \mapsto A_x$ is an absolute complete classification of E , using $(2^{\mathbb{N}})^{<\omega_1}$ as invariants. The latter may be coded as members of $2^{<\omega_1}$. We conclude that E_0 admits an absolute complete classification using sets of ordinals as invariants, contradicting Proposition 9.2.¹⁰ \square

9.2. Generalized ergodicity.

Definition 9.5. Let \mathbb{P} be a forcing poset, τ a \mathbb{P} -name. Say that τ is **\mathbb{P} -ergodic** if for any $p, q \in \mathbb{P}$ there is an automorphism γ of \mathbb{P} so that $\gamma \cdot p$ is compatible with q and γ preserves τ , that is, for any \mathbb{P} -generic G , $\tau[G] = \tau[\gamma \cdot G]$.

Example 9.6. Let τ be a canonical \mathbb{P} -name for the emptyset. Then τ is \mathbb{P} -ergodic if and only if \mathbb{P} is weakly homogeneous.

Definition 9.7. Say that a \mathbb{P} -name τ is **non-trivial** if there are two generic filters G, H so that $\tau[G] \neq \tau[H]$. Equivalently, τ is non-trivial if it is not forced to be a member of the ground model.

Example 9.8. Let (X, ν) be a standard measure space and $\Gamma \curvearrowright X$ an ergodic group action by measure preserving transformations. Let \mathbb{P} be Random real forcing on X , \dot{x} the name for the generic random real. Then $\Gamma \cdot \dot{x}$ is \mathbb{P} -ergodic.

Example 9.9. Let X be a Polish space and $\Gamma \curvearrowright X$ a generically ergodic action by homeomorphisms. Let \mathbb{P} be Cohen forcing on X and \dot{x} the name for the generic Cohen real. Then $\Gamma \cdot \dot{x}$ is \mathbb{P} -ergodic.

Theorem 9.10 (ZF). Let \mathbb{P} be a forcing poset, \dot{A} a \mathbb{P} -name for a potential E -invariant. Assume that \dot{A} is \mathbb{P} -ergodic. If $c: X \rightarrow I$ is an absolute E -invariant map, where in any \mathbb{P} -generic extension the members of I are subsets of V , then there is a fixed $B \in I$ in the ground model so that $c(\dot{A}) = \dot{B}$ is forced by \mathbb{P} . (We may apply c to a potential invariant as in Section 8.)

Proof. Let σ be a \mathbb{P} -name so that it is forced by \mathbb{P} that $c(\dot{A}) = \sigma$. By assumption, σ is forced to be a subset of V .

Claim 9.11. For any b in V , if some condition in \mathbb{P} forces that $\check{b} \in \sigma$ then any condition in \mathbb{P} forces that $\check{b} \in \sigma$.

Given the claim, we may define B to be the set of all b so that $\check{b} \in \sigma$ is forced, and conclude that $c(\dot{A}) = \sigma = \dot{B}$ is forced by \mathbb{P} , as required.

To prove the claim, assume towards a contradiction that there are two conditions $p, q \in \mathbb{P}$ so that $p \Vdash \check{b} \in \sigma$ and $q \Vdash \check{b} \notin \sigma$. By assumption, there is an automorphism γ of \mathbb{P} so that $\gamma \cdot p$ is compatible with q , and γ preserves \dot{A} . Since $\sigma = c(\dot{A})$ is forced, γ preserves σ as well. We conclude that $\gamma \cdot p \Vdash \check{b} \in \sigma$ and $q \Vdash \check{b} \notin \sigma$, contradicting that they are compatible. \square

There are three types of generalizations here, beyond what is needed for Proposition 9.2, which will be crucial in the applications below:

- (1) that A is a potential invariant (not necessarily realized in the given model);

¹⁰Another way of reaching the contradiction is to show that the classification $x \mapsto A_x$ is an ‘Ulm-type classification’ (see [HK95]), contradicting the fact that E_0 does not admit such classification.

- (2) that invariants in I are arbitrary subsets of the ground model (not just ordinals);
- (3) that the result will be applied in specifically designed models of ZF.

A typical application of the theorem will be to prove irreducibility to some $=^{+\alpha}$, as follows. Here \mathcal{P}^α denotes the iterated powerset operation, and $\mathcal{P}(\mathbb{N})$ is identified with \mathbb{R} .

Theorem 9.12. Let \mathbb{P} be a forcing poset, \dot{A} a non-trivial \mathbb{P} -name for a potential E -invariant. Assume that \dot{A} is \mathbb{P} -ergodic. Assume further that forcing with \mathbb{P} adds no new subsets of $\mathcal{P}^\alpha(\mathbb{N})$. Then there is no absolute complete classification of E using subsets of $\mathcal{P}^\alpha(\mathbb{N})$ as invariants. In particular, $E \not\leq_B =^{+\alpha+1}$.

For example, if \mathbb{P} adds no real numbers and \dot{A} is an ergodic, non-trivial \mathbb{P} -name for a potential E -invariant, then $E \leq_B =^+$.

Proof. Assume for contradiction that $c: X \rightarrow I$ is a complete classification of E where I consists of subsets of $\mathcal{P}^\alpha(\mathbb{N})$. By assumption, the invariants in I are subsets of V in any \mathbb{P} -generic extension. By Theorem 9.10 there is a fixed set B in V so that $c(\dot{A}) = \dot{B}$ is forced by \mathbb{P} . Finally, for any two \mathbb{P} -generics G, H over V , as $c(\dot{A}[G]) = c(\dot{A}[H])$ and c is a complete classification, we conclude that $\dot{A}[G] = \dot{A}[H]$, contradicting the assumption that \dot{A} is not trivial. \square

9.3. Irreducibility to $=^+$. We present here two examples proving irreducibility to $=^+$ (unclassifiability by countable sets of reals).

9.3.1. A proof of $=^{++} \not\leq_B =^+$. Let \mathbb{P} be the poset of all countable partial functions $p: \mathbb{R} \rightarrow 2$, ordered by extension. We identify a generic filter with a ‘generic function’ $\mathbb{R} \rightarrow 2$, and in turn with a ‘generic subset’ $G \subseteq \mathbb{R}$. Let \dot{B} be the \mathbb{P} -name for the set $B = \{H \subseteq \mathbb{R} : H \Delta G \text{ is countable}\}$, all subsets of \mathbb{R} which have countable symmetric difference with G . Note that in the extension $V[G]$, B is a potential invariant for $=^{++}$, as it is a set of sets of reals. We will use the following basic facts:

- Forcing with \mathbb{P} does not add reals;
- The generic subset G is not in the ground model.

Claim 9.13. \dot{B} is \mathbb{P} -ergodic.

Proof. Given $p, q \in \mathbb{P}$, consider the automorphism of \mathbb{P} acting on $r \in \mathbb{P}$ by flipping the value of $r(x)$ for any $x \in \text{dom } p \cap \text{dom } q$ for which $p(x) \neq q(x)$. Since only countably many values of G are changed, the automorphism fixes \dot{B} . \square

We conclude from Theorem 9.10 that there is no absolute complete classification of $=^{++}$ using invariants which are subsets of V in any \mathbb{P} -generic extension. In any \mathbb{P} -generic extension, since no reals are added, a set of reals is a subset of V .

Corollary 9.14. There is no absolute complete classification of $=^{++}$ using invariants which are sets of reals. In particular, $=^{++} \not\leq_B =^+$.

9.3.2. Subgroups of $(\mathbb{R}, +)$. Let $\mathcal{A} \subseteq \mathbb{R}^{\mathbb{N}}$ be all injective sequences $(x_n : n \in \mathbb{N})$ so that $x_0 = 0$ and $(\{x_n : n \in \mathbb{N}\}, +)$ is a subgroup of $(\mathbb{R}, +)$. \mathcal{A} is a closed subset of $\mathbb{R}^{\mathbb{N}}$, equipped with the product topology.

Definition 9.15. Define the equivalence relation $\sim_{\mathcal{A}}$ on \mathcal{A} as follows.

$$(x_n : n \in \mathbb{N}) \sim_{\mathcal{A}} (y_n : n \in \mathbb{N}) \iff$$

$\exists \lambda > 0$ so that the map $x \mapsto \lambda \cdot x$ is a bijection from $\{x_n : n \in \mathbb{N}\}$ to $\{y_n : n \in \mathbb{N}\}$.

This equivalence relation was studied in [CMRS23]. The main motivation is that $\sim_{\mathcal{A}}$ is equivalent to the classification problem for countable Archimedean groups. (See Proposition 3.1 in [CMRS23] and the discussion following it.) In particular, it is established in [CMRS23, Section 4] that

$$=^+ <_B \sim_{\mathcal{A}} <_B =^{++}$$

We provide here a proof of the lower bound.

Proposition 9.16. $\sim_{\mathcal{A}}$ is not Borel reducible to $=^+$.

This lower bound was recently used by [EGL24], where they study the complexity of classification for extremely amenable groups.

We identify a member $(x_n : n \in \mathbb{N}) \in \mathcal{A}$ with the subgroup $G = \{x_n : n \in \mathbb{N}\}$. Given an additive subgroup G of \mathbb{R} , define

$$G/a = \{g/a : g \in G\} \text{ and } A_G = \{G/a : a \in G \setminus \{0\}\}.$$

Fact 9.17 (See [CMRS23, Proposition 3.14]). The map

$$(x_n : n \in \mathbb{N}) \mapsto A_{\{x_n : n \in \mathbb{N}\}}$$

is an absolute complete classification of $\sim_{\mathcal{A}}$.

We will apply Theorem 9.10 using this classification. Note that a classifying invariant is a countable set of countable subgroups of $(\mathbb{R}, +)$.

Remark 9.18. We note that $\kappa(\sim_{\mathcal{A}}) = \kappa(=^+) = |\mathbb{R}|^+$, so the methods of sections 4 and 6 cannot be used to prove Proposition 9.16.

Suppose A is a potential $\sim_{\mathcal{A}}$ -invariant, using the above classification. Then A is a set of sets of reals and there is $G \in A$ so that $A = \{G/a : a \in G \setminus \{0\}\}$ (though A and G may be uncountable). Since $G \subseteq \mathbb{R}$, after collapsing $|\mathbb{R}|$ to be countable we have some $(x_n : n \in \mathbb{N}) \in \mathcal{A}$ enumerating G , so that A is the classifying invariant of $(x_n : n \in \mathbb{N})$. That is, any potential $\sim_{\mathcal{A}}$ -invariant is trivial after collapsing $|\mathbb{R}|$ to be countable. It follows from Corollary 5.8 that $\kappa(\sim_{\mathcal{A}}) \leq |\mathbb{R}|^+$. Note that $A = \{\mathbb{R}\} = \{\mathbb{R}/a : a \in \mathbb{R} \setminus \{0\}\}$ is a potential $\sim_{\mathcal{A}}$ -invariant, and so $\kappa(\sim_{\mathcal{A}}) = |\mathbb{R}|^+$

We introduce some terminology from [CMRS23, 4.3].

Given a set $S \subseteq \mathbb{R}$, define $D(S)$ to be the set of all numbers of the form $a_1^{l_1}, \dots, a_n^{l_n}$, where l_1, \dots, l_n are integers and a_1, \dots, a_n are in S . Let Y be a perfect set of reals so that any $a_1, \dots, a_n \in Y$ are algebraic independent over $Y \setminus \{a_1, \dots, a_n\}$. In particular, for any two subsets $S_1, S_2 \subseteq Y$,

$$D(S_1) \cap D(S_2) = D(S_1 \cap S_2).$$

Definition 9.19. Let \mathbb{P} be the poset of conditions $p = (S_p, G_p)$ where S_p is a countable subset of Y and G_p is a countable subset of $D(S_p)$. For $p, q \in \mathbb{P}$ say that q extends p if $S_q \supseteq S_p$ and $G_q \cap D(S_p) = G_p$.

\mathbb{P} is a partial order with maximal element (\emptyset, \emptyset) . Given a generic filter $F \subseteq \mathbb{P}$, let G be the additive subgroup of \mathbb{R} generated by $\bigcup_{p \in F} G_p$. Note that in the extension A_G is a potential invariant for $\sim_{\mathcal{A}}$: in any further generic extension in which G is countable, A_G is equal to A_x for any injective enumeration $x \in \mathcal{A}$ of G .

Claim 9.20. For any generic filter F , $G \notin V$. In particular $A_G \notin V$.

Proof. Fix a condition $p \in \mathbb{P}$. We need to find two extensions q_1, q_2 of p and some $x \in \mathbb{R}$ so that $q_1 \Vdash \check{x} \in \dot{G}$ and $q_2 \Vdash \check{x} \notin \dot{G}$. Since S_p is countable, we may find some $x \in Y \setminus S_p$. By the algebraic independence of the members of Y , it follows that $x \notin D(S_p)$. Define

$$S_{q_1} = S_{q_2} = S_p \cup \{x\}, \quad G_{q_1} = G_p \cup \{x\}, \quad G_{q_2} = G_p.$$

Then $q_1 = (S_{q_1}, G_{q_1})$ and $q_2 = (S_{q_2}, G_{q_2})$ are as required. \square

Claim 9.21. \mathbb{P} is countably closed. In particular, no reals are added when forcing with \mathbb{P} .

Claim 9.22. \dot{A}_G is \mathbb{P} -ergodic.

Proof. Fix $a \in Y$. Let $\mathbb{P}_a = \{p \in \mathbb{P} : a \in S_p\}$, a dense open subset of \mathbb{P} . Define $\pi_a(p) = (S_p, a \cdot G_p)$. Then π_a is an automorphism of \mathbb{P}_a , which preserves \dot{A}_G .

Given any $p, q \in \mathbb{P}$, fix some $a \in Y \setminus (S_p \cup S_q)$. By the algebraic independence of the members of Y , it follows that $a \cdot D(S_p) \cap D(S_q) = \emptyset$, and therefore $\pi_a(p)$ and q are compatible. \square

We conclude from Theorem 9.12 that $\sim_{\mathcal{A}}$ is not Borel reducible to $=^+$.

9.4. Forcing over choiceless models. Let $\mathcal{P}(On)$ denote all subsets of ordinals, $\mathcal{P}^2(On)$ denote all sets of sets of ordinals, and so on. As in Proposition 9.2, often times an irreducibility to $=^+$ can be strengthened to ‘unclassifiability using sets of sets of ordinals as invariants’. We show this for $=^{++}$ in Corollary 9.26 below. First, note that in Theorem 9.12, we may replace \mathbb{N} by some ordinal, and the same proof works. Furthermore, the proof did not use the axiom of choice.

Theorem 9.23 (ZF). Let \mathbb{P} be a forcing poset, \dot{A} a non-trivial \mathbb{P} -name for a potential E -invariant. Assume that \dot{A} is \mathbb{P} -ergodic. Assume further that forcing with \mathbb{P} adds no new subsets of $\mathcal{P}^\alpha(On)$. Then there is no absolute complete classification of E using subsets of $\mathcal{P}^\alpha(On)$ as invariants.

When $\alpha = 0$ we recover Proposition 9.2. When $\alpha = 1$ we conclude a generalization to ‘irreducibility to $=^+$ ’: unclassifiability by sets of ordinals as invariants.

For $\alpha \geq 1$, for the hypothesis in Theorem 9.23 to hold, the axiom of choice must fail. We present below some instances of this scenario, and how it naturally leads to an irreducibility result.

9.4.1. Forcing over the basic Cohen model. Recall the ‘the basic Cohen model’ in which the axiom of choice fails (see [Kan08a]). Let \mathbb{C} be Cohen forcing for adding a member to $\mathbb{R}^{\mathbb{N}}$. A condition in \mathbb{C} is a finite sequence p so that $p(i)$ is a non-empty open subset of \mathbb{R} , for each i in its domain. Say that a condition p extends q if the domain of p extends the domain of q and $p(i) \subseteq q(i)$ for each $i \in \text{dom } q$.¹¹

Let $x \in \mathbb{R}^{\mathbb{N}}$ be \mathbb{C} -generic over V , $A = \{x(n) : n \in \mathbb{N}\}$ and consider the model $V(A)$. This model has been extensively studied in the literature. A basic fact is that in $V(A)$, A is a Dedekind-finite set, that is, there is no sequence of distinct members of A . In particular, there is no $y \in \mathbb{R}^{\mathbb{N}}$ enumerating A . It follows that A is a non-trivial potential invariant for $=^+$, in $V(A)$.

¹¹ \mathbb{C} is (equivalent to) the topological version of Cohen forcing for the space $\mathbb{R}^{\mathbb{N}}$. See [Jec03, Section 26].

Working in $V(A)$, let \mathbb{P} be the poset of all finite partial functions $p: A \rightarrow 2$, ordered by extension. We identify a generic filter with a ‘generic function’ $A \rightarrow 2$, and in turn with a ‘generic subset’ $G \subseteq A$. Let \dot{B} be a name for the set $B = \{H \subseteq A : H \Delta G \text{ is finite}\}$, all subsets of A which have finite symmetric difference with G . Note that, in the extension $V(A)[G]$, B is a potential invariant for $=^{++}$, as it is a set of sets of reals.

Claim 9.24 (In $V(A)$). \dot{B} is \mathbb{P} -ergodic.

Proof. Given a finite subset $F \subseteq A$, there is an automorphism π_F of \mathbb{P} , flipping the values on F . That is, for $a \in F$, $p \in \mathbb{P}$, and $a \in \text{dom } p$, $\pi_F(p)(a) = 1 - p(a)$. Note that π_F preserves \dot{B} . Finally, for any two conditions $p, q \in \mathbb{P}$ there is a finite set F so that $\pi_F(p)$ is compatible with q . \square

We conclude from Theorem 9.10 that, in $V(A)$, there is no absolute complete classification of $=^{++}$ using invariants which are subsets of $V(A)$ in any \mathbb{P} -generic extension. The key point here is the following, due to Monro [Mon73].

Lemma 9.25. Forcing with \mathbb{P} adds no subsets of V to $V(A)$.

In particular, no sets of ordinals are added. It follows that any set of sets of ordinals, in a \mathbb{P} -generic extension of $V(A)$, is a subset of $V(A)$.

Corollary 9.26. There is no absolute complete classification of $=^{++}$ using invariants which are sets of sets of ordinals. In particular, $=^{++} \not\leq_B =^+$.

We prove Lemma 9.25 below. First, we present a more subtle irreducibility result, which does not follow from the methods considered earlier in this note.

9.4.2. *An equivalence relation between $=^+$ and $=^{++}$.*

Definition 9.27. Consider the space $X = \mathbb{R}^{\mathbb{N}} \times 2^{\mathbb{N}}$. For $(x, t) \in X$, define $A_x = \{x(n) : n \in \mathbb{N}\}$ and $G_{(x,t)} = \{x(n) : t(n) = 1\}$. Define

$$(x, t) E (y, s) \iff A_x = A_y \text{ and } G_{(x,t)} \Delta G_{(y,s)} \text{ is finite.}$$

This can be seen as a version of E_0 , with the base space \mathbb{N} replaced by $\mathbb{R}^{\mathbb{N}}/ =^+$. For $(x, t) \in X$, define

$$B_{(x,t)} = \{G \subseteq A_x : G \Delta G_{(x,t)} \text{ is finite}\}.$$

Note that the map $x \mapsto B_x$ is an absolute complete classification of E .

It is readily seen that $=^+ \leq_B E \leq_B =^{++}$. E is an example of an equivalence relation with *potential complexity* $D(\mathbf{\Pi}_3^0)$.¹² It follows from the results in [HKL98] that $E <_B =^{++}$. We show here that the reduction $=^+ <_B E$ is strict as well.

Working again in the basic Cohen model $V(A)$, we see that \dot{B} is a name for a potential invariant for E . The same argument above shows:

Corollary 9.28. There is no absolute complete classification of E using invariants which are sets of sets of ordinals. In particular, $E \not\leq_B =^+$.

Proof. This follows from Claim 9.24 and 9.25. \square

¹²See [HKL98] for a treatment of potential complexity. The equivalence relation E here is Borel reducible to the equivalence relation $\cong_{3,1}^*$ there. See also [Sha21, Sha].

Remark 9.29. One can verify that $\kappa(E) = |\mathbb{R}|^+ = \kappa(=^+)$, so the methods of sections 4 and 6 cannot be used to prove the irreducibility $E \not\leq_B =^+$. Furthermore, the method from Section 9.3 (collapsing over a ZFC model) does not seem to work here.

As discussed right before Section 9.4.1, the key point here is Lemma 9.25, witnessing the failure of choice in $V(A)$ in a strong way. We will use the following Continuity Lemma. See [Fel71, p.133], also [CMRS23, p.19] for a proof in this presentation of the basic Cohen model.

Lemma 9.30 (Continuity Lemma). Let ϕ be a formula, $\bar{a} = a_0, \dots, a_{n-1}$ a finite sequence of distinct members of A , and $v \in V$. Suppose $\phi(A, \bar{a}, v)$ holds in $V(A)$. Then there are open sets U_0, \dots, U_{n-1} such that $a_i \in U_i$ and for any $\bar{b} = b_0, \dots, b_{n-1}$ consisting of distinct elements from A , if $b_i \in U_i$ for all $i \leq n-1$, then $\phi(A, \bar{b}, v)$ holds in $V(A)$.

Proof of Lemma 9.25. Suppose τ is a \mathbb{P} -name for a subset of V , $\tau \in V(A)$. As in Fact 7.1, let $\bar{a} \subseteq A$ be finite, $w \in V$ a parameter, and ψ a formula so that in $V(A)$, τ is defined as the unique solution to $\psi(\tau, A, \bar{a}, w)$.

We will show that for any condition $p \in \mathbb{P}$ and any $v \in V$, if p forces $\check{v} \in \tau$ then $p \upharpoonright \bar{a}$ forces the same. It then follows that for any generic filter $G \subseteq P$ over $V(A)$, the values of $\tau[G]$ are determined by the single condition $q = G \upharpoonright \bar{a}$. That is, $\tau[G]$ may be defined in $V(A)$ as the set of all v for which $q \Vdash \check{v} \in \tau$.

Fix a condition $p \in \mathbb{P}$ and $v \in V$ so that $p \Vdash \check{v} \in \tau$. Assume that the domain of p is of the form \bar{a}, \bar{b} , with \bar{b} disjoint from \bar{a} . By the Continuity Lemma, there are infinitely many distinct tuples \bar{b}' , disjoint from \bar{a} , such that $p[\bar{b}'] \Vdash v \in \tau$ as well, where $p[\bar{b}']$ is the condition with domain \bar{a}, \bar{b}' defined on \bar{b}' as p is defined on \bar{b} . Note that $p[\bar{b}']$ extends $p \upharpoonright \bar{a}$.

Now for any r extending $p \upharpoonright \bar{a}$, there is some \bar{b}' such that $p[\bar{b}'] \Vdash \check{v} \in \tau$ and $p[\bar{b}']$ is compatible with r (take \bar{b}' disjoint from the domain of r). It follows that $p \upharpoonright \bar{a}$ forces $\check{v} \in \tau$. \square

9.5. The \mathbb{Z}^2 -jump of E_0 .

REFERENCES

- [BK96] Howard Becker and Alexander S. Kechris. *The descriptive set theory of Polish group actions*, volume 232 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1996.
- [CK11] Andrés Eduardo Caicedo and Richard Ketchersid. A trichotomy theorem in natural models of AD^+ . In *Set theory and its applications*, volume 533 of *Contemp. Math.*, pages 227–258. Amer. Math. Soc., Providence, RI, 2011.
- [CMRS23] F. Calderoni, D. Marker, L. Motto Ros, and A. Shani. An anti-classification result for groups acting freely on the line. *Advances in Mathematics*, 418, 2023.
- [CT19] Filippo Calderoni and Simon Thomas. The bi-embeddability relation for countable abelian groups. *Transactions of the American Mathematical Society*, 371(3):2237–2254, March 2019. Publisher Copyright: © 2018 American Mathematical Society.
- [Cum10] James Cummings. *Iterated Forcing and Elementary Embeddings*. Springer Netherlands, Dordrecht, 2010.
- [EGL24] Mahmood Etedadialabadi, Su Gao, and Feng Li. Extremely amenable automorphism groups of countable structures, 2024.
- [Fel71] Ulrich Felgner. *Models of ZF-set theory*. Lecture Notes in Mathematics, Vol. 223. Springer-Verlag, Berlin-New York, 1971.

- [FM95] Matthew Foreman and Menachem Magidor. Large cardinals and definable counterexamples to the continuum hypothesis. *Annals of Pure and Applied Logic*, 76(1):47–97, 1995.
- [For18] Matthew D. Foreman. What is ... a borel reduction? *Notices of the American Mathematical Society*, 2018.
- [FS89] Harvey Friedman and Lee Stanley. A borel reducibility theory for classes of countable structures. *J. Symbolic Logic*, 54(3):894–914, 09 1989.
- [Fuc73] László Fuchs. *Infinite Abelian Groups, Vol. II*. Pure and Applied Mathematics. Academic Press, New York, 1973.
- [Fuc15] László Fuchs. *Abelian Groups*. Springer Monographs in Mathematics. Springer, Cham, 2015.
- [Gao09] S. Gao. *Invariant descriptive set theory*. CRC Press, 2009.
- [Git08] Victoria Gitman. Scott’s problem for proper scott sets. *The Journal of Symbolic Logic*, 73(3):845–860, 2008.
- [Hal17] Lorenz J. Halbeisen. *Combinatorial Set Theory: With a Gentle Introduction to Forcing*. Springer Monographs in Mathematics. Springer, 2 edition, 2017.
- [Hjo95] Greg Hjorth. A dichotomy for the definable universe. *J. Symbolic Logic*, 60(4):1199–1207, 1995.
- [Hjo97] Greg Hjorth. Vaught’s conjecture on analytic sets. *Journal of the American Mathematical Society*, 14:125–143, 1997.
- [Hjo98] Greg Hjorth. An absoluteness principle for borel sets. *The Journal of Symbolic Logic*, 63(2):663–693, 1998.
- [Hjo99] Greg Hjorth. Orbit cardinals: on the effective cardinalities arising as quotient spaces of the form X/G where G acts on a Polish space X . *Israel J. Math.*, 111:221–261, 1999.
- [Hjo00] Greg Hjorth. *Classification and orbit equivalence relations*, volume 75 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2000.
- [Hjo01] Greg Hjorth. Vaught’s conjecture on analytic sets. *J. Amer. Math. Soc.*, 14(1):125–143, 2001.
- [HK95] Greg Hjorth and Alexander S. Kechris. Analytic equivalence relations and Ulm-type classifications. *J. Symbolic Logic*, 60(4):1273–1300, 1995.
- [HK01] Greg Hjorth and Alexander S. Kechris. Recent developments in the theory of Borel reducibility. volume 170, pages 21–52. 2001. Dedicated to the memory of Jerzy Łoś.
- [HKL90] L. A. Harrington, A. S. Kechris, and A. Louveau. A Glimm-Effros dichotomy for Borel equivalence relations. *J. Amer. Math. Soc.*, 3(4):903–928, 1990.
- [HKL98] Greg Hjorth, Alexander S. Kechris, and Alain Louveau. Borel equivalence relations induced by actions of the symmetric group. *Ann. Pure Appl. Logic*, 92(1):63–112, 1998.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.
- [Kan03] Akihiro Kanamori. *The Higher Infinite: Large Cardinals in Set Theory from Their Beginnings*. Perspectives in Logic. Springer, Berlin, Heidelberg, second edition, 2003.
- [Kan06] Akihiro Kanamori. Levy and set theory. *Ann. Pure Appl. Logic*, 140(1-3):233–252, 2006.
- [Kan08a] Akihiro Kanamori. Cohen and set theory. *Bull. Symbolic Logic*, 14(3):351–378, 2008.
- [Kan08b] Vladimir Kanovei. *Borel equivalence relations*, volume 44 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2008. Structure and classification.
- [Kec95] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [Kec99] Alexander S. Kechris. New directions in descriptive set theory. *The Bulletin of Symbolic Logic*, 5(2):161–174, 1999.
- [KMS16] JULIA KNIGHT, ANTONIO MONTALBÁN, and NOAH SCHWEBER. Computable structures in generic extensions. *The Journal of Symbolic Logic*, 81(3):814–832, 2016.
- [KN82a] Julia Knight and Mark Nadel. Expansions of models and turing degrees. *The Journal of Symbolic Logic*, 47(3):587–604, 1982.
- [KN82b] Julia Knight and Mark Nadel. Models of arithmetic and closed ideals. *The Journal of Symbolic Logic*, 47(4):833–840, 1982.

- [KS16] Itay Kaplan and Saharon Shelah. Forcing a countable structure to belong to the ground model. *Mathematical Logic Quarterly*, 62(6):530–546, 2016.
- [KTD12] Alexander S. Kechris and Robin D. Tucker-Drob. *The complexity of classification problems in ergodic theory*, page 265–300. London Mathematical Society Lecture Note Series. Cambridge University Press, 2012.
- [Kun80] Kenneth Kunen. *Set Theory: An Introduction to Independence Proofs*, volume 102 of *Studies in Logic and the Foundations of Mathematics*. North-Holland, Amsterdam, 1980.
- [Lar17] Paul B. Larson. Scott processes. In *Beyond first order model theory*, pages 23–76. CRC Press, Boca Raton, FL, 2017.
- [LS93] M. C. Laskowski and S. Shelah. On the existence of atomic models. *The Journal of Symbolic Logic*, 58(4):1189–1194, 1993.
- [LZ20] Paul B. Larson and Jindřich Zapletal. *Geometric set theory*, volume 248 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2020.
- [Mil17] Arnold W. Miller. *Descriptive Set Theory and Forcing: How to Prove Theorems about Borel Sets the Hard Way*. Lecture Notes in Logic. Cambridge University Press, 2017.
- [Mon73] G. P. Monro. Models of ZF with the same sets of sets of ordinals. *Fund. Math.*, 80(2):105–110, 1973.
- [MR21] Luca Motto Ros. Classification problems from the descriptive set theoretical perspective. arxiv: 2105.14865. 2021.
- [PS24] Aristotelis Panagiotopoulos and Assaf Shani. Strong ergodicity phenomena for bernoulli shifts of bounded algebraic dimension. *Annals of Pure and Applied Logic*, 175(5):103412, 2024.
- [Sam94] Ramez L. Sami. Polish group actions and the Vaught conjecture. *Trans. Amer. Math. Soc.*, 341(1):335–353, 1994.
- [Sha] Assaf Shani. The finite friedman-stanley jumps: generic dichotomies for borel homomorphisms. arxiv:2405.18360.
- [Sha21] Assaf Shani. Borel reducibility and symmetric models. *Trans. Amer. Math. Soc.*, 374(1):453–485, 2021.
- [Sha24a] Assaf Shani. Classifying invariants for E_1 : A tail of a generic real. *Notre Dame Journal of Formal Logic*, 65(3):333–356, August 2024.
- [Sha24b] Assaf Shani. Strong ergodicity around countable products of countable equivalence relations. *Israel Journal of Mathematics*, 263:871–908, 2024.
- [Ste84] Jacques Stern. On lusin’s restricted continuum problem. *Annals of Mathematics*, 120(1):7–37, 1984.
- [URL17] Douglas Ulrich, Richard Rast, and Michael C. Laskowski. Borel complexity and potential canonical Scott sentences. *Fund. Math.*, 239(2):101–147, 2017.
- [Wil80] George Wilmers. Minimally saturated models. In Leszek Pacholski, Jędrzej Wierzejewski, and Alec J. Wilkie, editors, *Model Theory of Algebra and Arithmetic*, pages 370–380, Berlin, Heidelberg, 1980. Springer Berlin Heidelberg.
- [Zap11] Jindřich Zapletal. Pinned equivalence relations. *Math. Res. Lett.*, 18(3):559–564, 2011.
- [Zip35] Leo Zippin. Countable torsion groups. *Annals of Mathematics*, 36(1):86–99, 1935.