

ON THE VANISHING OF TWISTED L -FUNCTIONS OF ELLIPTIC CURVES OVER FUNCTION FIELDS

ANTOINE COMEAU-LAPOINTE, CHANTAL DAVID, MATILDE LALIN AND WANLIN LI

ABSTRACT. We investigate in this paper the vanishing at $s = 1$ of the twisted L -functions of elliptic curves E defined over the rational function field $\mathbb{F}_q(t)$ (where \mathbb{F}_q is a finite field of q elements and characteristic ≥ 5) for twists by Dirichlet characters of prime order $\ell \geq 3$, from both a theoretical and numerical point of view. In the case of number fields, it is predicted that such vanishing is a very rare event, and our numerical data seems to indicate that this is also the case over function fields for non-constant curves. For constant curves, we adapt the techniques of [Li18, DL21] who proved vanishing at $s = 1/2$ for infinitely many Dirichlet L -functions over $\mathbb{F}_q(t)$ based on the existence of one, and we can prove that if there is one χ_0 such that $L(E, \chi_0, 1) = 0$, then there are infinitely many. Finally, we provide some examples which show that twisted L -functions of constant elliptic curves over $\mathbb{F}_q(t)$ behave differently than the general ones.

1. INTRODUCTION

Let E be an elliptic curve over \mathbb{Q} with L -function $L(E, s) = \sum_n a_n n^{-s}$, and χ be a Dirichlet character. Let $L(E, \chi, s) = \sum_n a_n \chi(n) n^{-s}$ be the twisted L -function. By the Birch and Swinnerton-Dyer conjecture, the vanishing of $L(E, \chi, s)$ at $s = 1$ is related to the growth of the rank of the Mordell-Weil group of E in the abelian extension of \mathbb{Q} associated to χ . Heuristics based on the distribution of modular symbols and random matrix theory ([DFK07, Conjecture 1.2], [MR]) have led to conjectures predicting that the vanishing of the twisted L -functions $L(E, \chi, s)$ at $s = 1$ is a very rare event as χ ranges over characters of prime order $\ell \geq 3$. For instance, it is predicted that there are only finitely many characters χ of order $\ell > 5$ such that $L(E, \chi, 1) = 0$. Mazur and Rubin rephrased this in terms of “Diophantine Stability”, and conjectured that if E is an elliptic curve over \mathbb{Q} and K/\mathbb{Q} is any real abelian extension such that K contains only finitely many subfields of degree 2, 3, or 5 over \mathbb{Q} , then the group of K -rational points $E(K)$ is finitely generated. They also proved that for each ℓ (under some hypotheses that can be shown to hold in certain contexts), there are infinitely many cyclic extensions K/\mathbb{Q} of order ℓ such that $E(K) = E(\mathbb{Q})$ (and then, assuming the Birch and Swinnerton-Dyer conjecture, such that the twisted L -functions $L(E, \chi, s)$ associated to the extensions K/\mathbb{Q} do not vanish) [MR18].

We remark that the case of vanishing of quadratic twists is very different from the higher order case $\ell \geq 3$ considered in this work, as the L -function of E twisted by a quadratic character of conductor D corresponds to the L -function of another elliptic curve E_D , and for half of the quadratic twists, $L(E, \chi_D, 1) = 1$. Goldfeld has conjectured that half of the

Date: May 26, 2022.

2020 Mathematics Subject Classification. Primary 11G05; Secondary 11G40, 14H25.

Key words and phrases. non-vanishing of L -functions; twisted L -functions of elliptic curves; function fields; elliptic curve rank in extensions.

twists E_D/\mathbb{Q} have rank 0, and half have rank 1 (asymptotically) [Gol74]). Furthermore, Gouvea and Mazur [GM91] have shown that the analytic rank of E_D is at least two for $\gg X^{1/2-\epsilon}$ of the quadratic discriminants $|D| \leq X$. The case of nonabelian extensions K/\mathbb{Q} of degree d with Galois group S_d is also different from the abelian extensions of order $\ell \geq 3$: in recent work, Lemke Oliver and Thorne [LOT21] showed that there are infinitely many such extensions where $\text{rank}(E(K)) > \text{rank}(E(\mathbb{Q}))$, for each $d \geq 2$, and Fornea [For19] has shown that for some curves E/\mathbb{Q} , the analytic rank of E increases for a positive proportion of the quintic fields with Galois group S_5 .

The vanishing (and non-vanishing) of twisted L -functions of elliptic curves is closely related to the one-level density, which is the study of low-lying zeroes, or the average analytic rank. This was studied over number fields and function fields, for quadratic and higher order twists. For quadratic twists, it is possible to prove results on the one-level density strong enough to deduce that a positive proportion of twists with even (respectively odd) analytic rank do not vanish (respectively vanish of order 1) at the central critical point [HB04, CL22]. The one-level density, or average rank, of higher order twists for elliptic curves L -functions was studied by [Cho] over number fields and [MS, CL22] over function fields. Quadratic twists of elliptic curve over function fields were also studied by [BFKRG20] who obtained results on the correlation of the analytic ranks of two twisted elliptic curves. The behavior of the algebraic rank of elliptic curves in cyclic extensions of \mathbb{Q} was investigated by Beneish, Kundu, and Ray [BKR].

We investigate in this article the vanishing at $s = 1$ of the twisted L -functions of elliptic curves E defined over the rational function field $\mathbb{F}_q(t)$,¹ for twists by Dirichlet characters of prime order $\ell \geq 3$, from both a theoretical and numerical point of view. It is natural to ask if the recent results of Li [Li18] and Donepudi and Li [DL21], who have found infinitely many instances of vanishing for L -functions of Dirichlet characters at $s = 1/2$, can be extended to L -functions of elliptic curves twisted by Dirichlet characters. We find that this is the case when E is a constant elliptic curve over $\mathbb{F}_q(t)$ ², and we can produce infinitely many cases of vanishing at the central critical point for characters of order ℓ provided we find one (Theorem 1.2). Then, the conjectures of [DFK07, MR21] do not hold in the special case of constant elliptic curves, and we present specific numerical examples in Section 6.

We also study non-constant elliptic curves over $\mathbb{F}_q(t)$ where q is a power of a prime $p \geq 5$, say $E : y^2 = x^3 + a(t)x + b(t)$, for some polynomials $a(t), b(t) \in \mathbb{F}_q[t]$. The L -function of $E/\mathbb{F}_q(t)$ is defined analogously as for E/\mathbb{Q} , by an infinite Euler product over the primes of $\mathbb{F}_q(t)$ (see (2.4)), but in this case, it follows from the work of Weil and Deligne that, after setting $u = q^{-s}$, $L(E, s) = \mathcal{L}(E, u)$, a polynomial in $\mathbb{Z}[u]$. Similarly, the twisted L -function $\mathcal{L}(E, \chi, u)$ is a polynomial in $\mathbb{Z}[\zeta_\ell][u]$, where χ is a Dirichlet character of order ℓ over $\mathbb{F}_q(t)$. More details and all relevant definitions are given in Section 2.

We present in Section 7 computational results for the vanishing of numerous twists of two base elliptic curves over $\mathbb{F}_q(t)$, the Legendre curve and a second curve, chosen to have good reduction at infinity. The data seems to indicate that the conjectures of [DFK07, MR21] also hold for non-constant elliptic curves over function fields, while presenting some unexpected

¹Throughout this article, we assume that \mathbb{F}_q is a finite field of q elements and characteristic ≥ 5 .

²Constant elliptic curves, i.e. an elliptic curve over \mathbb{F}_q considered as a curve over $\mathbb{F}_q(t)$, were studied by many authors because of their special properties. In particular, Milne showed that the Birch and Swinnerton-Dyer conjecture is true for constant elliptic curves [Mil68].

features. To our knowledge, this is the first data about the vanishing of L -functions of elliptic curves twisted by characters of order $\ell \geq 3$, over function fields. The case of quadratic twists of elliptic curves over function fields was considered by Baig and Hall [BH12] to test Goldfeld’s conjecture in that context, and our numerical computations are similar.

The case of a constant curve $E/\mathbb{F}_q(t)$ is defined by taking an elliptic curve E_0/\mathbb{F}_q and considering its base change to $\mathbb{F}_q(t)$, denoted by $E = E_0 \times_{\mathbb{F}_q} \mathbb{F}_q(t)$. In this case, the roots of $\mathcal{L}(E, \chi, u)$ can be described in terms of the roots of the L -functions $\mathcal{L}(E_0, u)$ and $\mathcal{L}(C_\chi, u)$, where the L -functions are associated to the elliptic curve E_0/\mathbb{F}_q and the ℓ -cyclic cover $C_\chi/\mathbb{P}_{\mathbb{F}_q}^1$ corresponding to the Dirichlet character χ respectively (see Section 3). This allows us to use a generalized version of the results of Li [Li18] and Donepudi–Li [DL21] about vanishing of the Dirichlet L -functions $\mathcal{L}(\chi, u)$ to obtain some vanishing for $\mathcal{L}(E, \chi, u)$ at $u = q^{-1}$. The argument of [Li18, DL21] has two distinct parts, first finding one character χ_0 such that $\mathcal{L}(\chi_0, u_0) = 0$ for some fixed u_0 , and then sieving to produce infinitely many such characters. The order of $q \bmod \ell$ is related to the presence/absence of ℓ -th roots of unity in $\mathbb{F}_q(t)$, which makes the study of the characters of order ℓ delicate, and the authors of [Li18, DL21] restrict to the Kummer case where $q \equiv 1 \pmod{\ell}$. As we need to treat all the cases (in particular, we often work over the finite field \mathbb{F}_p where p is prime), we generalize their sieving beyond the Kummer case. We also need to consider vanishing at any u_0 where $\mathcal{L}(E_0, u_0) = 0$, and not only $u_0 = q^{-1/2}$ as in their work.

We recall that an algebraic integer α is called a q -Weil integer if $|\alpha| = q^{1/2}$ under every complex embedding.

Theorem 1.1. *Let ℓ be a prime and q be a prime power coprime to ℓ . Let u_0 be a q -Weil integer. Suppose there exists a Dirichlet character χ_0 over $\mathbb{F}_q(t)$ of order ℓ and with conductor of degree d_0 such that $\mathcal{L}(\chi_0, u_0^{-1}) = 0$. Then, there are at least $\gg q^{2n/d_0}$ Dirichlet characters χ of order ℓ over $\mathbb{F}_q(t)$ with conductor of degree bounded by n such that $\mathcal{L}(\chi, u_0^{-1}) = 0$.*

We prove the above theorem in Section 4. The next result is then a direct consequence of Theorem 1.1, using the properties of constant elliptic curves discussed in Section 3.

Theorem 1.2. *Let E_0 be an elliptic curve over \mathbb{F}_q , and let $E = E_0 \times_{\mathbb{F}_q} \mathbb{F}_q(t)$. Suppose there exists a Dirichlet character χ_0 over $\mathbb{F}_q(t)$ of order ℓ and with conductor of degree d_0 such that $\mathcal{L}(E, \chi_0, q^{-1}) = 0$. Then, there are at least $\gg q^{2n/d_0}$ Dirichlet characters χ of order ℓ over $\mathbb{F}_q(t)$ with conductor of degree bounded by n such that $\mathcal{L}(E, \chi, q^{-1}) = 0$.*

Then, to guarantee that a constant elliptic curve $E/\mathbb{F}_q(t)$ has infinitely many twists of order ℓ such that $L(E, \chi, u)$ vanishes at q^{-1} , it suffices to find one. Using the results of Section 3, this can be rephrased in terms of finding curves C/\mathbb{F}_q which are ℓ -cyclic covers of $\mathbb{P}_{\mathbb{F}_q}^1$ and such that $\mathcal{L}(E_0, u)$ divides $\mathcal{L}(C, u)$, and we investigate this question numerically in Section 6.

In Section 6, we present various isogeny classes of elliptic curves E_0 over different prime fields such that $\mathcal{L}(E, \chi, q^{-1}) = 0$ for characters χ of prime order $\ell = 3, 5, 7, 11$. One observation from the data is the existence of supersingular curves defined over prime fields \mathbb{F}_p which admit a degree ℓ cyclic map to \mathbb{P}^1 ramifying at 4 points where $p \equiv -1 \pmod{\ell}$. The existence of such curves does not follow from previous results on the topic and one may hope to prove this statement following the strong evidence presented in Table 1.

It is natural to ask if the same dichotomy (no instances of vanishing or infinitely many cases of vanishing) also holds for non-constant elliptic curves over $\mathbb{F}_q(t)$, but there is no

reason to believe it would be the case. The ideas leading to the proof of Proposition 1.2 for constant curves do not apply to the general case, as the change of variable trick used to produce infinitely many extensions where E acquires points would send points on E to points on a different elliptic curve when E is not constant. However, there are results of that type for an elliptic curve E over \mathbb{Q} due to Fearnley, Kisilevsky, and Kuwata [FKK12], where the authors prove that if there is one cyclic cubic field K such that $E(K)$ is infinite, then there are infinitely many, and there are always infinitely many such K when $E(\mathbb{Q})$ contains at least 6 points. On the non-vanishing side, Brubaker, Bucur, Chinta, Frechette and Hoffstein [BBC⁺04] use the method of multiple Dirichlet series to prove that if there exists a single non-vanishing order ℓ twist of an L -function associated to a cuspidal automorphic representation of $GL(2, \mathbb{A}_K)$, then there are infinitely many.

The structure of this article is as follows: we define in Section 2 the L -functions attached to Dirichlet characters and elliptic curves over $\mathbb{F}_q(t)$, and we recall their properties. We discuss in Section 3 the case of the L -functions of constant elliptic curves. We describe the ℓ -cyclic covers of $\mathbb{P}_{\mathbb{F}_q}^1$ and their characters in Section 4, for all cases (not only the Kummer case $q \equiv 1 \pmod{\ell}$) using the work of Bary-Soroker and Meisner [BSM19], and we then generalize the sieves of [Li18, DL21] to those general ℓ -cyclic covers. We then use those results to prove Theorems 1.1 and 1.2. Finally, we describe our computations in Section 5, and we present our numerical data in Sections 6 and 7.

Acknowledgments. The authors would like to thank Patrick Meisner for helpful discussions, and the anonymous referees for helpful comments that greatly improved the exposition of this paper. This work is supported by the Natural Sciences and Engineering Research Council of Canada (NSERC Discovery Grant 155635-2019 to CD, 335412-2013 to ML), by the Fonds de recherche du Québec - Nature et technologies (Projet de recherche en équipe 300951 to CD and ML), and by the Centre de recherches mathématiques and the Institut des sciences mathématiques (CRM-ISM postdoctoral fellowship to WL). Some of the computations were checked using the computational software MAGMA.

2. DIRICHLET CHARACTERS, ELLIPTIC CURVES AND L -FUNCTIONS OVER $\mathbb{F}_q(t)$

2.1. Dirichlet characters. We review here the theory of Dirichlet characters of order ℓ over $\mathbb{F}_q(t)$, and their L -functions. We refer the reader to [DFL] and [BSM19] for more details.

Let n_q be the multiplicative order of q modulo ℓ . We say that we are in the Kummer case if $n_q = 1$ and in the non-Kummer case otherwise. We also say that a monic irreducible polynomial $P \in \mathbb{F}_q[t]$ is n_q -divisible if $n_q \mid \deg P$.

We first consider the Kummer case and we fix once and for all an isomorphism Ω from the ℓ -th roots of unity in \mathbb{F}_q^* to the ℓ -th roots of unity in \mathbb{C}^* . We first define the ℓ -th order residue symbol χ_P , for P an irreducible monic polynomial in $\mathbb{F}_q[t]$. For any $a \in \mathbb{F}_q[t]$, if $P \mid a$, then $\chi_P(a) = 0$, and otherwise $\chi_P(a) = \alpha$, where α is the unique ℓ -th root of unity in \mathbb{C}^* such that

$$a^{\frac{q^{\deg(P)} - 1}{\ell}} \equiv \Omega^{-1}(\alpha) \pmod{P}.$$

We extend the definition multiplicatively to any monic polynomial $F \in \mathbb{F}_q[t]$ by defining for $F = P_1^{e_1} \dots P_s^{e_s}$, with distinct primes P_i ,

$$\chi_F = \chi_{P_1}^{e_1} \cdots \chi_{P_s}^{e_s}.$$

Then, χ_F is a character of order ℓ modulo $P_1 \dots P_s$, and the primitive characters of conductor $P_1 \dots P_s$ are given by taking all choices $1 \leq e_i \leq \ell - 1$, i.e. the conductors of the primitive characters are the square-free monic polynomials $F \in \mathbb{F}_q[t]$, and for each such conductor, there are $(\ell - 1)^{\omega(F)}$ characters, where $\omega(F)$ is the number of primes dividing F .

For the non-Kummer case, the ℓ -th residue symbols χ_P are defined similarly, but only for the n_q -divisible primes P , since we must have $\ell \mid q^{\deg P} - 1$. The conductors of such characters are then square-free polynomials F supported on n_q -divisible primes and for each such F , there are $(\ell - 1)^{\omega(F)}$ primitive characters of order ℓ , as above.

In both cases, we can write each character of conductor F as

$$(2.1) \quad \chi_F = \chi_{F_1} \chi_{F_2}^2 \dots \chi_{F_{\ell-1}}^{\ell-1}$$

corresponding to a decomposition $F = F_1 \dots F_{\ell}$ where the F_i 's are square-free and coprime.

Let χ be a Dirichlet character, and let $\mathcal{L}(\chi, u)$ be the Dirichlet L -function defined by

$$\mathcal{L}(\chi, u) = \prod_P (1 - \chi(P)u^{\deg P})^{-1},$$

where the product includes the prime at infinity, and we give the value of $\chi(P_{\infty})$ in Lemma 2.1. If the restriction of χ to \mathbb{F}_q is trivial, we say that χ is even, otherwise, we say that χ is odd. We define δ_{χ} by

$$(2.2) \quad \delta_{\chi} := \begin{cases} 0 & \text{when } \chi \text{ is even,} \\ 1 & \text{when } \chi \text{ is odd.} \end{cases}$$

For χ a primitive character of conductor F , it follows from the work of Weil that $\mathcal{L}(\chi, u)$ is a polynomial of degree $\deg(F) - 2 + \delta_{\chi}$ and satisfies the functional equation

$$(2.3) \quad \mathcal{L}(\chi, u) = \omega_{\chi} (\sqrt{q}u)^{\deg(F)-2+\delta_{\chi}} \mathcal{L}(\bar{\chi}, 1/(qu)).$$

The sign of the functional equation is

$$\omega_{\chi} = \begin{cases} \frac{G(\chi)}{|G(\chi)|} & \text{when } \chi \text{ is even,} \\ \frac{\sqrt{q}}{\tau(\chi)} \frac{G(\chi)}{|G(\chi)|} & \text{when } \chi \text{ is odd,} \end{cases}$$

where if χ odd,

$$\tau(\chi) = \sum_{a \in \mathbb{F}_q^*} \chi(a) e^{2\pi i \text{tr}_{\mathbb{F}_q/\mathbb{F}_p}(a)/p},$$

and for any χ , $G(\chi)$ is the Gauss sum

$$G(\chi) = \sum_{a \bmod F} \chi(a) e_q \left(\frac{a}{F} \right).$$

Here e_q is the exponential defined by Hayes [Hay66] for any $b \in \mathbb{F}_q((1/T))$:

$$e_q(b) = e^{\frac{2\pi i \text{tr}_{\mathbb{F}_q/\mathbb{F}_p}(b_1)}{p}},$$

where b_1 is the coefficient of $1/T$ in the Laurent expansion of b . We refer the reader to [DFL] for a proof of those results.

Lemma 2.1. *Let χ be a Dirichlet character of order ℓ defined over $\mathbb{F}_q(t)$ with conductor F , where F is a monic squarefree polynomial in $\mathbb{F}_q[t]$.*

If $q \not\equiv 1 \pmod{\ell}$, then χ does not ramify at infinity, $\chi(P_\infty) = 1$, and χ is even.

If $q \equiv 1 \pmod{\ell}$, let $\chi = \chi_{F_1} \chi_{F_2}^2 \cdots \chi_{F_{\ell-1}}^{\ell-1}$ as in (2.1). Then,

χ ramifies at $P_\infty \iff \ell \nmid \deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1}) \iff \chi$ is odd, and

$$\chi(P_\infty) = \begin{cases} 1 & \ell \mid \deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1}), \\ 0 & \ell \nmid \deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1}). \end{cases}$$

Proof. We first discuss when the characters are odd and even. Let P be an n_q -divisible prime. As in the Kummer case, we can fix an isomorphism Ω from the ℓ -th roots of unity in $\mathbb{F}_{q^{n_q}}$ to the ℓ -th roots of unity in \mathbb{C} . We remark that for $\alpha \in \mathbb{F}_q^*$,

$$\chi_P(\alpha) = \Omega \left(\alpha^{\frac{q^{\deg(P)} - 1}{\ell}} \right) = \Omega \left(\alpha^{\frac{\deg(P)(q^{n_q} - 1)}{n_q \ell}} \right).$$

Indeed, writing $\deg(P) = n_q k$, we have

$$\frac{q^{\deg(P)} - 1}{\ell} = \frac{q^{n_q k} - 1}{\ell} = \frac{q^{n_q} - 1}{\ell} (1 + q^{n_q} + \cdots + q^{n_q(k-1)})$$

and we use the fact that $1 + q^{n_q} + \cdots + q^{n_q(k-1)} \equiv k \pmod{\ell}$.

First assume that $q \equiv 1 \pmod{\ell}$. Then by multiplicativity,

$$\chi_F(\alpha) = \Omega \left(\alpha^{\frac{\deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1})(q-1)}{\ell}} \right),$$

and χ is trivial over \mathbb{F}_q iff $\ell \mid \deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1})$.

Now suppose that $n_q > 1$. We have as before

$$\chi(\alpha) = \Omega \left(\alpha^{\frac{\deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1})(q^{n_q} - 1)}{n_q \ell}} \right),$$

and since $n_q > 1$, we have that $\ell \nmid (q - 1)$. In particular $(\ell, q - 1) = 1$ since ℓ is prime. Now we have that both $\ell \mid (q^{n_q} - 1)$ and $(q - 1) \mid (q^{n_q} - 1)$. It follows that $(q - 1) \mid \frac{q^{n_q} - 1}{\ell}$. Taking $\alpha \in \mathbb{F}_q$, we have

$$\chi(\alpha) = \Omega \left(\alpha^{\frac{\deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1})(q^{n_q} - 1)}{n_q \ell}} \right) = 1,$$

and therefore χ is an even character.

The statement that P_∞ does not ramify for the non-Kummer case follows from the fact that the cyclic field extension can only ramify at primes of degree divisible by $n_q > 1$ and P_∞ is a prime of degree 1. In the Kummer case, the character χ_F is associated with the cyclic cover $y^\ell = F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1}$, and there is ramification at ∞ iff $\ell \nmid \deg(F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1})$, and $\chi(P_\infty) = 0$ in this case. If χ does not ramify at P_∞ , then $\chi(P_\infty) = 1$ since we are only considering the case in which $F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1}$ is monic. \square

2.2. **L -functions of elliptic curves over $\mathbb{F}_q(t)$.** Let E be an elliptic curve over $\mathbb{F}_q(t)$. Let P be a prime of $\mathbb{F}_q(t)$, i.e $P = P(t) \in \mathbb{F}_q[t]$ is an irreducible polynomial or $P = P_\infty$, the prime at infinity. If P is a prime of good reduction, then the reduction of E (which we also denote by E) is an elliptic curve over the finite field $\mathbb{F}_P = \mathbb{F}_q[t]/(P) \cong \mathbb{F}_{q^{\deg P}}$ (where $\mathbb{F}_\infty \cong \mathbb{F}_q$ since the prime at infinity has degree 1), and

$$\#E(\mathbb{F}_P) = q^{\deg P} + 1 - a_P, \quad a_P = \alpha_P + \bar{\alpha}_P, \quad |\alpha_P| = \sqrt{q^{\deg P}}.$$

Let

$$\mathcal{L}_P(E, u) := 1 - a_P u + q^{\deg P} u^2 = (1 - \alpha_P u)(1 - \bar{\alpha}_P u)$$

be the L -function of E/\mathbb{F}_P . Let N_E be the conductor of E , which is the product of the primes of bad reduction with the appropriate powers.³ Let M_E (respectively A_E) be the product of the multiplicative (respectively additive) primes of E . Then $N_E = M_E A_E^2$.

The L -function of E is defined by

$$(2.4) \quad \mathcal{L}(E, u) := \prod_{P \nmid N_E} \mathcal{L}_P(E, u^{\deg P})^{-1} \prod_{P \mid N_E} \mathcal{L}_P(E, u^{\deg P})^{-1},$$

where for primes dividing N_E , we define

$$\mathcal{L}_P(E, u^{\deg P}) = (1 - a_P u^{\deg P})$$

and $a_P = 0, 1, -1$ depending on the type of bad reduction (additive, split multiplicative, and non-split multiplicative respectively). It is proven by Weil [Kat02, BH12] that $\mathcal{L}(E, u)$ is a polynomial of degree⁴ $\deg N_E - 4$ for any non-constant elliptic curve defined over the rational function field $\mathbb{F}_q(t)$ and it satisfies the functional equation

$$(2.5) \quad \mathcal{L}(E, u) = \omega_E (qu)^{\deg(N_E)-4} \mathcal{L}(E, 1/(q^2 u)),$$

where $\omega_E = \pm 1$ is the sign of the functional equation. We refer the reader to [Bru92, Appendix] and [BH12] for more details.

Let χ be a Dirichlet character of order ℓ and conductor F , and suppose that $(F, N_E) = 1$. If χ is odd, we also assume that E has good reduction at ∞ (since the prime at infinity is not included in the conductor of the Dirichlet character, we need this additional condition to ensure that the places where χ ramifies and the places of bad reduction for E are disjoint). The L -function of E twisted by χ is defined by

$$(2.6) \quad \begin{aligned} \mathcal{L}(E, \chi, u) &:= \prod_{P \nmid N_E} (1 - \chi(P) \alpha_P u^{\deg(P)})^{-1} (1 - \chi(P) \bar{\alpha}_P u^{\deg(P)})^{-1} \\ &\times \prod_{P \mid N_E} (1 - \chi(P) a_P u^{\deg(P)})^{-1}. \end{aligned}$$

Let K be the cyclic field extension of degree ℓ over $\mathbb{F}_q(t)$ corresponding to χ . Then,

$$(2.7) \quad \mathcal{L}(E/K, u) = \mathcal{L}(E, u) \prod_{i=1}^{\ell-1} \mathcal{L}(E, \chi^i, u).$$

³We emphasize that we include the prime at infinity in the conductor of the elliptic curve (if the curve has bad reduction at infinity of course). Our conductor is an effective divisor, written multiplicatively.

⁴The formula for the degree of $\mathcal{L}(E, u)$ implies in particular that there are no non-constant elliptic curves over $\mathbb{F}_q(t)$ with conductor of degree smaller than 4, which can be thought of as the analogue to the fact that there are no elliptic curves over \mathbb{Q} with conductor smaller than 11.

It follows from the Riemann Hypothesis that

$$\mathcal{L}(E/K, u) = \prod_{j=1}^B (1 - qe^{i\theta_j} u).$$

Since there is no common ramification between E and χ , (2.7) and Theorem 2.2 (stated and proven below) imply that $B = \ell(\deg N_E - 4) + 2(\ell - 1)(\deg F + \delta_\chi)$.

It is well-known that $\mathcal{L}(E, \chi, u)$ satisfies a functional equation from the work of Weil [Wei71]. The explicit formula for the sign of the functional equation is contained in [Wei71] in a very general context, but we need a precise formula for the numerical computations, so we deduce it below from the work of Tan and Rockmore [Tan93, TR92].

Theorem 2.2. *Let ℓ be a prime, χ a primitive Dirichlet character of conductor F and order ℓ , and let E be a non-constant elliptic curve with conductor N_E such that $(N_E, F) = 1$. If $P_\infty \mid N_E$, we also assume that χ is even. The L -function $\mathcal{L}(E, \chi, u)$ is a polynomial of degree*

$$\mathfrak{n} := \deg N_E + 2 \deg F - 4 + 2\delta_\chi,$$

where δ_χ is given by (2.2). Each $\mathcal{L}(E, \chi, u)$ satisfies the functional equation

$$(2.8) \quad \mathcal{L}(E, \chi, u) = \omega_{E \otimes \chi} (qu)^{\mathfrak{n}} \mathcal{L}(E, \bar{\chi}, 1/(q^2 u)),$$

where $\omega_{E \otimes \chi}$ is the sign of the functional equation for $\mathcal{L}(E, \chi, u)$, given by

$$\omega_{E \otimes \chi} = \omega_\chi^2 \omega_E \chi(N_E).$$

Proof. The sign of the functional equation (and the functional equation itself) can be deduced from the modularity of elliptic curves over function fields. We follow [Tan93, TR92] who use modular symbols over function fields. They consider different normalizations, so we explain here how to adjust their work to get the result that we need. Let $K = \mathbb{F}_q(t)$. For any place v , let \mathcal{O}_v be the associated ring of integers. If $N = \sum_v N_v$ is an effective divisor over K , let

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \left(\begin{pmatrix} a_v & b_v \\ c_v & d_v \end{pmatrix}_v \right) \in \prod_v \mathrm{GL}_2(\mathcal{O}_v) : c \equiv 0 \pmod{N} \right\}.$$

Let \mathbb{A}_K be the ring of adèles over K . Then \mathbb{A}_K^* embeds in $\mathrm{GL}_2(\mathbb{A}_K)$ as diagonal matrices. Also $\mathrm{GL}_2(K)$ embeds in $\mathrm{GL}_2(\mathbb{A}_K)$ by the diagonal map.

A \mathbb{C} -valued function on $\mathrm{GL}_2(\mathbb{A}_K)$ is called a modular function of level N if it satisfies that $f(\gamma\tau\kappa) = f(\tau)$ for all $\tau \in \mathrm{GL}_2(\mathbb{A}_K)$, $\gamma \in \mathrm{GL}_2(K)$, and $\kappa \in \mathbb{A}_K^* \cdot \Gamma_0(N)$. It is a fundamental result that if E is a non-constant elliptic curve over K , then there is a normalized cuspidal modular function f of level N_E such that the L -function of E is the L -function of f . This also holds for the twisted L -functions. To make that statement precise, and use it to get the functional equation, we will follow the notation of [Tan93, TR92], where the L -functions are normalized differently (and we will go back to our L -function at the end). Let f be the normalized cuspidal modular function corresponding to E , χ a Dirichlet character of conductor coprime to N_E and we define as [Tan93, (1.10)]

$$L_f(\chi, s) = \sum_M \frac{c_f(M)\chi(M)}{|M|^{s-1}},$$

where M runs through all effective divisors and the $c_f(M)$ are the normalized coefficients obtained from the Fourier expansion of f . This is also true when χ is a quasi-character, which

for our purposes is the product of a Dirichlet character and a map χ_s given by $\chi_s(M) = |M|^{-s}$.

We now use the modular symbols $\Theta_{f,D}$ to get the functional equation. The modular symbols $\Theta_{f,D}$ are elements of the group ring $R[W_D]$, where $W_D = K^* \backslash \mathbb{A}_K^* / U_D$ is the Weil group of a divisor D of K , and R is a ring containing all the Fourier coefficients of f . We refer to [Tan93] for all the relevant definitions. The modular symbols are used to interpolate special values of the twisted L -functions, and we have [Tan93, Proposition 2],

$$(2.9) \quad L_f(\chi, 1) = \tau_\chi^{-1} \chi(\Theta_{f,D}),$$

where τ_χ is a Gauss sum. Using quasi-characters, we also have

$$(2.10) \quad L_f(\chi, s) = L_f(\chi\chi_{s-1}, 1) = \tau_{\chi\chi_{s-1}}^{-1} (\chi\chi_{s-1})(\Theta_{f,D}).$$

Using the Atkin–Lehner involution w_{N_E} , we have when $(D, N_E) = 1$ (including at ∞) [Tan93, Proposition 3]

$$(2.11) \quad \Theta_{f,D} = \Theta_{w_{N_E}(f),D}^t N_E,$$

where t is the involution on $R[W_D]$ sending $\sum_{w \in W_D} a_w w$ to $\sum_{w \in W_D} a_w w^{-1}$.

Applying a quasi-character χ to $\Theta = \sum_{w \in W_D} a_w w$ results in $\chi(\Theta) = \sum_{w \in W_D} a_w \chi(w)$, while applying χ together with the involution t results in $\chi(\Theta^t) = \sum_{w \in W_D} a_w \chi^{-1}(w) = \chi^{-1}(\Theta)$.

We apply $\chi\chi_{s-1}$ to (2.11), and we combine it with (2.10) to get

$$\begin{aligned} L_f(\chi, s) &= \tau_{\chi\chi_{s-1}}^{-1} (\chi\chi_{s-1})(\Theta_{f,D}) \\ &= \tau_{\chi\chi_{s-1}}^{-1} (\chi\chi_{s-1})(\Theta_{w_{N_E}(f),D}^t) \chi(N_E) |N_E|^{-(s-1)} \\ &= \frac{\tau_{\chi^{-1}\chi_{1-s}}}{\tau_{\chi\chi_{s-1}}} L_{w_{N_E}(f)}(\chi^{-1}\chi_{1-s}, 1) \chi(N_E) |N_E|^{-(s-1)} \\ &= \frac{\tau_{\chi^{-1}\chi_{1-s}}}{\tau_{\chi\chi_{s-1}}} L_{w_{N_E}(f)}(\chi^{-1}, 2-s) \chi(N_E) |N_E|^{-(s-1)}. \end{aligned}$$

The third line above follows from using (2.9) with f replaced by $w_{N_E}(f)$ and $\chi\chi_{s-1}$ replaced by $(\chi\chi_{s-1})^{-1}$, together with the observation that the involution t has the effect of inverting the character. Using the fact that f is an eigenvector for the self-dual Atkin–Lehner operator, we have $w_{N_E}(f) = \omega_E f$, where $\omega_E = \pm 1$ is the sign of the functional equation (2.5), and then $L_{w_{N_E}(f)}(\chi^{-1}, 2-s) = \omega_E L_f(\chi^{-1}, 2-s)$.

To compute the Gauss sums associated with the quasi-characters, we use [TR92, (2.2.3)]

$$\tau_{\chi\chi_s} = q^{s(\deg D - 2)} \tau_\chi,$$

where τ_χ is the Gauss sum of the Dirichlet character χ of conductor D . Replacing above, this gives

$$(2.12) \quad \tau_\chi L_f(\chi, s) = \omega_E \tau_{\chi^{-1}} \chi(N_E) q^{(1-s)(\deg(N_E) + 2 \deg(D) - 4)} L_f(\chi^{-1}, 2-s),$$

where [Tan93, (3.4)] is a particular case (for $s = 1$). The L -function of the elliptic curve is given by

$$L(E, \chi, s) = \sum_M \frac{c_f(M) |M| \chi(M)}{|M|^s} = \mathcal{L}(E, \chi, u)$$

for $u = q^{-s}$. The functional equation can be obtained by noticing that $L_f(\chi, s) = L(E, \chi, s)$, and replacing in (2.12). This leads to

$$\tau_\chi L(E, \chi, s) = \omega_E \tau_{\chi^{-1}} \chi(N_E) q^{(1-s)(\deg(N_E)+2\deg(D)-4)} L(E, \chi^{-1}, 2-s).$$

Using $u = q^{-s}$, we finally get

$$(2.13) \quad \mathcal{L}(E, \chi, u) = \omega_{E \otimes \chi} (qu)^{(\deg(N_E)+2\deg(D)-4)} \mathcal{L}(E, \chi^{-1}, 1/(q^2u)),$$

where

$$\omega_{E \otimes \chi} = \left(\frac{\overline{\tau_\chi}}{|D|^{1/2}} \right)^2 \omega_E \chi(N_E).$$

In order to get exactly the statement of the theorem, we need to take into account the difference of notation between [Tan93] and this paper. When χ is odd and there is ramification at ∞ , the conductor D of (2.13) is $P_\infty D'$, where $D' \in \mathbb{F}_q[t]$, and so D' is the definition of the conductor in this paper. Adjusting the formula to make it compatible with our notation, we get for all cases

$$\mathcal{L}(E, \chi, u) = \omega_{E \otimes \chi} (qu)^{(\deg(N_E)+2\deg(D)-4+2\delta_\chi)} \mathcal{L}(E, \chi^{-1}, q^2u^{-1}),$$

which is the functional equation (2.8). Finally, we remark that $\frac{\overline{\tau_\chi}}{|D|^{1/2}}$ is by definition the sign of the functional equation of $\mathcal{L}(\chi, u)$, since it is the product of the same local Gauss sums because $(D, N_E) = 1$, and we have $\omega_{E \otimes \chi} = \omega_\chi^2 \omega_E \chi(N_E)$. □

Remark 2.3. *When E is a constant elliptic curve, we prove in the next section that $\mathcal{L}(E, \chi, u)$ satisfies the same functional equation with $\mathfrak{n} = 2 \deg F - 4 + 2\delta_\chi$ and $\omega_{E \otimes \chi} = \omega_\chi^2$.*

3. L-FUNCTIONS OF CONSTANT ELLIPTIC CURVES OVER $\mathbb{F}_q(t)$

Let C be a ℓ -cyclic cover of $\mathbb{P}_{\mathbb{F}_q}^1$ of genus g , and let $K = \mathbb{F}_q(C)$ be the corresponding extension of $\mathbb{F}_q(t)$. The zeta function of C can be expressed as

$$(3.1) \quad \mathcal{Z}(C, u) = \mathcal{Z}(u) \mathcal{L}(C, u) = \frac{\prod_{j=1}^{2g} (1 - \beta_j u)}{(1-u)(1-qu)},$$

where $|\beta_j| = q^{1/2}$ for $1 \leq j \leq 2g$. We also have

$$\mathcal{L}(C, u) = \prod_{i=1}^{\ell-1} \mathcal{L}(\chi^i, u),$$

where the χ^i are the characters of order ℓ associated to the extension $K/\mathbb{F}_q(t)$.

Let E_0 be an elliptic curve over \mathbb{F}_q with L -function

$$\mathcal{L}(E_0, u) = (1 - \alpha_1 u)(1 - \alpha_2 u).$$

Theorem 3.1. Let $E = E_0 \times_{\mathbb{F}_q} \mathbb{F}_q(t)$, and let C, K and α_1, α_2 , and the β_j 's be as above. Then,

$$(3.2) \quad \begin{aligned} \mathcal{L}(E/K, u) &= \mathcal{Z}(C, \alpha_1 u) \mathcal{Z}(C, \alpha_2 u) \\ &= \frac{\prod_{\substack{1 \leq i \leq 2 \\ 1 \leq j \leq 2g}} (1 - \alpha_i \beta_j u)}{\prod_{1 \leq i \leq 2} (1 - \alpha_i u)(1 - \alpha_i q u)}. \end{aligned}$$

Moreover, $\mathcal{L}(E, \chi, u) = \mathcal{L}(\chi, \alpha_1 u) \mathcal{L}(\chi, \alpha_2 u)$, and writing

$$\mathcal{L}(\chi, u) = \prod_{1 \leq j \leq 2g/(\ell-1)} (1 - \gamma_j u),$$

then

$$\mathcal{L}(E, \chi, u) = \prod_{\substack{1 \leq i \leq 2 \\ 1 \leq j \leq 2g/(\ell-1)}} (1 - \alpha_i \gamma_j u).$$

Proof. We refer the reader to [Mil68, Section 3] and to [Oes90, Section 3.2] for the general proof. To illustrate the ideas, we prove (3.2) when $K = \mathbb{F}_q(t)$. Since $\#E_0(\mathbb{F}_{q^n}) = q^n + 1 - \alpha_1^n - \alpha_2^n$, if P is a prime, then

$$\#E(\mathbb{F}_P) = \#E_0(\mathbb{F}_P) = q^{\deg(P)} + 1 - \alpha_1^{\deg(P)} - \alpha_2^{\deg(P)}.$$

Since all the primes are of good reduction, we have

$$\begin{aligned} \mathcal{L}(E/\mathbb{F}_q(t), u) &= \mathcal{L}(E, u) = \prod_P (1 - (\alpha_1^{\deg(P)} + \alpha_2^{\deg(P)})u^{\deg(P)} + q^{\deg(P)}u^{2\deg(P)})^{-1} \\ &= \prod_P (1 - \alpha_1^{\deg(P)}u^{\deg(P)})^{-1} (1 - \alpha_2^{\deg(P)}u^{\deg(P)})^{-1} \\ &= \frac{1}{(1 - \alpha_1 u)(1 - q\alpha_1 u)(1 - \alpha_2 u)(1 - q\alpha_2 u)} \\ &= \mathcal{Z}(\alpha_1 u) \mathcal{Z}(\alpha_2 u). \quad \square \end{aligned}$$

Remark 3.2. From the above result, it is easy to get the functional equation for $\mathcal{L}(E, \chi, u)$ when E is a constant curve, using the functional equation of $\mathcal{L}(\chi, u)$ given by (2.3). Let $m = \deg_u \mathcal{L}(\chi, u) = 2g/(\ell - 1)$. In the notation of Section 2, we have $m = 2g/(\ell - 1) = \deg F - 2 + \delta_\chi$, and

$$\begin{aligned} \mathcal{L}(E, \chi, u) &= \mathcal{L}(\chi, \alpha_1 u) \mathcal{L}(\chi, \alpha_2 u) = \omega_\chi(\sqrt{q}\alpha_1 u)^m \mathcal{L}(\bar{\chi}, 1/q\alpha_1 u) \omega_\chi(\sqrt{q}\alpha_2 u)^m \mathcal{L}(\bar{\chi}, 1/q\alpha_2 u) \\ &= \omega_\chi^2(q^2 u^2)^m \mathcal{L}(\bar{\chi}, \alpha_2/(q^2 u)) \mathcal{L}(\bar{\chi}, \alpha_1/(q^2 u)) \\ &= \omega_\chi^2(qu)^{2m} \mathcal{L}(E, \bar{\chi}, 1/(q^2 u)) = \omega_\chi^2(qu)^{2\deg F - 4 + 2\delta_\chi} \mathcal{L}(E, \bar{\chi}, 1/(q^2 u)) \end{aligned}$$

Corollary 3.3. Let $E = E_0 \times_{\mathbb{F}_q} \mathbb{F}_q(t)$, and let χ be a Dirichlet character over $\mathbb{F}_q(t)$ with associated curve C and function field $K = \mathbb{F}_q(C)$ respectively. Then, $\mathcal{L}(E/K, q^{-1}) = 0$ if and only if $\mathcal{L}(C, \alpha_1^{-1}) = \mathcal{L}(C, \alpha_2^{-1}) = 0$,

Proof. From equation (3.2) in Theorem 3.1, $\mathcal{L}(E/K, q^{-1}) = 0$ if and only if there is one $\beta_j = q/\alpha_1 = \alpha_2$ or $\beta_j = q/\alpha_2 = \alpha_1$, where the β_j 's are given by (3.1), and both α_1^{-1} and α_2^{-1} are roots of $\mathcal{L}(C, u)$, because of the functional equation of $\mathcal{L}(C, u)$. \square

4. CYCLIC EXTENSIONS OF DEGREE ℓ OVER $\mathbb{F}_q(t)$

We prove in this section the following result which extends the result of [DL21] to general q and ℓ (removing the restrictions $q \equiv 1 \pmod{\ell}$ and $y^\ell = F(t)$ with $\ell \mid \deg F$).

Proposition 4.1. *Let ℓ be an odd prime. Fix an ℓ -cyclic cover C_0 over $\mathbb{P}_{\mathbb{F}_q(t)}^1$ with conductor of degree d_0 . Then there are at least $\gg q^{2n/d_0}$ ℓ -cyclic covers C with conductor of degree bounded by n admitting a non-constant map from C to C_0 .*

The proof of this result is fairly long and will require several intermediate steps.

The affine equations for ℓ -cyclic covers are well-known in the Kummer case $q \equiv 1 \pmod{\ell}$, which is the case treated in [DL21]. In this case, C has an affine equation $y^\ell = F_1 F_2^2 \cdots F_{\ell-1}^{\ell-1}$, where $F_i \in \mathbb{F}_q[t]$ are square-free and pairwise co-prime of degree d_i . The conductor of the ℓ -cyclic cover is $F_1 \cdots F_{\ell-1}$ and by the Riemann–Hurwitz formula, the genus is $\frac{\ell-1}{2}(d_1 + \cdots + d_{\ell-1} - 2)$ if $\ell \mid (d_1 + 2d_2 + \cdots + (\ell-1)d_{\ell-1})$ and $\frac{\ell-1}{2}(d_1 + \cdots + d_{\ell-1} - 1)$ otherwise. In this later case, there is ramification at infinity since $\ell \nmid (d_1 + 2d_2 + \cdots + (\ell-1)d_{\ell-1})$ by Lemma 2.1.

To treat the general case and prove Proposition 4.1, we use the work of Bary-Soroker and Meisner [BSM19], who give explicitly the affine equations of general ℓ -cyclic covers. We summarize their results below.

As before, let n_q be the multiplicative order of q modulo ℓ . Then, the conductors of the ℓ -cyclic covers over \mathbb{F}_q are monic square-free polynomials in $\mathbb{F}_q[t]$ supported on n_q -divisible primes. In order to count all the ℓ -cyclic covers (or characters of order ℓ) with such conductors, let

$$\mathcal{F}_{q,\ell} := \{F \in \mathbb{F}_q[t] : F = P_1^{e_1} \cdots P_s^{e_s}, n_q \mid \deg P_i, 1 \leq e_i \leq \ell - 1\},$$

where the P_i are monic irreducible n_q -divisible polynomials in $\mathbb{F}_q[t]$.

Let ϕ_q be the Frobenius automorphism of \mathbb{F}_q . Then, ϕ_q acts on $f(t) \in \mathbb{F}_{q^{n_q}}[t]$ by acting on the coefficients, and we define

$$N_{n_q}(f) := f \phi_q(f) \phi_q^2(f) \cdots \phi_q^{n_q-1}(f) \in \mathbb{F}_q[t].$$

By hypothesis, each prime P_i in the factorization of $F \in \mathcal{F}_{q,\ell}$ splits as n_q primes in $\mathbb{F}_{q^{n_q}}[t]$, and we can write any $F \in \mathcal{F}_{q,\ell}$ as

$$(4.1) \quad F = \mathfrak{F}_1 \cdots \mathfrak{F}_{n_q}, \quad \mathfrak{F}_i \in \mathbb{F}_{q^{n_q}}[t], \quad \phi_q(\mathfrak{F}_i) = \mathfrak{F}_{i+1} \quad 1 \leq i \leq n_q - 1, \quad \phi_q(\mathfrak{F}_{n_q}) = \mathfrak{F}_1.$$

In other words, for $F \in \mathcal{F}_{q,\ell}$, $F = N_{n_q}(\mathfrak{F}_i)$ for any i . Since \mathfrak{F}_1 determines \mathfrak{F}_i for all i , it suffices to work with \mathfrak{F}_1 . We will write $\mathfrak{F}_1 \in \mathcal{F}_{q,\ell}^{(1)}$ when $F \in \mathcal{F}_{q,\ell}$. We have

$$(4.2) \quad \mathfrak{F}_1 = f_1 f_2^2 \cdots f_{\ell-1}^{\ell-1},$$

where the $f_i \in \mathbb{F}_{q^{n_q}}[t]$ are pairwise co-prime and square-free.

For any vector $\mathbf{v} = (v_1, \dots, v_{n_q}) \in \mathcal{V} = \{0, 1, 2, \dots, \ell - 1\}^{n_q}$, and any $F \in \mathcal{F}_{q,\ell}$ written as in (4.1), let $F_{\mathbf{v}} = \mathfrak{F}_1^{v_1} \cdots \mathfrak{F}_{n_q}^{v_{n_q}}$. For $0 \leq k \leq n_q - 1$, let $\mathbf{v}_k = (q^k, q^{k-1}, \dots, q^{k+1-n_q})$. Reducing modulo ℓ , we have $\mathbf{v}_k \in \mathcal{V}$. Let $\zeta_\ell \in \mathbb{F}_{q^{n_q}}$ be a fixed primitive ℓ th root of unity. For any $F \in \mathcal{F}_{q,\ell}$, let C_F be the curve over \mathbb{F}_q with affine model

$$(4.3) \quad C_F : \prod_{j=0}^{\ell-1} \left(y - \sum_{k=0}^{n_q-1} \zeta_\ell^{jq^k} \sqrt[\ell]{F_{\mathbf{v}_k}} \right) = 0.$$

Notice that there is no canonical choice for $\sqrt[\ell]{F_{\mathbf{v}_k}}$, but the above equation is still well defined, since the factors include all the Galois conjugates.

In the Kummer case $n_q = 1$, $F_{\mathbf{v}_0} = \mathfrak{F}_1 = F$, and C_F has affine model $y^\ell = F(t)$. In the case $\ell = 3$ and $q \equiv 2 \pmod{3}$, $F = \mathfrak{F}_1\mathfrak{F}_2$ and by (4.3), C_F has equation

$$\begin{aligned} C_F : & \left(y - \sqrt[3]{\mathfrak{F}_1\mathfrak{F}_2^2} - \sqrt[3]{\mathfrak{F}_1^2\mathfrak{F}_2} \right) \left(y - \zeta_3 \sqrt[3]{\mathfrak{F}_1\mathfrak{F}_2^2} - \zeta_3^2 \sqrt[3]{\mathfrak{F}_1^2\mathfrak{F}_2} \right) \\ & \times \left(y - \zeta_3^2 \sqrt[3]{\mathfrak{F}_1\mathfrak{F}_2^2} - \zeta_3 \sqrt[3]{\mathfrak{F}_1^2\mathfrak{F}_2} \right) = 0 \\ \iff & y^3 - 3\mathfrak{F}_1\mathfrak{F}_2 y - \mathfrak{F}_1\mathfrak{F}_2(\mathfrak{F}_1 + \mathfrak{F}_2) = 0, \end{aligned}$$

which is defined over \mathbb{F}_q . In general, C_F is birationally equivalent to $y^\ell = F_{\mathbf{v}_0}$ over $\overline{\mathbb{F}_q}$.

We now give more information about the equation (4.3) in the general case, including a precise formula for $n_q = 2$, using the work of Gupta and Zagier [GZ93]. We used these general formulas for $n_q = 2$ to obtain the equations for the curves C_1, C_2 and C_3 in Section 6.

Following [GZ93], we define the polynomial the complex polynomial

$$(4.4) \quad \Psi_{\ell, n_q}(y) = \prod_{j \in \mathcal{R}_{\ell, n_q}} \left(y - \sum_{k=0}^{n_q-1} \omega_\ell^{jq^k} \right),$$

where ℓ is an odd prime number, ω_ℓ denotes a complex ℓ -root of unity, and \mathcal{R}_{ℓ, n_q} denotes a set of coset representatives of $(\mathbb{Z}/\ell\mathbb{Z})^*$ modulo the cyclic subgroup $\langle q \rangle$. This is a polynomial of degree $\frac{\ell-1}{n_q}$. Notice that for $n_q = 1$, $\Psi_{\ell, 1}(y)$ gives the ℓ th cyclotomic polynomial and for $n_q = 2$, $\Psi_{\ell, 2}(y)$ gives the ℓ th real cyclotomic polynomial.

Gupta and Zagier prove various results regarding the coefficients of $\Psi_{\ell, n_q}(y)$, and in particular, they recover a formula of Gauss:

$$(4.5) \quad \Psi_{\ell, 2}(y) = \sum_{n=0}^{\frac{\ell-1}{2}} (-1)^{\lfloor \frac{\ell-1-2n}{4} \rfloor} \binom{\lfloor \frac{\ell-1+2n}{4} \rfloor}{n} y^n.$$

In the following result we relate the coefficients in the equation defining C_F in (4.3) to those of Ψ_{ℓ, n_q} . Together with the results of [GZ93], and (4.5) in particular, this allows us to compute a more explicit formula for equation (4.3) in the case $n_q = 2$.

Proposition 4.2. *Let ℓ be an odd prime and let $\Psi_{\ell, n_q}(y)$ be defined as in (4.4). Let a_m be the coefficients of the following polynomial*

$$(4.6) \quad y^\ell + \sum_{m=0}^{\ell-1} a_m y^m := \Psi_{\ell, n_q}(y)^{n_q} (y - n_q).$$

Then, $a_m \in \mathbb{Z}$, and there exists certain coefficients $b_{s_0, \dots, s_{n_q-1}} \in \mathbb{F}_p \subseteq \mathbb{F}_q$ such that the equation defining C_F in (4.3) can be written as

$$(4.7) \quad C_F : y^\ell + \sum_{m=0}^{\ell-1} \sum_{\substack{0 \leq s_k \\ \sum_{k=0}^{n_q-1} s_k = \ell - m \\ \sum_{k=0}^{n_q-1} q^k s_k \equiv 0 \pmod{\ell}}} b_{s_0, \dots, s_{n_q-1}} \mathfrak{F}_1^{\frac{1}{\ell} \sum_{k=0}^{n_q-1} s_k [q^k]_\ell} \mathfrak{F}_2^{\frac{1}{\ell} \sum_{k=0}^{n_q-1} s_k [q^{k-1}]_\ell} \dots \mathfrak{F}_{n_q}^{\frac{1}{\ell} \sum_{k=0}^{n_q-1} s_k [q^{k+1-n_q}]_\ell} y^m = 0,$$

where $[\alpha]_\ell \equiv \alpha \pmod{\ell}$, and $0 \leq [\alpha]_\ell \leq \ell - 1$. Furthermore, the $b_{s_0, \dots, s_{n_q-1}}$ satisfy

$$(4.8) \quad \sum_{\substack{0 \leq s_k \\ \sum_{k=0}^{n_q-1} s_k = \ell - m \\ \sum_{k=0}^{n_q-1} q^k s_k \equiv 0 \pmod{\ell}}} b_{s_0, \dots, s_{n_q-1}} = a_m,$$

where the a_m are given by (4.6) and the equality takes place in $\mathbb{F}_p \subseteq \mathbb{F}_q$ after reducing the a_m modulo p (the characteristic of \mathbb{F}_q).

In particular, for $n_q = 2$, we have

$$(4.9) \quad C_F : y^\ell + \sum_{r=1}^{\frac{\ell-1}{2}} a_{2r-1} (\mathfrak{F}_1 \mathfrak{F}_2)^{\frac{\ell+1}{2}-r} y^{2r-1} - \mathfrak{F}_1 \mathfrak{F}_2 (\mathfrak{F}_1^{\ell-2} + \mathfrak{F}_2^{\ell-2}) = 0.$$

Before proceeding to the proof, we remark that the condition $\sum_{k=0}^{n_q-1} q^k s_k \equiv 0 \pmod{\ell}$ implies that $\sum_{k=0}^{n_q-1} q^{k-j} s_k \equiv 0 \pmod{\ell}$ (since $(q, \ell) = 1$), and therefore each of the exponents of the \mathfrak{F}_j in (4.7) is an integer number. One can also see that the $b_{s_0, \dots, s_{n_q-1}}$ are invariant by cyclic permutation of the subindexes. Each of these cyclic permutations results in a permutation in the exponents of the \mathfrak{F}_j . Thus, the final polynomial is symmetric in the \mathfrak{F}_j .

Proof. The initial step of the proof follows from the elementary fact that

$$\Psi_{\ell, n_q}(y)^{n_q} (y - n_q) = \prod_{j=0}^{\ell-1} \left(y - \sum_{k=0}^{n_q-1} \omega_\ell^{jq^k} \right).$$

We remark that the above polynomial has coefficients in the algebraic integers $\overline{\mathbb{Z}}$, and that it is invariant by Galois conjugation of the coefficients. Therefore, we conclude that $\Psi_{\ell, n_q}(y)^{n_q} (y - n_q) \in \mathbb{Z}[y]$ and $a_m \in \mathbb{Z}$.

Following some ideas from [GZ93], we consider more generally

$$f_{\ell, n_q}(A_0, \dots, A_{n_q-1}) = \prod_{j=0}^{\ell-1} \left(1 - \sum_{k=0}^{n_q-1} \omega_\ell^{jq^k} A_k \right),$$

and we remark again that this polynomial has coefficients in \mathbb{Z} .

Taking the formal logarithm,

$$\begin{aligned}
-\log f_{\ell, n_q}(A_0, \dots, A_{n_q-1}) &= \sum_{j=0}^{\ell-1} \sum_{m=1}^{\infty} \frac{\left(\sum_{k=0}^{n_q-1} \omega_{\ell}^{jq^k} A_m \right)^m}{m} \\
&= \sum_{j=0}^{\ell-1} \sum_{m=1}^{\infty} \frac{1}{m} \sum_{\substack{h_0 + \dots + h_{n_q-1} = m \\ h_i \geq 0}} \binom{m}{h_0, \dots, h_{n_q-1}} \omega_{\ell}^{\sum_{k=0}^{n_q-1} jq^k h_k} A_0^{h_1} \dots A_{n_q-1}^{h_{n_q-1}} \\
&= \sum_{m=1}^{\infty} \frac{1}{m} \sum_{\substack{h_0 + \dots + h_{n_q-1} = m \\ h_i \geq 0}} \binom{m}{h_1, \dots, h_{n_q}} A_0^{h_0} \dots A_{n_q-1}^{h_{n_q-1}} \sum_{j=0}^{\ell-1} \omega_{\ell}^{j \sum_{k=0}^{n_q-1} q^k h_k}
\end{aligned}$$

and the innermost sum is zero unless $\sum_{k=0}^{n_q-1} q^k h_k \equiv 0 \pmod{\ell}$.

In conclusion, the only powers of A_0, \dots, A_{n_q-1} appearing in the Taylor series of $\log f_{\ell, n_q}(A_0, \dots, A_{n_q-1})$ and consequently in the Taylor series of $f_{\ell, n_q}(A_0, \dots, A_{n_q-1})$ are of the form $A_0^{s_0} \dots A_{n_q-1}^{s_{n_q-1}}$ such that

$$(4.10) \quad \sum_{k=0}^{n_q-1} q^k s_k \equiv 0 \pmod{\ell}.$$

But the total degree of f_{ℓ, n_q} is ℓ , and therefore $0 \leq s_0 + \dots + s_{n_q-1} \leq \ell$. Putting this information together, we obtain

$$(4.11) \quad f_{\ell, n_q}(A_0, \dots, A_{n_q-1}) = 1 + \sum_{m=0}^{\ell-1} \sum_{\substack{0 \leq s_k \\ \sum_{k=0}^{n_q-1} s_k = \ell - m \\ \sum_{k=0}^{n_q-1} q^k s_k \equiv 0 \pmod{\ell}}} b_{s_0, \dots, s_{n_q-1}} A_0^{s_0} \dots A_{n_q-1}^{s_{n_q-1}}.$$

Reducing modulo p (the characteristic of \mathbb{F}_q), making the change of variables

$$A_k = \frac{\sqrt{\ell F_{\mathbf{v}_k}}}{y} = \frac{1}{y} \mathfrak{F}_1^{\lfloor \frac{q^k}{\ell} \rfloor} \mathfrak{F}_2^{\lfloor \frac{q^{k-1}}{\ell} \rfloor} \dots \mathfrak{F}_{n_q}^{\lfloor \frac{q^{k+1-n_q}}{\ell} \rfloor},$$

and multiplying by y^{ℓ} , we obtain equation (4.7). Identity (4.8) follows from comparing with (4.6).

When $n_q = 2$, we have $q \equiv -1 \pmod{\ell}$. Equation (4.10) and condition $\sum_{k=0}^{n_q-1} s_k = \ell - m$ reduce the choices of s_0, s_1 to two cases: either $s_0 = s_1$ and $m \neq 0$ or $(s_0, s_1) = (0, \ell), (\ell, 0)$ and $m = 0$.

For the case $s_0 = s_1$, we can set $A_0 = A_1$ and reduce to the case of [GZ93, Theorem 3] to find the coefficients of each $(A_0 A_1)^{s_1}$. We then replace $A_0 = \frac{\sqrt{\ell \mathfrak{F}_1 \mathfrak{F}_2^{\ell-1}}}{y}$, $A_1 = \frac{\sqrt{\ell \mathfrak{F}_1^{\ell-1} \mathfrak{F}_2}}{y}$ (or equivalently, we replace $A_0 A_1$ by $\frac{\mathfrak{F}_1 \mathfrak{F}_2}{y}$), and obtain the coefficients a_m for $m \neq 0$ from the statement. In this case one can see from working with $\Psi_{\ell, 2}(y)$ that $a_m = 0$ for m even different from 0.

The cases $(s_0, s_1) = (0, \ell), (\ell, 0)$ only occur for the constant coefficient in (4.7) which is

$$(-1)^{\ell} \omega_{\ell}^{0 + \dots + (\ell-1)} (A_0^{\ell} + A_1^{\ell}) = -(A_0^{\ell} + A_1^{\ell}).$$

Replacing again $A_0 = \frac{\sqrt[\ell]{\mathfrak{F}_1 \mathfrak{F}_2^{\ell-1}}}{y}$, $A_1 = \frac{\sqrt[\ell]{\mathfrak{F}_1^{\ell-1} \mathfrak{F}_2}}{y}$ and multiplying by y^ℓ gives equation (4.9). \square

Proposition 4.3. [BSM19, Proposition 2.14] *Let $B = \{b \in \mathbb{F}_{q^{n_q}}^*/(\mathbb{F}_{q^{n_q}}^*)^\ell\}$. There is a $(\ell - 1)$ -to-1 correspondence between $\mathcal{F}_{q,\ell} \times B$ and the ℓ -cyclic covers of $\mathbb{F}_q(t)$, and then a 1-to-1 correspondence between $\mathcal{F}_{q,\ell} \times B$ and the characters of order ℓ over $\mathbb{F}_q(t)$.*

Notice that following Proposition 4.3, it will suffice for us to work with monic polynomials, and therefore we will restrict ourselves to work with the set $\mathcal{F}_{q,\ell}$.

Lemma 4.4. *With notation as above, assume $n_q > 1$. Then for each $0 \leq k \leq n_q - 1$, we have $\ell \mid \deg(F_{\mathbf{v}_k})$.*

Proof. By construction,

$$\begin{aligned} \deg(F_{\mathbf{v}_k}) &= \sum_{j=1}^{n_q} \mathbf{v}_{k,j} \deg(\mathfrak{F}_j) = \sum_{j=1}^{n_q} \mathbf{v}_{k,j} \deg(\phi^{j-1}(f_1 f_2^2 \cdots f_{\ell-1}^{\ell-1})) \\ &\equiv \sum_{j=1}^{n_q} q^{k+1-j} \sum_{h=1}^{\ell-1} h \deg(\phi^{j-1}(f_h)) \equiv \sum_{h=1}^{\ell-1} h \deg(f_h) \sum_{j=1}^{n_q} q^{k+1-j} \pmod{\ell}. \end{aligned}$$

Since $n_q > 1$,

$$\sum_{j=1}^{n_q} q^{k+1-j} = \frac{q^{k+1-n_q}(q^{n_q} - 1)}{q - 1} \equiv 0 \pmod{\ell}. \quad \square$$

Given an ℓ -cyclic cover C_0 , we can build ℓ -cyclic covers C with a non-constant map to C_0 by a change of variables, as done in [DL21, Lemma 3.2] for the Kummer case when $\ell \mid \deg F$. We can detect the curves C_F with $F \in \mathcal{F}_{q,\ell}$ using the following lemma.

Lemma 4.5. *Let $f \in \mathbb{F}_{q^{n_q}}[t]$. Then, $N_{n_q}(f)$ is square-free iff $f = \mathbf{p}_1 \cdots \mathbf{p}_s$ where the \mathbf{p}_i are such that $N_{n_q}(\mathbf{p}_i)$ are distinct n_q -divisible primes of $\mathbb{F}_q[t]$.*

Proof. If $f = \mathbf{p}_1 \cdots \mathbf{p}_s$, where the \mathbf{p}_i are such that $N_{n_q}(\mathbf{p}_i)$ are distinct n_q -divisible primes of $\mathbb{F}_q[t]$, then it is clear that $N_{n_q}(f) = N_{n_q}(\mathbf{p}_1) \cdots N_{n_q}(\mathbf{p}_s)$ is square-free.

Now assume that $N_{n_q}(f) = N_{n_q}(\mathbf{p}_1) \cdots N_{n_q}(\mathbf{p}_s)$ is square-free. Then it is clear that the $N_{n_q}(\mathbf{p}_i)$ are distinct primes in $\mathbb{F}_q[t]$. Finally, they are n_q -divisible, since they are the result of taking the N_{n_q} -norm. \square

Definition 4.6. *For a one-variable polynomial $f(t) \in \overline{\mathbb{F}_q}[t]$, let $f^*(u, v) := v^{\deg(f)} f(u/v)$ denote the homogeneous polynomial in variables u, v resulting from the change of variables $t = u/v$.*

Lemma 4.7. *Let $F \in \mathcal{F}_{q,\ell}$, with $\mathfrak{F}_1 \in \mathcal{F}_{q,\ell}^{(1)}$ given by (4.1) and C_F given by (4.3). As in (4.2), we write $\mathfrak{F}_1 = f_1 f_2^2 \cdots f_{\ell-1}^{\ell-1}$, where $f_i \in \mathbb{F}_{q^{n_q}}[t]$ are pairwise co-prime and square-free.*

- *Let $h(t)$ be a non-constant polynomial in $\mathbb{F}_q[t]$ such that*

$$N_{n_q}(f_1(h(t)) f_2(h(t)) \cdots f_{\ell-1}(h(t)))$$

is square-free. Then, $(F \circ h)(t) = N_{n_q}(\mathfrak{F}_1(h(t))) \in \mathcal{F}_{q,\ell}$. Let $C_{F \circ h}$ be given by (4.3). Then,

$$\begin{aligned} C_{F \circ h} &\longrightarrow C_F \\ (t, y) &\mapsto (h(t), y) \end{aligned}$$

is a non-constant map from $C_{F_{\text{oh}}}$ to C_F .

- Assume that $n_q > 1$. Let $u(t), v(t)$ be non-constant polynomials in $\mathbb{F}_q[t]$ such that

$$N_{n_q} (f_1^*(u, v) \cdots f_{\ell-1}^*(u, v))$$

is square-free. Then $G(t) = N_{n_q} (\mathfrak{F}_1^*(u(t), v(t))) \in \mathcal{F}_{q, \ell}$. Let C_G be given by (4.3). Then

$$\begin{aligned} C_G &\longrightarrow C_F \\ (t, y) &\mapsto (u(t)/v(t), yv(t)^{-\deg(F_{\mathbf{v}_0)}/\ell}) \end{aligned}$$

is a non-constant map from C_G to C_F .

- Assume that $n_q = 1$ and write $\deg F = A\ell - \delta$, where $0 \leq \delta \leq \ell - 1$. Let $u(t), v(t)$ be non-constant polynomials in $\mathbb{F}_q[t]$ such that $f_1^*(u, v)f_2^*(u, v) \cdots f_{\ell-1}^*(u, v)$ is square-free. Let $g_i^* = f_i^*$ for $i \neq \delta$ and $g_\delta^* = vf_\delta^*$. Then, $g_1^*(u, v)g_2^*(u, v) \cdots g_{\ell-1}^*(u, v)$ is also square-free and $G(t) = g_1^*(u, v)g_2^*(u, v)^2 \cdots g_{\ell-1}^*(u, v)^{\ell-1} \in \mathcal{F}_{q, \ell}$. Let $C_G : y^\ell = G(t)$. Then

$$\begin{aligned} C_G &\longrightarrow C_F \\ (t, y) &\mapsto (u(t)/v(t), yv(t)^{-A}) \end{aligned}$$

is a non-constant map from C_G to C_F .

Proof. We prove the second and third point in the statement, as the first point is a consequence of them. First consider the case where $n_q > 1$. We replace t by $u(t)/v(t)$ in equation (4.3) and we get

$$\prod_{j=0}^{\ell-1} \left(y - \sum_{k=0}^{n_q-1} \zeta_\ell^{jq^k} \sqrt[\ell]{\frac{F_{\mathbf{v}_k}^*(u, v)}{v^{\deg(F_{\mathbf{v}_k})}}} \right) = 0.$$

Recall from Lemma 4.4 that for the non-Kummer case, $\ell \mid \deg(F_{\mathbf{v}_k})$. Notice also that the \mathbf{v}_k are all permutations of each other. In fact, \mathbf{v}_{k+1} can be constructed from \mathbf{v}_k by shifting each element one place to the right cyclically and using the fact that $q^{n_q} \equiv 1 \pmod{\ell}$. Writing $A = \frac{\deg(F_{\mathbf{v}_k})}{\ell}$, and making the change of variables $Y = v^A y$, we finally have

$$\prod_{j=0}^{\ell-1} \left(Y - \sum_{k=0}^{n_q-1} \zeta_\ell^{jq^k} \sqrt[\ell]{F_{\mathbf{v}_k}^*(u, v)} \right) = 0,$$

which is C_G for $G(t) = N_{n_q} (\mathfrak{F}_1^*(u(t), v(t)))$.

We now consider the Kummer case. We replace t by $u(t)/v(t)$ in $y^\ell = F(t)$ to get

$$v^{A\ell} y^\ell = v^\delta F^*(u, v) = g_1^*(u, v)g_2^*(u, v)^2 \cdots g_{\ell-1}^*(u, v)^{\ell-1},$$

and with the change of variables $Y = v^A y$, we get

$$Y^\ell = g_1^*(u, v)g_2^*(u, v)^2 \cdots g_{\ell-1}^*(u, v)^{\ell-1},$$

which is C_G for $G(t) = g_1^*(u, v)g_2^*(u, v)^2 \cdots g_{\ell-1}^*(u, v)^{\ell-1}$. \square

Then Lemma 4.7 translates the conditions for finding curves C_G with a map to C_F to detecting when $N_{n_q} (f_1^*(u, v) \cdots f_{\ell-1}^*(u, v))$ is square-free. We can now proceed to the proof of Proposition 4.1.

Proof of Proposition 4.1. Our proof follows the argument of [DL21], but without restricting to the particular case where $n_q = 1$ and $\ell \mid \deg F$. We concentrate on the parts of their argument where using the general setting explained above introduces some changes, and we just refer to their article for the parts of their argument that can be directly used.

Let $F = F_0$ be as in Lemma 4.7 and let $C_0 = C_{F_0}$ be the curve (4.3). Let d_0 be the degree of the conductor. We now give a lower bound for the number of ℓ -cyclic covers with conductor of degree smaller than n that can be obtained by the process of Lemma 4.7 applied to F_0 , by using the square-free sieve over $\mathbb{F}_q[t]$.

Let

$$\begin{aligned} \mathcal{P}(n) &= \{(D_1, \dots, D_{\ell-1}) \in (\mathbb{F}_{q^{n_q}}[t])^{\ell-1} : D_1, \dots, D_{\ell-1} \text{ pairwise co-prime, monic, square-free,} \\ &\quad \mathfrak{F}_1 = D_1 \cdots D_{\ell-1}^{\ell-1} \in \mathcal{F}_{q,\ell}^{(4)}, \deg(D_1 \cdots D_{\ell-1}) \leq n\} \\ &= \{(D_1, \dots, D_{\ell-1}) \in (\mathbb{F}_{q^{n_q}}[t])^{\ell-1} : D_1, \dots, D_{\ell-1} \text{ monic, } N_{n_q}(D_1 \cdots D_{\ell-1}) \text{ square-free,} \\ &\quad \deg(D_1 \cdots D_{\ell-1}) \leq n\}, \end{aligned}$$

where the second line follows from Lemma 4.5.

By the above discussion, each tuple $(D_1, \dots, D_{\ell-1}) \in \mathcal{P}(n)$ gives rise to the ℓ -cyclic cover C_F where $\mathfrak{F}_1 = D_1 D_2^2 \cdots D_{\ell-1}^{\ell-1}$ and $F = N_{n_q}(\mathfrak{F}_1)$. The conductor is $N_{n_q}(D_1 \cdots D_{\ell-1})$ of degree $\leq n_q n$, and then the genus is such that $g \leq \frac{\ell-1}{2}(n_q n - 2)$.

We write $\mathfrak{F}_1^0 = f_1 f_2^2 \cdots f_{\ell-1}^{\ell-1}$ where $f_i \in \mathbb{F}_{q^{n_q}}[t]$ and $N_{n_q}(\mathfrak{F}_1^0) = F_0$. Notice that $d_0 = \deg(N_{n_q}(f_1 \cdots f_{\ell-1})) = n_q(\deg(f_1) + \cdots + \deg(f_{\ell-1}))$. We count the number of distinct $(D_1, \dots, D_{\ell-1}) \in \mathcal{P}(n)$ such that there exists $(u, v) \in \mathbb{F}_q[t]^2$ with

$$(4.12) \quad D_1(t) = f_1^*(u(t), v(t)), \dots, D_{\ell-1}(t) = f_{\ell-1}^*(u(t), v(t)).$$

We then need to detect when $N_{n_q}(D_1 \cdots D_{\ell-1})$ is square-free. Let $G(u, v)$ denote the homogeneous polynomial such that

$$N_{n_q}(f_1^*(u, v) \cdots f_{\ell-1}^*(u, v)) = G(u, v).$$

We now apply a result of Poonen [Poo03] which counts the number of square-free values of $G(u, v)$ as u, v runs over polynomials in $\mathbb{F}_q[t]$, as given in [DL21] in a form suitable for our application.

Proposition 4.8. [Poo03, Theorem 8.1] [DL21, Proposition 3.4] *Let P be a finite set of primes in $\mathbb{F}_q[t]$, B be the localization of $\mathbb{F}_q[t]$ by inverting the primes in P , $K = \mathbb{F}_q(t)$, $f \in B[x_1, \dots, x_m]$ be a polynomial that is square-free as an element of $K[x_1, \dots, x_m]$ and for a choice of $x \in \mathbb{F}_q[t]^m$, we say that $f(x)$ is square-free in B if the ideal $(f(x))$ is a product of distinct primes in B . For $b \in B$, define $|b| = |B/(b)|$ and for $b = (b_1, \dots, b_n) \in B^n$, define $|b| = \max |b_i|$. Let*

$$\begin{aligned} S_f &:= \{x \in \mathbb{F}_q[t]^m : f(x) \text{ is square-free in } B\}, \\ \mu_{S_f} &:= \lim_{N \rightarrow \infty} \frac{|\{b \in S_f : |b| < N\}|}{N^m}. \end{aligned}$$

For each nonzero prime π of B , let c_π be the number of $x \in (A/\pi^2)^m$ that satisfy $f(x) = 0$ in A/π^2 . The limit μ_{S_f} exists and is equal to $\prod_\pi (1 - c_\pi/|\pi|^{2m})$.

We then apply Proposition 4.8 to $G(u, v)$. Following [DL21, Remark 3.5], let B be the localization of $\mathbb{F}_q[t]$ by the set of primes π with $|\pi| \leq \deg(N_{n_q}(f_1 \cdots f_{\ell-1})) = d_0$. This guarantees that

$$\mu_{SG} = \lim_{N \rightarrow \infty} \frac{|\{b \in \mathbb{F}_q[t]^2, |b| \leq N : G(b) \text{ is square-free in } B\}|}{N^2} > 0.$$

The curve C_F associated to $F = N_{n_q}(D_1 D_2^2 \cdots D_{\ell-1}^{\ell-1}) = F_0^*(u(t), v(t))$ as in (4.12) has genus bounded by $\frac{\ell-1}{2}(d_0 \deg(u(t)/v(t)) - 2)$, and therefore, if we want to guarantee that the genus of C_F is less or equal than g , we can prescribe that

$$(4.13) \quad \deg(u(t)/v(t)) := \max\{\deg u(t), \deg v(t)\} \leq \frac{g + \ell - 1}{g_0 + \ell - 1},$$

where g_0 is the genus of C_{F_0} .

Now we want to give an upper bound for the $b = (u, v) \in \mathbb{F}_q[t]^2$ satisfying condition (4.13) such that equation (4.12) is satisfied. Now take $N = q^n$, with $n = \frac{2g}{\ell-1} + 2$, and we impose the condition $\max\{\deg u, \deg v\} \leq n/d_0$. Notice that

$$\frac{n}{d_0} = \frac{2g + 2(\ell - 1)}{d_0(\ell - 1)} = \frac{2g + 2(\ell - 1)}{(d_0 - 2)(\ell - 1) + 2(\ell - 1)} = \frac{g + \ell - 1}{g_0 + \ell - 1},$$

and therefore condition (4.13) is satisfied. Applying Proposition 4.8, we get a positive proportion of $\gg \mu N^{2/d_0} = \mu q^{2n/d_0}$ such that $N_{n_q}(D_1 \cdots D_{\ell-1})$ is square-free.

To conclude, for a fixed tuple $(D_1, \dots, D_{\ell-1})$ we need to find an upper bound on the number of pairs $(u(t), v(t))$ such that (4.12) is satisfied in order to correct a double counting. Following a similar reasoning to [DL21], we bound this number by $qn^2 q^{\varepsilon n}$.

In total, for n sufficiently large, we have

$$\gg \mu q^{n(2/d_0 - \varepsilon)}$$

elements in $\mathcal{P}(n)$ corresponding to ℓ -cyclic covers of $\mathbb{P}_{\mathbb{F}_q}^1$ with conductor of degree bounded by n that admit a non-constant map to C_0 . \square

We then need a geometric condition for the vanishing of $\mathcal{L}(C, u)$ at some point $u = u_0^{-1}$, where C is a curve over \mathbb{F}_q . This is given by the following theorem of Li [Li18, Section 2] relating the existence of a rational map between curves to the divisibility of the L -functions. The proof uses Honda–Tate theory, which states that every q -Weil number is an eigenvalue of the geometric Frobenius acting on the ℓ -adic Tate module of a simple abelian variety over \mathbb{F}_q , which is unique up to isogeny. We refer the reader to [Li18, Section 2] for the details, and the proof of the following theorem.

Theorem 4.9. *Let u_0 be a q -Weil number and let A_0 be (the isogeny class of) the unique simple Abelian variety over \mathbb{F}_q having u_0 as a Frobenius eigenvalue, as guaranteed by the theorem of Honda–Tate. Let C be a curve over \mathbb{F}_q . Then, $\mathcal{L}(C, u_0^{-1}) = 0$ if and only if there exists a non-trivial map $C \rightarrow A_0$ if and only if $\mathcal{L}(A_0, u)$ divides $\mathcal{L}(C, u)$.*

Proof of Theorems 1.1 and 1.2. The proof of Theorem 1.1 follows directly from Proposition 4.1 and Theorem 4.9: let C_0 be the ℓ -cyclic cover associated to χ_0 , i.e. $\mathcal{L}(C_0, u_0^{-1}) = 0$. By Proposition 4.1 and Theorem 4.9, there are at least q^{2n/d_0} ℓ -cyclic covers with conductor of degree $\leq n$ such that $\mathcal{L}(C_0, u) \mid \mathcal{L}(C, u) = \prod_{i=1}^{\ell-1} \mathcal{L}(\chi^i, u)$, and then at least q^{2n/d_0} characters of order ℓ and conductor of degree $\leq n$ such that $\mathcal{L}(\chi, u_0^{-1}) = 0$.

The proof of Theorem 1.2 follows directly from Corollary 3.3 and the above. Indeed, if $E = E_0 \times_{\mathbb{F}_q} \mathbb{F}_q(t)$ and there exists χ_0 such that $\mathcal{L}(E, \chi_0, q^{-1}) = 0$, then by Corollary 3.3, $\mathcal{L}(C_{\chi_0}, \alpha_1^{-1}) = 0$, and we reason as above. \square

5. DESCRIPTION OF THE CODE

We want to compute L -functions $\mathcal{L}(E, \chi, u)$ described by (2.6), where χ is a character of conductor F . To simplify, we are choosing $q = p$ to be prime.

Following Section 2, the L -functions are polynomials of degree $\mathbf{n} = \deg N_E + 2 \deg F - 4 + 2\delta_\chi$, and

$$\mathcal{L}(E, \chi, u) = \sum_{n=0}^{\mathbf{n}} \left(\sum_{f \in \mathcal{M}_n} a_f \chi(f) \right) u^n = \sum_{n=0}^{\mathbf{n}} c_n u^n,$$

where \mathcal{M}_n is the set of monic polynomials of degree n in $\mathbb{F}_p[t]$.

Using the functional equation (2.8), we get

$$(5.1) \quad c_n = \omega_{E \otimes \chi} p^{2(n - \lfloor \mathbf{n}/2 \rfloor - 1)} \overline{c_{\mathbf{n}-n}}, \quad 0 \leq n \leq \mathbf{n},$$

and it suffices to compute c_i for $0 \leq i \leq \lfloor \mathbf{n}/2 \rfloor$.⁵

We then need to compute the a_f appearing in (2.6), for $\deg f \leq \mathbf{n}/2$. It follows from the Euler product that $a_{fg} = a_f a_g$ for $(f, g) = 1$, and for $P \in \mathbb{F}_q[t]$ and $n \geq 1$,

$$a_{P^n} = \begin{cases} a_P a_{P^{n-1}} - p a_{P^{n-2}}, & \text{if } P \nmid N_E, \\ a_P a_{P^{n-1}}, & \text{if } P \mid N_E, \end{cases}$$

where p is the characteristic of \mathbb{F}_q .

We now turn to the computation of the a_P of a fixed curve $E : y^2 = x^3 + a(t)x^2 + b(t)x + c(t)$. For P prime, we compute a_P using

$$a_P = - \sum_{\substack{x \in \mathbb{F}_p[t] \\ \deg(x) < \deg(P)}} \left(\frac{x^3 + a(t)x^2 + b(t)x + c(t)}{P} \right).$$

After we have computed all a_f for $\deg f \leq (\deg N_E + 2d - 4 + 2\delta_\chi)/2$, we can evaluate $\mathcal{L}(E, \chi, u)$ for any Dirichlet character with conductor of degree d over $\mathbb{F}_p[t]$. We go through the characters of order ℓ and conductor degree d in the following way. Let n_p be the multiplicative order of p modulo ℓ as before. Let $F \in \mathbb{F}_p[t]$ be a polynomial of degree d supported on n_p -divisible primes. We can enumerate all characters of order ℓ and conductor F by choosing only one character per cyclic extension of order ℓ of $\mathbb{F}_q(t)$, since the L -functions of the $\ell - 1$ characters associated to the same extension K vanish together. Writing $F = P_1 \cdots P_k$, where the P_i are distinct n_p -divisible primes, and $P_i = \mathfrak{P}_{i,1} \cdots \mathfrak{P}_{i,n_p}$ over $\mathbb{F}_{p^{n_p}}(t)$, we consider

⁵It follows from (5.1) that we can compute numerically the sign of the functional equation by computing $c_{\mathbf{n}/2}$ when \mathbf{n} is even, and $c_{\lfloor \mathbf{n}/2 \rfloor}$ and $c_{\lfloor \mathbf{n}/2 \rfloor + 1}$ when \mathbf{n} is odd. We used this in the numerical data to compute twists of the Legendre curve by odd characters, as in this case Theorem 2.2 does not apply. Of course, this requires $c_{\mathbf{n}/2} \neq 0$. When $c_{\mathbf{n}/2} = 0$, we computed the next coefficient $c_{(\mathbf{n}/2)+1}$ to get the sign of the functional equation. In all the cases considered, $c_{(\mathbf{n}/2)+1}$ was not zero (when $c_{\mathbf{n}/2} = 0$), so this was enough.

the (non-conjugate) characters of conductor F over $\mathbb{F}_p(t)$ given by

$$(5.2) \quad \chi(A) = \chi_{\mathfrak{P}_{1,1}}(A) \prod_{j=2}^k \chi_{\mathfrak{P}_{j,1}}^{a_j}(A),$$

for $a_j \in \{1, \dots, \ell - 1\}$, and where each $\chi_{\mathfrak{P}_{j,1}}$ is the ℓ th-power residue symbol modulo $\mathfrak{P}_{j,1}$ over $\mathbb{F}_{p^{n_p}}(t)$ defined in Section 2.

6. VANISHING OF TWISTS OF CONSTANT CURVES: NUMERICAL DATA

Let E_0 be an elliptic curve over \mathbb{F}_p with $\mathcal{L}(E_0, u) = (1 - \alpha_0 u)(1 - \bar{\alpha}_0 u)$, and let $E = E_0 \times_{\mathbb{F}_p} \mathbb{F}_p(t)$. By (2.7), $\mathcal{L}(E, \chi, p^{-1}) = 0$ for some character χ associated to $K/\mathbb{F}_q(t)$ if and only if $\mathcal{L}(E/K, p^{-1}) = 0$, and using the results of Section 3, this is equivalent to

$$\mathcal{L}(E_0, u) \mid \mathcal{L}(C_\chi, u) = \prod_{j=1}^{\ell-1} \mathcal{L}(\chi^j, u).$$

By Theorem 1.2, once we have found one χ_0 such that $\mathcal{L}(C_{\chi_0}, \alpha_0^{-1}) = 0$, then there are infinitely many, so we concentrate on finding χ_0 . We examined degree 2 factors of $\mathcal{L}(\chi^j, u)$ which arise as $\mathcal{L}(E_0, u)$ for some E_0 over \mathbb{F}_p .

In particular, we considered the case where $\mathcal{L}(\chi, u)$ has degree 2, which in the case of even (respectively odd) characters means that the conductor of χ is a polynomial of degree 4 (respectively 3) in $\mathbb{F}_q[t]$. Table 1 presents results for this case: for fixed values of ℓ and p , we computed $\mathcal{L}(\chi, u)$ for all characters such that $\mathcal{L}(\chi, u)$ is a polynomial of degree 2, and we listed all the cases that we found where $\mathcal{L}(\chi, u) = \mathcal{L}(E_0, u)$ for some elliptic curve E_0/\mathbb{F}_p . Notice that this means $\mathcal{L}(C_\chi, u) = \mathcal{L}(E_0, u)^{\ell-1}$. Each entry in Table 1 may correspond to many characters χ . We did not count them, but our program keeps an instance for each case. For example, the curve C_1/\mathbb{F}_5 given by

$$y^3 + (2t^4 + 2t^3 + t^2 + 4t + 4)y + (3t^6 + 2t^5 + 2t^4 + 2t^3 + t^2 + t + 3) = 0$$

has L -function $\mathcal{L}(C_1, u) = (1 + 5u^2)^2$; the curve C_2/\mathbb{F}_{59} given by

$$y^5 + (54t^4 + 18t^3 + 34t^2 + 18t + 39)y^3 + (5t^8 + 23t^7 + 44t^6 + 20t^5 + 35t^4 + 30t^3 + 17t^2 + 33t + 21)y + (57t^{10} + 18t^9 + 24t^8 + 58t^7 + 14t^6 + 9t^5 + 41t^4 + 17t^3 + 38t^2 + 48t + 44) = 0$$

has L -function $\mathcal{L}(C_2, u) = (1 + 59u^2)^4$; and the curve C_3/\mathbb{F}_{13} given by

$$y^7 + (6t^4 + 6t^3 + 6t^2 + 12t + 1)y^5 + (t^8 + 2t^7 + 3t^6 + 6t^5 + t^4 + 5t + 4)y^3 + (6t^{12} + 5t^{11} + 10t^{10} + 7t^8 + 2t^7 + 3t^6 + 9t^5 + 3t^4 + 2t^3 + 6t^2 + t + 4)y + (11t^{14} + 6t^{13} + 12t^{12} + 10t^{11} + 5t^{10} + 8t^9 + 6t^8 + 2t^7 + 2t^6 + 10t^5 + 7t^4 + 12t^3 + 3t^2 + 3t + 9) = 0$$

has L -function $\mathcal{L}(C_3, u) = (1 + 13u^2)^6$.

Of course, it would be interesting to prove some criteria which guarantees the existence of a character of degree ℓ over \mathbb{F}_p such that $\mathcal{L}(E_0, u)$ divides $\mathcal{L}(\chi, u)$. From the data, we are led to believe that this could always be the case when $n_p = 2$ and $\mathcal{L}(E_0, u) = 1 + pu^2$, corresponding to the isogeny class of supersingular elliptic curves over \mathbb{F}_p , but we currently do not have a proof. We present further evidence for larger values of ℓ in Table 2. Since this becomes more time-consuming, we only consider a thin family of the characters of order ℓ , where $a_j = 1$ for all j in (5.2). In some cases $((\ell, p) = (13, 103), (17, 101), (31, 61),$ and

(37, 73)), we did not go over all characters in the thin family, we stopped after we found $\mathcal{L}(\chi, u) = (1 + pu^2)$, so there might be other characters where $\mathcal{L}(\chi, u) = (1 + a_p u + pu^2)$. In summary, the following is true for all the cases that we tested: for every ℓ, p such that $n_p = 2$, there exists a character χ of order ℓ over \mathbb{F}_p such that $\mathcal{L}(\chi, u) = 1 + pu^2$.

Remark 6.1. *There is a large amount of work in the literature on Newton polygons of cyclic covers of \mathbb{P}^1 , in particular on the existence of supersingular and superspecial curves. See for example, [LMPT19b, LMPT19a, LMPT20]. But the existence of the curves we present in this paper does not follow from previous work. In fact, the existence of supersingular curves in families of cyclic covers which ramify at 4 points with growing degree ℓ is surprising from a dimension counting perspective. More surprisingly, these curves are defined over the prime field \mathbb{F}_p .*

ℓ	p	n_p	$\mathcal{L}(\chi, u) = 1 + a_p u + pu^2$
3	5	2	0, 3
	7	1	-2, -1, 1, 2, 4
	11	2	-3, 0, 3, 6
	13	1	-5, -4, -2, -1, 1, 2, 4, 5
	17	2	-6, -3, 0, 3, 6
	19	1	-8, -7, -5, -4, -2, -1, 1, 2, 4, 5, 7, 8
5	3	4	\emptyset
	7	4	3
	11	1	-2, 2, 3
	13	4	-1, 4
	19	2	0, 5
	29	2	0
	31	1	-2, 2, 3, 8
7	13	2	0
	29	1	-2, 2, 5
11	23	1	\emptyset
	43	2	0
13	5	4	\emptyset
61	11	4	\emptyset

TABLE 1. All instances of E_0 for which there is a χ of order ℓ over \mathbb{F}_p such that $\mathcal{L}(\chi, u) = \mathcal{L}(E_0, u)$ for some elliptic curve E_0/\mathbb{F}_p .

ℓ	p	n_p	$\mathcal{L}(\chi, u) = 1 + a_p u + pu^2$
13	103	2	0
17	67	2	0
	101	2	0
19	37	2	0
31	61	2	0
37	73	2	0

TABLE 2. More cases where there is a character χ of order ℓ over \mathbb{F}_p such that $\mathcal{L}(\chi, u) = (1 + p^2u)$. For the cases $(\ell, p) = (17, 67)$ and $(19, 37)$, we considered all characters in the thin family, and we did not find any other cases where $\mathcal{L}(\chi, u) = \mathcal{L}(E_0, u)$ except for $\mathcal{L}(E_0, u) = (1 + p^2u)$. For the other cases, we stopped after finding χ such that $\mathcal{L}(\chi, u) = (1 + p^2u)$, and we did not find any other $\mathcal{L}(E_0, u)$ up to that point.

7. VANISHING OF TWISTS OF NON-CONSTANT CURVES: NUMERICAL DATA

We now present data for the vanishing of $\mathcal{L}(E, \chi, p^{-1})$, where χ varies over characters of order ℓ over the finite field \mathbb{F}_p , and where E is a non-constant curve. We used the Legendre curve $E_1 : y^2 = x(x-1)(x-t)$ of conductor $N_1 = t(t-1)P_\infty^2$, and the curve $E_2 : y^2 = (x-1)(x-2t^2-1)(x-t^2)$ of conductor $N_2 = t(t-1)(t+1)(t^2+1)$. Notice that the Legendre curve has bad reduction at P_∞ , while the curve E_2 has good reduction at P_∞ . Both E_1 and E_2 define rational elliptic surfaces over $\mathbb{C}(t)$, since the coefficients of their Weierstrass forms satisfy $\deg a_j(t) \leq j$. Moreover, E_1 has rank 0 over $\mathbb{C}(t)$, while E_2 has rank 1 over $\mathbb{C}(t)$ with generator

$$P = ((1+i)t^2 + (1+i)t + 1, (-1+i)t(t+1)(t-i)).$$

Then, E_2 has rank 1 over $\mathbb{F}_q(t)$ when \mathbb{F}_q contains a primitive fourth root of unity.

To prove that P is a generator of E_2 over $\mathbb{C}(t)$, one can follow the method of [LM19, Theorem 7], which uses the theory of Mordell–Weil lattices (see the book of Shioda and Schütt [SS19] for a general reference). Tate’s algorithm ([Sil94], IV.9) shows that the rational elliptic surface E_2 has singularities at $t = 0$ of type I_4 in the Kodaira classification, and at $t = \pm 1, \pm i$ of type I_2 . Then the Shioda–Tate formula ([Shi72, Corollary 1.5]) implies that the rank is 1, since the Picard number is known to be 10 for rational elliptic surfaces. One can prove that the torsion group over $\mathbb{C}(t)$ is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ by examining Table (4.5) in [MP89]. The point P is found by inspection. To prove that it is a generator, one predicts the absolute value of the discriminant of the Mordell–Weil lattice by using the Determinant formula ([SS19, Corollary 6.39]) and the fact that the Néron–Severi lattice has discriminant -1 for rational elliptic surfaces. Since the Mordell–Weil lattice has rank 1, the discriminant gives the Néron height of a generator, in this case $1/4$. We can then compute the height of P (coming from the height pairing on the Mordell–Weil lattice, [SS19, Definition 6.21]) and check that it is also $1/4$, showing that P is a generator.

We present in Tables 3, 4, and 5 our results for twists of the Legendre curve with characters of order 3, 5, and 7 respectively, and various ground fields $\mathbb{F}_p(t)$. For the curve given by $y^2 = (x-1)(x-2t^2-1)(x-t^2)$, we present in Tables 7, 8, and 9 our results for twists of this curve with characters of order 3, 5, and 7 respectively, and various ground fields $\mathbb{F}_p(t)$. We have also tested higher order twists ($\ell = 11, 13$ for E_1 and $\ell = 11, 31, 71$ for E_2) but without finding any vanishing. This data is presented in Tables 6 and 10.

Each table has the same format: the first three columns are the values of ℓ , p and n_p and the fourth column is the degree d of the conductors of the characters of order ℓ over $\mathbb{F}_p(t)$ considered (then, n_p always divides d). The L -functions $\mathcal{L}(E, \chi, u)$ are then computed for all χ of order ℓ over $\mathbb{F}_p(t)$ with conductor of degree d , and they are classified according to their analytic rank, which is defined as $\text{rank}(\chi) = r_{\text{an}}(E, \chi) = \text{ord}_{u=q^{-1}} \mathcal{L}(E, \chi, u)$. Since $\text{rank}(\chi^i) = \text{rank}(\chi^j)$, we only count one power per character in our data. Then, the next columns give the number of such χ where the analytic rank is 0, or 1, or 2, \dots . The most extensive computation that we did was for twists of order $\ell = 3$ of the curve E_2 for conductors of degree 8 over $\mathbb{F}_5(t)$, where we needed to compute a_p for primes of degree ≤ 8 , which is the most involved part of computing the twisted L -functions $\mathcal{L}(E_2, \chi, u)$ for characters with conductors of degree 8. This took approximately 20 days on an Intel(R) Core(TM) i5-4300U CPU. This is also the only case where we found a twist of analytic rank 3.

The data for the Legendre curve is very compatible with the conjectures of [DFK04] and [MR21], as we have found no instances of vanishing for any character of order 7 or higher. For the curve E_2 , we have found many instances of vanishing for characters of order 7, but none for characters of higher order.

twist order	p	n_p	deg conductor d	rank 0	rank 1	rank 2
3	5	2	2	6	4	0
			4	205	32	3
			6	5784	260	16
			8	302640	116	4
	7	1	1	5	0	0
			2	37	4	0
			3	324	37	1
			4	2935	73	0

TABLE 3. Twists of order 3 for the Legendre curve.

twist order	p	n_p	deg conductor d	rank 0	rank 1
5	7	4	4	585	3
	11	1	1	9	0
			2	199	0
			3	3759	5
			4	65143	11
	19	2	2	170	1

TABLE 4. Twists of order 5 for the Legendre curve.

twist order	p	n_p	deg conductor d	rank 0
7	5	6	6	2580
	11	3	3	440
	13	2	2	78
			4	25116
	23	3	3	4048
	29	1	1	27
			2	2512
			3	179192
	41	2	2	820
	197	1	1	195
	337	1	1	335
	379	1	1	377

TABLE 5. Twists of order 7 for the Legendre curve. We have found no instances of vanishing in this case.

twist order	p	n_p	deg conductor d	rank 0
11	5	5	5	624
	23	1	1	21
	43	2	2	903
	67	1	1	65
	89	1	1	87
13	5	4	4	150
	29	3	3	8120
	53	1	1	51
			2	16678

TABLE 6. Twists of order 11 and 13 for the Legendre curve. We have found no instances of vanishing in this case.

twist order	p	n_p	deg conductor d	rank 0	rank 1	rank 2	rank 3
3	5	2	2	8	2	0	0
			4	214	26	0	0
			6	5780	280	0	0
			8	149222	2136	20	2
	7	1	1	4	0	0	0
			2	30	2	0	0
			3	264	22	2	0
			4	2299	49	4	0
			5	18670	240	2	0
			6	148537	1343	32	0
	11	2	2	53	0	1	0
	13	1	1	8	0	0	0
			2	122	12	0	0
			3	2140	56	4	0
	17	2	2	116	20	0	0
	19	1	1	14	2	0	0
			2	380	28	2	0
	23	2	2	244	6	2	0
	29	2	2	364	42	0	0
	31	1	1	26	2	0	0
2			1190	24	6	0	
103	1	1	100	0	0	0	
109	1	1	104	0	0	0	
151	1	1	146	2	0	0	

TABLE 7. Twists of order 3 for the curve $y^2 = (x - 1)(x - 2t^2 - 1)(x - t^2)$.

twist order	p	n_p	deg conductor d	rank 0	rank 1	rank 2
5	7	4	4	587	0	1
	11	1	1	8	0	0
			2	166	0	0
			3	3064	0	0
	19	2	2	170	0	0
	29	2	2	388	18	0
	31	1	1	28	0	0
			2	1975	0	1
	41	1	1	36	0	0
	101	1	1	96	0	0
131	1	1	128	0	0	

TABLE 8. Twists of order 5 for the curve $y^2 = (x - 1)(x - 2t^2 - 1)(x - t^2)$.

twist order	p	n_p	deg conductor d	rank 0	rank 1
7	5	6	6	2560	20
	11	3	3	440	0
	13	2	2	72	6
			4	24984	132
	29	1	1	24	0
			2	2046	16
	41	2	2	800	20

TABLE 9. Twists of order 7 for the curve $y^2 = (x - 1)(x - 2t^2 - 1)(x - t^2)$.

twist order	p	n_p	deg conductor d	rank 0
11	5	5	5	624
	23	1	1	20
			2	2152
			3	168448
	43	2	2	902
	67	1	1	64
			2	22370
	89	1	1	84
199	1	1	196	
31	5	3	3	40
71	5	5	5	624

TABLE 10. Twists of order 11, 31, and 71 for the curve $y^2 = (x-1)(x-2t^2-1)(x-t^2)$. We have found no instances of vanishing in this case.

REFERENCES

- [BBC⁺04] Ben Brubaker, Alina Bucur, Gautam Chinta, Sharon Frechette, and Jeffrey Hoffstein, *Nonvanishing twists of $GL(2)$ automorphic L -functions*, Int. Math. Res. Not. (2004), no. 78, 4211–4239. MR 2111362
- [BFKRG20] Hung M. Bui, Alexandra Florea, Jonathan P. Keating, and Edva Roditty-Gershon, *Moments of quadratic twists of elliptic curve L -functions over function fields*, Algebra Number Theory **14** (2020), no. 7, 1853–1893. MR 4150252
- [BH12] Salman Baig and Chris Hall, *Experimental data for Goldfeld’s conjecture over function fields*, Exp. Math. **21** (2012), no. 4, 362–374. MR 3004252
- [BKR] Lea Beneish, Debanjana Kundu, and Anwesh Ray, *Rank jumps and growth of shafarevich–tate groups for elliptic curves in $\mathbb{Z}/p\mathbb{Z}$ -extensions*, arXiv:2107.09166.
- [Bru92] Armand Brumer, *The average rank of elliptic curves. I*, Invent. Math. **109** (1992), no. 3, 445–472. MR 1176198
- [BSM19] Lior Bary-Soroker and Patrick Meisner, *On the distribution of the rational points on cyclic covers in the absence of roots of unity*, Mathematika **65** (2019), no. 3, 719–742. MR 3940228
- [Cho] Peter J. Cho, *Analytic ranks of elliptic curves over number fields*, arXiv:2005.07909.
- [CL22] Antoine Comeau-Lapointe, *One-level density of the family of twists of an elliptic curve over function fields*, Journal of Number Theory (2022).
- [DFK04] Chantal David, Jack Fearnley, and Hershky Kisilevsky, *On the vanishing of twisted L -functions of elliptic curves*, Experiment. Math. **13** (2004), no. 2, 185–198. MR 2068892
- [DFK07] ———, *Vanishing of L -functions of elliptic curves over number fields*, Ranks of elliptic curves and random matrix theory, London Math. Soc. Lecture Note Ser., vol. 341, Cambridge Univ. Press, Cambridge, 2007, pp. 247–259. MR 2322350
- [DFL] Chantal David, Alexandra Florea, and Matilde Lalin, *The mean values of cubic L -functions over function fields*, arXiv e-prints, arXiv:1901.00817.

- [DL21] Ravi Donepudi and Wanlin Li, *Vanishing of Dirichlet L -functions at the central point over function fields*, Rocky Mountain J. Math. **51** (2021), no. 5, 1615–1628. MR 4382986
- [FKK12] Jack Fearnley, Hershy Kisilevsky, and Masato Kuwata, *Vanishing and non-vanishing Dirichlet twists of L -functions of elliptic curves*, J. Lond. Math. Soc. (2) **86** (2012), no. 2, 539–557. MR 2980924
- [For19] Michele Fornea, *Growth of the analytic rank of modular elliptic curves over quintic extensions*, Math. Res. Lett. **26** (2019), no. 6, 1571–1586. MR 4078688
- [GM91] F. Gouvêa and B. Mazur, *The square-free sieve and the rank of elliptic curves*, J. Amer. Math. Soc. **4** (1991), no. 1, 1–23. MR 1080648
- [Gol74] Dorian M. Goldfeld, *A simple proof of Siegel’s theorem*, Proc. Nat. Acad. Sci. U.S.A. **71** (1974), 1055. MR 0344222
- [GZ93] S. Gupta and D. Zagier, *On the coefficients of the minimal polynomials of Gaussian periods*, Math. Comp. **60** (1993), no. 201, 385–398. MR 1155574
- [Hay66] D. R. Hayes, *The expression of a polynomial as a sum of three irreducibles*, Acta Arith. **11** (1966), 461–488. MR 0201422
- [HB04] D. R. Heath-Brown, *The average analytic rank of elliptic curves*, Duke Math. J. **122** (2004), no. 3, 591–623. MR 2057019
- [Kat02] Nicholas M. Katz, *Twisted L -functions and monodromy*, Annals of Mathematics Studies, vol. 150, Princeton University Press, Princeton, NJ, 2002. MR 1875130
- [Li18] Wanlin Li, *Vanishing of hyperelliptic L -functions at the central point*, J. Number Theory **191** (2018), 85–103. MR 3825462
- [LM19] Matilde Lalín and Xinchun Ma, *θ -triangle and ω -parallelogram pairs with common area and common perimeter*, J. Number Theory **202** (2019), 1–26. MR 3958062
- [LMPT19a] Wanlin Li, Elena Mantovan, Rachel Pries, and Yunqing Tang, *Newton polygons arising from special families of cyclic covers of the projective line*, Res. Number Theory **5** (2019), no. 1, Paper No. 12, 31. MR 3897613
- [LMPT19b] ———, *Newton polygons of cyclic covers of the projective line branched at three points*, Research directions in number theory—Women in Numbers IV, Assoc. Women Math. Ser., vol. 19, Springer, Cham, [2019] ©2019, pp. 115–132. MR 4069381
- [LMPT20] ———, *Newton Polygon Stratification of the Torelli Locus in Unitary Shimura Varieties*, International Mathematics Research Notices (2020), rnaa306.
- [LOT21] Robert J. Lemke Oliver and Frank Thorne, *Rank growth of elliptic curves in non-abelian extensions*, Int. Math. Res. Not. IMRN (2021), no. 24, 18411–18441. MR 4365991
- [Mil68] J. S. Milne, *The Tate-Šafarevič group of a constant abelian variety*, Invent. Math. **6** (1968), 91–105. MR 244264
- [MP89] Rick Miranda and Ulf Persson, *Torsion groups of elliptic surfaces*, Compositio Math. **72** (1989), no. 3, 249–267. MR 1032336
- [MR] Barry Mazur and Karl Rubin, *Arithmetic conjectures suggested by the statistical behavior of modular symbols*, arXiv:1910.12798.
- [MR18] Barry Mazur and Karl Rubin, *Diophantine stability*, Amer. J. Math. **140** (2018), no. 3, 571–616, With an appendix by Michael Larsen. MR 3805014
- [MR21] Barry Mazur and Karl Rubin, *Arithmetic conjectures suggested by the statistical behavior of modular symbols*, Experimental Mathematics (2021).
- [MS] Patrick Meisner and Anders Södergren, *Low-lying zeros in families of elliptic curve L -functions over function fields*, arXiv:2110.00102.
- [Oes90] Joseph Oesterlé, *Empilements de sphères*, no. 189–190, 1990, Séminaire Bourbaki, Vol. 1989/90, pp. Exp. No. 727, 375–397. MR 1099882
- [Poo03] Bjorn Poonen, *Squarefree values of multivariable polynomials*, Duke Math. J. **118** (2003), no. 2, 353–373. MR 1980998
- [Shi72] Tetsuji Shioda, *On elliptic modular surfaces*, J. Math. Soc. Japan **24** (1972), 20–59. MR 0429918
- [Sil94] Joseph H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 151, Springer-Verlag, New York, 1994. MR 1312368

- [SS19] Matthias Schütt and Tetsuji Shioda, *Mordell-Weil lattices*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 70, Springer, Singapore, 2019. MR 3970314
- [Tan93] Ki-Seng Tan, *Modular elements over function fields*, J. Number Theory **45** (1993), no. 3, 295–311. MR 1247386
- [TR92] Ki-Seng Tan and Daniel Rockmore, *Computation of L-series for elliptic curves over function fields*, J. Reine Angew. Math. **424** (1992), 107–135. MR 1144162
- [Wei71] André Weil, *Dirichlet series and automorphic forms*, Lect. Notes. Math., vol. 189, Berlin–Heidelberg–New York, 1971.

ANTOINE COMEAU-LAPOINTE: DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY, 1455 DE MAISONNEUVE WEST, MONTRÉAL, QUÉBEC, CANADA H3G 1M8

Email address: antoine.comeau-lapointe@concordia.ca

CHANTAL DAVID: DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY, 1455 DE MAISONNEUVE WEST, MONTRÉAL, QUÉBEC, CANADA H3G 1M8

Email address: chantal.david@concordia.ca

MATILDE LALÍN: DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL. CP 6128, SUCC. CENTRE-VILLE. MONTREAL, QC H3C 3J7, CANADA

Email address: mlalin@dms.umontreal.ca

WANLIN LI: CENTRE DE RECHERCHES MATHÉMATIQUES, UNIVERSITÉ DE MONTRÉAL. CP 6128, SUCC. CENTRE-VILLE. MONTREAL, QC H3C 3J7, CANADA

Email address: liwanlin@crm.umontreal.ca